

# Reading notes



Linfan

This document contains all notes taken while reading materials (e.g., textbooks, literature) in preparation for the capstone project. Black text are information consolidated from the readings; blue text are notes (proofs, explanations); red text are questions.

## Contents

<b>1</b>	<b>Theory of Ordinary Differential Equations (Coddington Levinson)</b>	<b>2</b>
1.1	Chapter 11: Algebraic properties of linear boundary-value problems on a finite interval	2
1.1.1	Introduction	2
1.1.2	Boundary form formula	5

## Todo list

	Why the $(-1)^n$ ?	5
	Is the direction correct? e.g., from the existence of $J$ , construct $U^+$ and $U_c^+$ .	10

# 1 Theory of Ordinary Differential Equations (Coddington Levinson)

## 1.1 Chapter 11: Algebraic properties of linear boundary-value problems on a finite interval

### 1.1.1 Introduction

**Definition 1.1.1.** Let  $L$  be the **linear differential operator of order  $n$**  ( $n \geq 1$ ) defined by

$$Lx = p_0x^{(n)} + p_1x^{(n-1)} + \cdots + p_{n-1}x' + p_nx$$

where the  $p_k$  are complex-valued functions of class  $C^{n-k}$  on a closed bounded interval  $[a, b]$  (i.e., derivatives  $p_k, p'_k, \dots, p_k^{(n-k)}$  exist on  $[a, b]$  and are continuous) and  $p_0(t) \neq 0$  on  $[a, b]$ .

**Definition 1.1.2.** **Homogeneous boundary conditions** refer to a set of equations/constraints of the type

$$\sum_{k=1}^n (M_{jk}x^{(k-1)}(a) + N_{jk}x^{(k-1)}(b)) = 0 \quad (j = 1, \dots, m) \quad (1.1.1)$$

where  $M_{jk}, N_{jk}$  are complex constants.

**Definition 1.1.3.** A **homogeneous boundary-value problem** concerns finding the solutions of

$$Lx = 0$$

on  $[a, b]$  which satisfy some homogeneous boundary conditions defined above.

**Definition 1.1.4.** For any homogeneous boundary value problem, an **adjoint problem** refers to the problem of finding the solutions of

$$L^+x := (-1)^n(\bar{p}_0x)^{(n)} + (-1)^{n-1}(\bar{p}_1x)^{(n-1)} + \cdots + \bar{p}_nx = 0$$

on  $[a, b]$  which satisfy some homogeneous boundary conditions “complementary” to the conditions associated with the solutions of  $Lx = 0$ .

**Theorem 1.1.5.** (Green’s formula) For  $u, v \in C^n$  on  $[a, b]$ ,

$$\int_{t_1}^{t_2} (Lu)\bar{v} dt - \int_{t_1}^{t_2} u(L^+\bar{v}) dt = [uv](t_2) - [uv](t_1) \quad (1.1.2)$$

where  $a \leq t_1 < t_2 \leq b$  and  $[uv](t)$  is the form in  $(u, u', \dots, u^{(n-1)})$  and  $(v, v', \dots, v^{(n-1)})$  given by

$$[uv](t) = \sum_{m=1}^n \sum_{j+k=m-1} (-1)^j u^{(k)}(t) (p_{n-m}\bar{v})^{(j)}(t).$$

*Remark 1.1.6.* Alternatively,  $[uv](t)$  can be written as (checked for  $n = 2$ )

$$[uv](t) = \sum_{j,k=1}^n B_{jk}(t) u^{(k-1)}(t) \bar{v}^{(j-1)}(t) \quad (1.1.3)$$

where  $B_{jk}$  are the  $j, k$ -entry of the  $n \times n$  matrix

$$B(t) = \begin{bmatrix} B_{11} & B_{12} & \cdots & \cdots & p_0(t) \\ \vdots & \vdots & \cdots & -p_0(t) & 0 \\ (-1)^{n-1}p_0(t) & 0 & \cdots & 0 & 0. \end{bmatrix}$$

Since  $B(t)$  is square with  $\det B(t) = (p_0(t))^n$  where  $p_0(t) \neq 0$  on  $[a, b]$  (as in the definition of  $L$ ),  $B(t)$  is nonsingular/invertible for  $t \in [a, b]$ .

**Definition 1.1.7.** For vectors  $f = (f_1, \dots, f_k)$ ,  $g = (g_1, \dots, g_k)$ , define the product

$$f \cdot g := \sum_{i=1}^k f_i \bar{g}_i.$$

**Definition 1.1.8.** A **semibilinear form** is a complex-valued function  $\mathcal{S}$  defined for pairs of vectors  $f = (f_1, \dots, f_k)$ ,  $g = (g_1, \dots, g_k)$  satisfying

$$\begin{aligned}\mathcal{S}(\alpha f + \beta g, h) &= \alpha \mathcal{S}(f, h) + \beta \mathcal{S}(g, h) \\ \mathcal{S}(f, \alpha g + \beta h) &= \bar{\alpha} \mathcal{S}(f, g) + \bar{\beta} \mathcal{S}(f, h)\end{aligned}$$

for any complex numbers  $\alpha, \beta$  and vectors  $f, g, h$ .

Note that  $\mathcal{S}$  is linear in the first argument but not the second one. If  $\mathcal{S}$  were bilinear, it would be linear in each argument.

*Remark 1.1.9.* If

$$S = \begin{bmatrix} s_{11} & \cdots & s_{1k} \\ \vdots & & \vdots \\ s_{k1} & \cdots & s_{kk} \end{bmatrix},$$

then  $Sf \cdot g$  is a semibilinear form

$$\begin{aligned}\mathcal{S}(f, g) &:= Sf \cdot g \\ &= \begin{bmatrix} s_{11} & \cdots & s_{1k} \\ \vdots & & \vdots \\ s_{k1} & \cdots & s_{kk} \end{bmatrix} \begin{bmatrix} f_1 \\ \vdots \\ f_k \end{bmatrix} \cdot \begin{bmatrix} g_1 \\ \vdots \\ g_k \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=1}^k s_{1j} f_j \\ \vdots \\ \sum_{j=1}^k s_{kj} f_j \end{bmatrix} \cdot \begin{bmatrix} g_1 \\ \vdots \\ g_k \end{bmatrix} \\ &= \sum_{i=1}^k \left( \sum_{j=1}^k s_{ij} f_j \right) \bar{g}_i \\ &= \sum_{i,j=1}^k s_{ij} f_j \bar{g}_i.\end{aligned}\tag{1.1.4}$$

Indeed:

$$\begin{aligned}\mathcal{S}(\alpha f + \beta g, h) &= \sum_{i,j=1}^k s_{ij} (\alpha f_j + \beta g_j) \bar{h}_i \\ &= \alpha \sum_{i,j=1}^k s_{ij} f_j \bar{h}_i + \beta \sum_{i,j=1}^k s_{ij} g_j \bar{h}_i \\ &= \alpha Sf \cdot h + \beta Sg \cdot h \\ &= \alpha \mathcal{S}(f, h) + \beta \mathcal{S}(g, h).\end{aligned}$$

Similarly,

$$\mathcal{S}(f, \alpha g + \beta h) = \sum_{i,j=1}^k s_{ij} f_j (\alpha \bar{g}_i + \bar{\beta} \bar{h}_i)$$

$$\begin{aligned}
&= \bar{\alpha} \sum_{i,j=1}^k s_{ij} f_j \bar{g}_i + \bar{\beta} \sum_{i,j=1}^k f_j \bar{h}_i \\
&= \bar{\alpha} S f \cdot g + \bar{\beta} S f \cdot h \\
&= \bar{\alpha} \mathcal{S}(f, g) + \bar{\beta} \mathcal{S}(f, h).
\end{aligned}$$

*Remark 1.1.10.* Under a similar matrix framework, we see that  $[uv](t)$  is a semibilinear form with matrix  $B(t)$ :

Let  $\vec{u} = (u, u', \dots, u^{(n-1)})$  and  $\vec{v} = (v, v', \dots, v^{(n-1)})$ . Then we have

$$\begin{aligned}
[uv](t) &= \sum_{j,k=1}^n B_{jk}(t) u^{(k-1)}(t) \bar{v}^{(j-1)}(t) \quad (\text{by (1.1.3)}) \\
&= \sum_{i,j=1}^n (B_{ij} u^{(j-1)} \bar{v}^{(i-1)})(t) \\
&= (B \vec{u} \cdot \vec{v})(t) \\
&= \mathcal{S}(\vec{u}, \vec{v})(t).
\end{aligned}$$

With this notation, we can rewrite the right hand side of Green's formula as a semibilinear form below:

$$\begin{aligned}
[uv](t_2) - [uv](t_1) &= \sum_{j,k=1}^n B_{jk}(t_2) u^{(k-1)}(t_2) \bar{v}^{(j-1)}(t_2) - \sum_{j,k=1}^n B_{jk}(t_1) u^{(k-1)}(t_1) \bar{v}^{(j-1)}(t_1) \\
&= B(t_2) \vec{u}(t_2) \cdot \vec{v}(t_2) - B(t_1) \vec{u}(t_1) \cdot \vec{v}(t_1) \\
&= \begin{bmatrix} B_{11}(t_2) & \cdots & B_{1n}(t_2) \\ \vdots & & \vdots \\ B_{n1}(t_2) & \cdots & B_{nn}(t_2) \end{bmatrix} \begin{bmatrix} u(t_2) \\ \vdots \\ u^{(n-1)}(t_2) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_2) \\ \vdots \\ \bar{v}^{(n-1)}(t_2) \end{bmatrix} - \\
&\quad \begin{bmatrix} B_{11}(t_1) & \cdots & B_{1n}(t_1) \\ \vdots & & \vdots \\ B_{n1}(t_1) & \cdots & B_{nn}(t_1) \end{bmatrix} \begin{bmatrix} u(t_1) \\ \vdots \\ u^{(n-1)}(t_1) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_1) \\ \vdots \\ \bar{v}^{(n-1)}(t_1) \end{bmatrix} \\
&= \begin{bmatrix} -B_{11}(t_1) & \cdots & -B_{1n}(t_1) \\ \vdots & & \vdots \\ -B_{n1}(t_1) & \cdots & -B_{nn}(t_1) \end{bmatrix} \begin{bmatrix} u(t_1) \\ \vdots \\ u^{(n-1)}(t_1) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_1) \\ \vdots \\ \bar{v}^{(n-1)}(t_1) \end{bmatrix} + \\
&\quad \begin{bmatrix} B_{11}(t_2) & \cdots & B_{1n}(t_2) \\ \vdots & & \vdots \\ B_{n1}(t_2) & \cdots & B_{nn}(t_2) \end{bmatrix} \begin{bmatrix} u(t_2) \\ \vdots \\ u^{(n-1)}(t_2) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_2) \\ \vdots \\ \bar{v}^{(n-1)}(t_2) \end{bmatrix} \\
&= \begin{bmatrix} -B_{11}(t_1) & \cdots & -B_{1n}(t_1) & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ -B_{n1}(t_1) & \cdots & -B_{nn}(t_1) & 0 & \cdots & 0 \\ 0 & \cdots & 0 & B_{11}(t_2) & \cdots & B_{1n}(t_2) \\ \vdots & & \vdots & \vdots & & \vdots \\ 0 & \cdots & 0 & B_{n1}(t_2) & \cdots & B_{nn}(t_2) \end{bmatrix} \begin{bmatrix} u(t_1) \\ \vdots \\ u^{(n-1)}(t_1) \\ u(t_2) \\ \vdots \\ u^{(n-1)}(t_2) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_1) \\ \vdots \\ \bar{v}^{(n-1)}(t_1) \\ \bar{v}(t_2) \\ \vdots \\ \bar{v}^{(n-1)}(t_2) \end{bmatrix} \\
&= \begin{bmatrix} -B(t_1) & 0_n \\ 0_n & B(t_2) \end{bmatrix} \begin{bmatrix} u(t_1) \\ \vdots \\ u^{(n-1)}(t_1) \\ u(t_2) \\ \vdots \\ u^{(n-1)}(t_2) \end{bmatrix} \cdot \begin{bmatrix} \bar{v}(t_1) \\ \vdots \\ \bar{v}^{(n-1)}(t_1) \\ \bar{v}(t_2) \\ \vdots \\ \bar{v}^{(n-1)}(t_2) \end{bmatrix} \\
&=: \hat{B} \begin{bmatrix} \vec{u}(t_1) \\ \vec{u}(t_2) \end{bmatrix} \cdot \begin{bmatrix} \vec{v}(t_1) \\ \vec{v}(t_2) \end{bmatrix}.
\end{aligned} \tag{1.1.5}$$

Since  $\det \hat{B} = (-1)^n \det B(t_1) \det B(t_2)$ ,  $\hat{B}$  is nonsingular for  $t_1, t_2 \in [a, b]$  (since  $B(t)$  is nonsingular for  $t \in [a, b]$ , as shown before).

Why the  $(-1)^n$ ?

### 1.1.2 Boundary form formula

**Definition 1.1.11.** Given any set of  $2mn$  complex constants  $M_{ij}, N_{ij}$  ( $i = 1, \dots, m; j = 1, \dots, n$ ), define  $m$  **boundary operators (boundary forms)**  $U_1, \dots, U_m$  for functions  $x$  on  $[a, b]$ , for which  $x^{(j)}$  ( $j = 1, \dots, n-1$ ) exists at  $a$  and  $b$ , by

$$U_i x = \sum_{j=1}^n (M_{ij} x^{(j-1)}(a) + N_{ij} x^{(j-1)}(b)) \quad (i = 1, \dots, m) \tag{1.1.6}$$

$U_i$  are **linearly independent** if the only set of complex constants  $c_1, \dots, c_m$  for which

$$\sum_{i=1}^m c_i U_i x = 0$$

for all  $x \in C^{n-1}$  on  $[a, b]$  is  $c_1 = c_2 = \dots = c_m = 0$ .

*Remark 1.1.12.* Note that for  $\alpha, \beta \in \mathbb{C}$  and  $x_1, x_2 \in C^{n-1}$  on  $[a, b]$ ,

$$\begin{aligned} U_i(\alpha x_1 + \beta x_2) &= \sum_{j=1}^n (M_{ij}(\alpha x_1 + \beta x_2)^{(j-1)}(a) + N_{ij}(\alpha x_1 + \beta x_2)^{(j-1)}(b)) \\ &= \alpha \sum_{j=1}^n (M_{ij}x_1^{(j-1)}(a) + N_{ij}x_1^{(j-1)}(b)) + \beta \sum_{j=1}^n (M_{ij}x_2^{(j-1)}(a) + N_{ij}x_2^{(j-1)}(b)) \quad (\text{by linearity of derivatives}) \\ &= \alpha U_i x_1 + \beta U_i x_2. \end{aligned}$$

So  $U_i$  are linear operators.

*Remark 1.1.13.* To describe (1.1.6) with matrices, define

$$\xi := \begin{bmatrix} x \\ x' \\ \vdots \\ x^{(n-1)} \end{bmatrix}; \quad U := \begin{bmatrix} U_1 \\ U_2 \\ \vdots \\ U_m \end{bmatrix}; \quad M := \begin{bmatrix} M_{11} & \cdots & M_{1n} \\ \vdots & & \vdots \\ M_{m1} & \cdots & M_{mn} \end{bmatrix}; \quad N := \begin{bmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & & \vdots \\ N_{m1} & \cdots & N_{mn} \end{bmatrix}.$$

Then (1.1.6) can be written as

$$Ux = M\xi(a) + N\xi(b).$$

Indeed:

$$\begin{aligned} M\xi(a) + N\xi(b) &= \begin{bmatrix} M_{11} & \cdots & M_{1n} \\ \vdots & & \vdots \\ M_{m1} & \cdots & M_{mn} \end{bmatrix} \begin{bmatrix} x(a) \\ x'(a) \\ \vdots \\ x^{(n-1)}(a) \end{bmatrix} + \begin{bmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & & \vdots \\ N_{m1} & \cdots & N_{mn} \end{bmatrix} \begin{bmatrix} x(b) \\ x'(b) \\ \vdots \\ x^{(n-1)}(b) \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=1}^n M_{1j}x^{(j-1)}(a) \\ \vdots \\ \sum_{j=1}^n M_{mj}x^{(j-1)}(a) \end{bmatrix} + \begin{bmatrix} \sum_{j=1}^n N_{1j}x^{(j-1)}(b) \\ \vdots \\ \sum_{j=1}^n N_{mj}x^{(j-1)}(b) \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=1}^n (M_{1j}x^{(j-1)}(a) + N_{1j}x^{(j-1)}(b)) \\ \vdots \\ \sum_{j=1}^n (M_{mj}x^{(j-1)}(a) + N_{mj}x^{(j-1)}(b)) \end{bmatrix} \\ &= \begin{bmatrix} U_1 x \\ \vdots \\ U_m x \end{bmatrix} = \begin{bmatrix} U_1 \\ U_2 \\ \vdots \\ U_m \end{bmatrix} x = Ux. \end{aligned}$$

Define the  $m \times 2n$  matrix

$$(M : N) := \begin{bmatrix} M_{11} & \cdots & M_{1n} & N_{11} & \cdots & N_{1n} \\ \vdots & & \vdots & \vdots & & \vdots \\ M_{m1} & \cdots & M_{mn} & N_{m1} & \cdots & N_{mn} \end{bmatrix}.$$

Then  $U_1, \dots, U_m$  are linearly independent if and only if  $\text{rank}(M : N) = m$ , or equivalently,  $\text{rank}(U) = m$ . Recall that the rank of a matrix is the largest number of linearly independent rows or columns in it. For a matrix  $A_{m \times n}$ ,  $\text{rank}(A) \leq \min\{m, n\}$  and  $\text{rank}(A) = \text{rank}(A^T)$ .

$Ux$  can be written as

$$\begin{aligned} Ux &= \begin{bmatrix} \sum_{j=1}^n (M_{1j}x^{(j-1)}(a) + N_{1j}x^{(j-1)}(b)) \\ \vdots \\ \sum_{j=1}^n (M_{mj}x^{(j-1)}(a) + N_{mj}x^{(j-1)}(b)) \end{bmatrix} \\ &= \begin{bmatrix} M_{11} & \cdots & M_{1n} & N_{11} & \cdots & N_{1n} \\ \vdots & & \vdots & \vdots & & \vdots \\ M_{m1} & \cdots & M_{mn} & N_{m1} & \cdots & N_{mn} \end{bmatrix} \begin{bmatrix} x(a) \\ \vdots \\ x^{(n-1)}(a) \\ x(b) \\ \vdots \\ x^{(n-1)}(b) \end{bmatrix} \\ &= (M : N) \begin{bmatrix} \xi(a) \\ \xi(b) \end{bmatrix}. \end{aligned}$$

**Definition 1.1.14.** If  $U = (U_1, \dots, U_m)$  is any boundary form with  $\text{rank}(U) = m$  and  $U_c = (U_{m+1}, \dots, U_{2n})$  any form with  $\text{rank}(U_c) = 2n - m$  such that  $(U_1, \dots, U_{2n})$  has rank  $2n$ , then  $U$  and  $U_c$  are **complementary boundary forms**. “Adjoining”  $U_{m+1}, \dots, U_{2n}$  to  $U_1, \dots, U_m$  is equivalent to imbedding the matrix  $(M : N)$  in a  $2n \times 2n$  nonsingular matrix (recall that for square matrices, nonsingular  $\iff$  full rank).

We wish to describe the right hand side of Green’s formula (1.1.2) as a linear combination of a boundary form  $U$  and a complementary form  $U_c$ . To do so, we consider the following results about the semibilinear form (1.1.4).

**Definition 1.1.15.** For a matrix  $A = (a_{ij})$ , its **adjoint** is defined as the conjugate transpose  $A^* = (\bar{a}_{ij})$ .

**Proposition 1.1.16.** In the context of the semibilinear form (1.1.4), we have

$$Sf \cdot g = f \cdot S^*g. \quad (1.1.7)$$

*Proof.*

$$\begin{aligned} Sf \cdot g &= \sum_{i,j=1}^k s_{ij} f_j \bar{g}_i \quad (\text{by (1.1.4)}); \\ f \cdot S^*g &= \begin{bmatrix} f_1 \\ \vdots \\ f_k \end{bmatrix} \cdot \begin{bmatrix} \bar{s}_{11} & \cdots & \bar{s}_{k1} \\ \vdots & & \vdots \\ \bar{s}_{1k} & \cdots & \bar{s}_{kk} \end{bmatrix} \begin{bmatrix} g_1 \\ \vdots \\ g_k \end{bmatrix} \\ &= \begin{bmatrix} f_1 \\ \vdots \\ f_k \end{bmatrix} \cdot \begin{bmatrix} \sum_{j=1}^k \bar{s}_{j1} g_j \\ \vdots \\ \sum_{j=1}^k \bar{s}_{jk} g_j \end{bmatrix} \\ &= \sum_{i=1}^k f_i \cdot \left( \sum_{j=1}^k \bar{s}_{ji} g_j \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^k f_i \cdot \left( \sum_{j=1}^k s_{ji} \bar{g}_j \right) \\
&= \sum_{i,j=1}^k s_{ji} f_i \bar{g}_j = Sf \cdot g.
\end{aligned}$$

□

**Proposition 1.1.17.** *Let  $\mathcal{S}$  be the semibilinear form associated with a nonsingular matrix  $S$ . Suppose  $\bar{f} := Ff$  where  $F$  is a nonsingular matrix. Then there exists a unique nonsingular matrix  $G$  such that if  $\bar{g} = Gg$ , then  $\mathcal{S}(f, g) = \bar{f} \cdot \bar{g}$  for all  $f, g$ .*

*Proof.* Let  $G := (SF^{-1})^*$ , then

$$\begin{aligned}
\mathcal{S}(f, g) &= Sf \cdot g \\
&= S(F^{-1}F)f \cdot g \\
&= SF^{-1}(Ff) \cdot g \\
&= SF^{-1}\bar{f} \cdot g \\
&= \bar{f} \cdot (SF^{-1})^*g \quad (\text{by (1.1.7)}) \\
&= \bar{f} \cdot Gg \\
&= \bar{f} \cdot \bar{g}.
\end{aligned}$$

To see that  $G$  is nonsingular, note that  $\det G = \det((SF^{-1})^T) = \det(\overline{SF^{-1}}) = \overline{\det(SF^{-1})} = \overline{\det(S)\det(F)^{-1}} \neq 0$  since  $S, F$  are nonsingular. □

**Proposition 1.1.18.** *Suppose  $\mathcal{S}$  is associated with the unit matrix  $E$ , i.e.,  $\mathcal{S}(f, g) = f \cdot g$ . Let  $F$  be a nonsingular matrix such that the first  $j$  ( $1 \leq j < k$ ) components of  $\bar{f} = Ff$  are the same as those of  $f$ . Then the unique nonsingular matrix  $G$  such that  $\bar{g} = Gg$  and  $\bar{f} \cdot \bar{g} = f \cdot g$  (as in Proposition 1.1.17) is such that the last  $k - j$  components of  $\bar{g}$  are linear combinations of the last  $k - j$  components of  $g$  with nonsingular coefficient matrix.*

*Proof.* We note that for the condition on  $F$  to hold,  $F$  must have the form

$$\begin{bmatrix} E_j & 0_+ \\ F_+ & F_{k-j} \end{bmatrix}_{k \times k}$$

where  $E_j$  is the  $j \times j$  identity matrix,  $0_+$  is the  $j \times (k - j)$  zero matrix,  $F_+$  is a  $(k - j) \times j$  matrix, and  $F_{k-j}$  a  $(k - j) \times (k - j)$  matrix. Let  $G$  be the unique nonsingular matrix in Proposition 1.1.17. Write  $G$  as

$$\begin{bmatrix} G_j & G_- \\ G_+ & G_{k-j} \end{bmatrix}_{k \times k}$$

where  $G_j, G_-, G_+, G_{k-j}$  are  $j \times j, j \times (k - j), (k - j) \times j, (k - j) \times (k - j)$  matrices, respectively. By the definition of  $G$ ,

$$f \cdot g = Ff \cdot Gg = \bar{f} \cdot Gg = G^* \bar{f} \cdot g = G^* Ff \cdot g,$$

(where the third equality follows from a reverse application of (1.1.7) with  $\bar{f}$  as  $f$ ,  $G^*$  as  $S$ ) which implies

$$G^* F = E_k.$$



Since

$$\begin{aligned} G^*F &= \begin{bmatrix} G_j^* & G_-^* \\ G_-^* & G_{k-j}^* \end{bmatrix} \begin{bmatrix} E_j & 0_+ \\ F_+ & F_{k-j} \end{bmatrix} \\ &= \begin{bmatrix} G_j^* + G_-^*F_+ & G_-^*F_{k-j} \\ G_-^* + G_{k-j}^*F_+ & G_{k-j}^*F_{k-j} \end{bmatrix} \\ &= \begin{bmatrix} E_j & 0_{j \times (k-j)} \\ 0_{(k-j) \times j} & E_{k-j} \end{bmatrix}. \end{aligned}$$

Thus,  $G_-^*F_{k-j} = 0_+$ , the  $j \times (k-j)$  zero matrix. But  $\det F = \det(E_j) \cdot \det(F_{k-j}) \neq 0$ , so  $\det F_{k-j} \neq 0$  and we must have  $G_-^* = 0_+$ , i.e.,  $G_- = 0_{(k-j) \times j}$ . Thus,  $G$  is upper-triangular, and so  $\det G = \det G_j \cdot \det G_{k-j} \neq 0$ , which implies  $\det G_{k-j} \neq 0$  and  $G_{k-j}$  is nonsingular. Hence,

$$\bar{g} = Gg = \begin{bmatrix} G_j & G_- \\ 0_{(k-j) \times j} & G_{k-j} \end{bmatrix} \begin{bmatrix} g_1 \\ \vdots \\ g_k \end{bmatrix}$$

where  $G_{k-j}$  is the nonsingular coefficient matrix such that

$$\begin{bmatrix} \bar{g}_{j-1} \\ \vdots \\ \bar{g}_k \end{bmatrix} = G_{k-j} \begin{bmatrix} g_{j-1} \\ \vdots \\ g_k \end{bmatrix}.$$

□

**Theorem 1.1.19.** (Boundary-form formula) Given any boundary form  $U$  of rank  $m$  (Definition 1.1.11), and any complementary form  $U_c$  (Definition 1.1.14), there exist unique boundary forms  $U_c^+$ ,  $U^+$  of rank  $m$  and  $2n - m$ , respectively, such that

$$[xy](b) - [xy](a) = Ux \cdot U_c^+y + U_cx \cdot U^+y. \quad (1.1.8)$$

If  $\tilde{U}_c$  is any other complementary form to  $U$ , and  $\tilde{U}_c^+, \tilde{U}^+$  the corresponding forms of rank  $m$  and  $2n - m$ , then

$$\tilde{U}^+y = C^*U^+y$$

for some nonsingular matrix  $C$ .

*Proof.* Recall from (1.1.5) that the left hand side of (1.1.8) can be considered as a semibilinear form  $\mathcal{S}(f, g) = \hat{B}f \cdot g$  for vectors

$$f = \begin{bmatrix} x(a) \\ \vdots \\ x^{(n-1)}(a) \\ x(b) \\ \vdots \\ x^{(n-1)}(b) \end{bmatrix}, \quad g = \begin{bmatrix} y(a) \\ \vdots \\ y^{(n-1)}(a) \\ y(b) \\ \vdots \\ y^{(n-1)}(b) \end{bmatrix}$$

with the nonsingular matrix

$$\hat{B} = \begin{bmatrix} -B(a) & 0_n \\ 0_n & B(b) \end{bmatrix}.$$

Recall from Remark 1.1.13 that

$$Ux = M\xi(a) + N\xi(b) = (M : N) \begin{bmatrix} \xi(a) \\ \xi(b) \end{bmatrix}$$

for  $M, N, \xi$  are as defined there. With the definition of  $f$ , we have  $f = \begin{bmatrix} \xi(a) \\ \xi(b) \end{bmatrix}$  and thus

$$Ux = (M : N)f.$$

By Definition 1.1.14,  $U_c x = (\tilde{M} : \tilde{N})f$  for two appropriate matrices  $\tilde{M}, \tilde{N}$  for which

$$H = \begin{bmatrix} M & N \\ \tilde{M} & \tilde{N} \end{bmatrix}_{2n \times 2n}$$

has rank  $2n$ . Thus,

$$\begin{bmatrix} Ux \\ U_c x \end{bmatrix} = \begin{bmatrix} (M : N)f \\ (\tilde{M} : \tilde{N})f \end{bmatrix} = \begin{bmatrix} M & N \\ \tilde{M} & \tilde{N} \end{bmatrix} f = Hf.$$

By Proposition 1.1.17, there exists a unique  $2n \times 2n$  nonsingular matrix  $J$  such that  $\mathcal{S}(f, g) = Hf \cdot Jg$ . Let  $U^+, U_c^+$  be such that

$$Jg = \begin{bmatrix} U_c^+ y \\ U^+ y \end{bmatrix},$$

then (1.1.8) holds since

$$[xy](b) - [xy](a) = \mathcal{S}(f, g) = Hf \cdot Jg = \begin{bmatrix} Ux \\ U_c x \end{bmatrix} \cdot \begin{bmatrix} U_c^+ y \\ U^+ y \end{bmatrix} = Ux \cdot U_c^+ y + U_c x \cdot U^+ y.$$

□

Is the direction correct? e.g., from the existence of  $J$  construct  $U^+$  and  $U_c^+$ .