

Referring Physician:


DOB: Age: Gender

Ref#: Hosp#: Provider Group:

Date of Service: Date Received: Outpatient
Room: Bed: Date Reported:

UID: 868F9544-F5A4-43EC-9031-6F701E82E6FE
TCGA-FY-A40M-01A-PR

Redacted



FINAL SURGICAL PATHOLOGY REPORT

Diagnosis:

1C0-0-3

path: carcinoma, papillary
Follicular variant 9340/3

COCF: carcinoma, papillary,
thyroid 8260/3

Site: thyroid, NOS

7-3-12 ED C73.4

A. THYROID GLAND, TOTAL THYROIDECTOMY:

- PAPILLARY CARCINOMA, FOLLICULAR VARIANT, MULTIFOCAL

- Largest focus, left lobe, 1.7 cm in maximum dimension.

- Second focus, right lobe, 0.1 cm in greatest dimension.

- MARGIN STATUS: NEGATIVE.

- Tumor is 0.01 cm from the posterior margin of the left lobe.

- EXTRATHYROIDAL EXTENSION: NOT IDENTIFIED.

- NO PARATHYROID TISSUE IDENTIFIED.

- SEE PATHOLOGIC TUMOR STAGING SUMMARY BELOW.

B. CENTRAL NODE TISSUE, EXCISION:

- FIVE LYMPH NODES, NEGATIVE FOR METASTASIS (0/5).

PATHOLOGIC TUMOR STAGING SUMMARY:

Tumor type: Papillary thyroid carcinoma, follicular variant.

Primary tumor: pT1b (largest focus, 1.7 cm in greatest dimension,
limited to the thyroid).

Regional lymph nodes: pN0 (0/5 lymph nodes).

Distant metastasis: Not applicable.

Tumor stage: I.

Lymphovascular invasion: Not identified.

Margin status: R0 (0.01 cm from posterior margin of left lobe).

Per TSS, follicular variant < 99%

Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
HIPAA Discrepancy		
Prior Malignancy History		
Dual/Synchronous Primary Notes		
Case R (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials		
Date reviewed		

Tumor Staging Information

Data derived from current specimen. Staging in accordance with or modified from AJCC Cancer Staging Handbook, Ed, and CAP protocol,

Case #:

Printed:


Phone

Page 1

This report continues... (FINAL)

Patient Narr

Patient: [REDACTED]



Case #: [REDACTED]

Procedure:

Specimen type:

Specimen integrity:

Specimen size:

Total thyroidectomy with central compartment dissection.

Thyroid gland.

Intact.

Right lobe: 3.7 x 1.8 x 1.3 cm.

Left lobe: 3.7 x 2.1 x 1.5 cm.

Isthmus: 1.8 x 1.5 x 0.5 cm.

TUMOR FEATURES:

Tumor size:

Largest focus, greatest dimension, 1.7 cm; additional dimensions, 1.5 x 1.4 cm.

Second focus, greatest dimension, 0.1 cm.

Tumor focality:

Multifocal

Tumor laterality:

Both lobes.

Histologic type:

Papillary carcinoma, follicular variant.

Lymphovascular invasion:

Not identified.

Tumor capsule:

Partially encapsulated.

Tumor capsular invasion:

Not identified.

Extrathyroidal extension:

Not identified.

LYMPH NODES:

Five lymph nodes, negative for metastasis (0/5).

MARGIN EVALUATION:

Distance to closest margin:

Negative (R0).

0.01 cm from posterior margin, left lobe.

Other margins:

0.06 cm from anterior left lobe margin; all other margins at least 0.1 cm.

PATHOLOGIC TUMOR STAGING DESCRIPTORS:

Primary tumor:

pT1b (largest lesion, 1.7 cm in greatest dimension, limited to the thyroid).

Regional lymph nodes:

pN0 (0/5 lymph nodes).

Distant metastasis:

Not applicable.

Margin status:

R0.

Pathologic stage:

I.

Additional pathologic findings:

Multiple adenomatoid nodules (multinodular goiter) with Hurthle cell change.

Comment:

Not applicable.

Case #: [REDACTED]

Printed: [REDACTED]

Page 2

This report continues... (FINAL)

Patient:

Case #:

FINAL SURGICAL PATHOLOGY REPORT

Source of Specimen:

- A. Thyroid
- B. Lymph node; Central node

Clinical History/Operative Dx:

Thyroid cancer

Gross Description:

A. The specimen is labeled total thyroid and is received without fixative. It consists of a total thyroidectomy specimen weighing 14 grams. A suture marks the left lobe. The right lobe measures 3.7 x 1.8 x 1.3 cm and the left lobe measures 3.7 x 2.1 x 1.5 cm. There is a triangular portion of isthmus which measures 1.8 x 1.5 x 0.5 cm. There are no grossly obvious parathyroid glands. The anterior right thyroid is inked blue, the anterior left thyroid is inked green, the anterior isthmus is inked yellow, and the posterior surfaces are inked black. The thyroid lobes and isthmus are serially sectioned to reveal a sharply circumscribed tan-white neoplasm in the superior third of the left thyroid lobe. It measures 1.7 x 1.5 x 1.4 cm. It appears contained within the thyroid gland. Representative sections of the neoplasm are obtained for research purposes. Elsewhere within the left thyroid lobe and also in the right thyroid lobe there is a beefy dark red thyroid gland parenchyma and areas of colloid-like change, but no other discrete masses. The specimen is entirely submitted as summarized below. Section summary:
A1-A6) left thyroid lobe submitted from superior-inferior,
A7-A11) right thyroid lobe submitted from superior-inferior,
A12-A13) isthmus.

B. The specimen is labeled central node and is received in formalin. It consists of multiple fragments of fibrofatty tissue which are 2 cm in aggregate. On gentle palpation there appears to be a possible 0.3 cm pale red lymph node. The specimen is submitted intact in cassette B1.

Microscopic Description:

A. Microscopic sections have been examined. The microscopic findings are reflected in the diagnosis rendered.

B. Microscopic sections have been examined. The microscopic findings are reflected in the diagnosis rendered.

Case #:

Printed:

Page 3

END OF REPORT (FINAL)