


Collection Date:



# **FINAL DIAGNOSIS:**

## **PART 1: THYROID, TOTAL THYROIDECTOMY (39 GRAMS) -**

- A. PAPILLARY THYROID CARCINOMA, THREE (3) FOCI (1.0 CM, 0.2 CM AND 0.1 CM), LEFT LOBE, NO ANGIOLYMPHATIC INVASION.
- B. LARGEST FOCUS, CONVENTIONAL TYPE WITH MICROSCOPIC EXTRATHYROIDAL EXTENSION; MARGINS FREE (CLOSEST 0.1 CM).
- C. PATHOLOGIC STAGE: pT3 N0.
- D. NODULAR THYROID HYPERPLASIA.

## **PART 2: LYMPH NODES, CENTRAL COMPARTMENT, SELECTIVE DISSECTION -**

- A. TWO (2) LYMPH NODES, NO TUMOR PRESENT (0/2).
- B. NORMOCELLULAR PARATHYROID.
- C. TISSUE ENTIRELY SUBMITTED FOR HISTOLOGIC EVALUATION.

## **CASE SYNOPSIS:**

### **SYNOPTIC DATA - PRIMARY THYROID TUMORS**

|                                 |                                                        |
|---------------------------------|--------------------------------------------------------|
| SPECIMEN TYPE:                  | Total Thyroidectomy                                    |
| TUMOR SITE:                     | Left Lobe                                              |
| TUMOR FOCALITY:                 | Multifocal                                             |
| TUMOR SIZE (largest nodule):    | Greatest Dimension: 1.0 cm                             |
| HISTOLOGIC TYPE**:              | Papillary carcinoma                                    |
| PRIMARY TUMOR (pT):             | pT3                                                    |
| REGIONAL LYMPH NODES (pN):      | pN0                                                    |
|                                 | Number of regional lymph nodes examined: 2             |
|                                 | Number of regional lymph nodes involved: 0             |
| DISTANT METASTASIS (pM):        | Not applicable                                         |
| EXTRATHYROIDAL EXTENSION:       | Present                                                |
|                                 | Extent: Minimal                                        |
| MARGINS:                        | Margins uninvolved by carcinoma                        |
|                                 | Distance of invasive carcinoma to closest margin: 1 mm |
| LYMPH-VASCULAR INVASION:        | Not identified                                         |
| ADDITIONAL PATHOLOGIC FINDINGS: | Nodular goiter                                         |

ICD-O-3

Carcinoma, papillary, thyroid 8260/3

Site: thyroid, NOS C73.9

pw  
9/29/12

| Criteria                            | Yes                                 | No                                  |
|-------------------------------------|-------------------------------------|-------------------------------------|
| Diagnosis Discrepancy               |                                     | <input checked="" type="checkbox"/> |
| Primary Tumor Site Discrepancy      | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> |
| HRPA Discrepancy                    |                                     | <input checked="" type="checkbox"/> |
| Prior Malignancy History            |                                     | <input checked="" type="checkbox"/> |
| Dual/Synchronous Primary Malignancy |                                     | <input checked="" type="checkbox"/> |
| Case is (circle):                   | QUALIFIED                           | DISQUALIFIED                        |
| Reviewer Initials                   | Date Reviewed: 9/29/12              |                                     |