


UUID: 28FED855-C6EB-479B-A359-28E4139D045A  
TCGA-EM-A2QW-01A-PR

Redacted



## Surgical Pathology Consultation Report

Patient

Name:

MRN:

DOB:

Gender: F

Service:

Visit #:

Location:

Facility:

Accession #:

Collected:

Received:

Reported:

Ordering

MD:

Copy To:

| Criteria                       | Yes       | No                                  |
|--------------------------------|-----------|-------------------------------------|
| Diagnosis Discrepancy          |           | <input checked="" type="checkbox"/> |
| Primary Tumor Site Discrepancy |           | <input checked="" type="checkbox"/> |
| HPA Discrepancy                |           | <input checked="" type="checkbox"/> |
| Prior Malignancy History       |           | <input checked="" type="checkbox"/> |
| Dual/Synchronous Primary Notes |           | <input checked="" type="checkbox"/> |
| Case Review                    | QUALIFIED | DISQUALIFIED                        |
| Reviewed                       | 8/31/11   | 8/18/11                             |

### Specimen(s) Received

1. Thyroid: RT.hemithyroid, stitch - rt. upper lobe

### Diagnosis

Multifocal papillary carcinoma, dominant oncocytic mixed follicular and macrofollicular variant, 1.3 cm; chronic lymphocytic thyroiditis and mild follicular nodular disease: Thyroid

No pathological diagnosis: Parathyroid, perithyroidal

- Right hemithyroidectomy specimen

ICD-O-3

carcinoma, papillary, follicular variant

8340/3

Site: thyroid, NOS C73.9

### Synoptic Data

Procedure: Right hemithyroidectomy

Received: Fresh

Specimen Integrity: Intact

Specimen Size: Right lobe:

5.9 cm

2.4 cm

1.4 cm

Isthmus +/- pyramidal lobe:

1.6 cm

0.8 cm

0.5 cm

Specimen Weight: 10.2 g

Tumor Focality: Multifocal, ipsilateral

----- DOMINANT TUMOR -----

Tumor Laterality: Right lobe

Tumor Size: Greatest dimension: 1.3cm

Additional dimension: 1.2 cm

Additional dimension: 1.0 cm

Histologic Type: Papillary carcinoma, follicular variant

Papillary carcinoma, macrofollicular variant

per ISS, follicular variant = 100%

|                                 |                                                                                                 |
|---------------------------------|-------------------------------------------------------------------------------------------------|
|                                 | Papillary carcinoma, oncocytic or oxyphilic variant                                             |
|                                 | Follicular architecture                                                                         |
|                                 | Macrofollicular architecture                                                                    |
|                                 | Classical cytomorphology                                                                        |
|                                 | Oncocytic or oxyphilic cytomorphology                                                           |
| Histologic Grade:               | Not applicable                                                                                  |
| Margins:                        | Margins uninvolved by carcinoma                                                                 |
| Tumor Capsule:                  | Totally encapsulated                                                                            |
| Tumor Capsular Invasion:        | Not identified                                                                                  |
| Lymph-Vascular Invasion:        | Not identified                                                                                  |
| Perineural Invasion:            | Not identified                                                                                  |
| Extrathyroidal Extension:       | Not identified                                                                                  |
| ----- SECOND TUMOR -----        |                                                                                                 |
| Tumor Laterality:               | Right lobe                                                                                      |
| Tumor Size:                     | Greatest dimension: 0.4cm                                                                       |
|                                 | Additional dimensions: 0.3 cm                                                                   |
| Histologic Type:                | Papillary carcinoma, follicular variant                                                         |
|                                 | Papillary carcinoma, microcarcinoma (occult, small or microscopic) variant                      |
|                                 | Papillary carcinoma, oncocytic or oxyphilic variant                                             |
|                                 | Follicular architecture                                                                         |
|                                 | Classical cytomorphology                                                                        |
|                                 | Oncocytic or oxyphilic cytomorphology                                                           |
| Histologic Grade:               | Not applicable                                                                                  |
| Margins:                        | Margins uninvolved by carcinoma                                                                 |
| Tumor Capsule:                  | None                                                                                            |
| Tumor Capsular Invasion:        | Not identified                                                                                  |
| Lymph-Vascular Invasion:        | Not identified                                                                                  |
| Perineural Invasion:            | Not identified                                                                                  |
| Extrathyroidal Extension:       | Not identified                                                                                  |
| TNM Descriptors:                | m (multiple primary tumors)                                                                     |
| Primary Tumor (pT):             | pT1b: Tumor more than 1 cm but not more than 2 cm in greatest dimension, limited to the thyroid |
| Regional Lymph Nodes (pN):      | pNX: Regional lymph nodes cannot be assigned                                                    |
|                                 | Number of regional lymph nodes examined: 0                                                      |
|                                 | Number of regional lymph nodes involved: 0                                                      |
| Distant Metastasis (pM):        | Not applicable                                                                                  |
| Additional Pathologic Findings: | Adenomatoid nodule(s) or Nodular follicular disease                                             |
|                                 | Thyroiditis                                                                                     |
|                                 | Parathyroid gland(s), within normal limits                                                      |

\*Pathologic Staging is based on AJCC/UICC TNM, 7th Edition

verified by:

Electronically

---

### **Gross Description**

1. The specimen labeled with the patient's name and as "Right Hemi thyroid stitch right upper lobe", consists of a right hemithyroidectomy specimen that is oriented with a suture and weighs 10.2 g. The lobe measures 5.9 cm SI x 2.4 cm ML x 1.4 cm AP and the isthmus measures 1.6 SI

x 0.8 cm ML x 0.5 cm AP. The external surfaces have fibrous adhesions and are painted with silver nitrate. No parathyroid glands or no lymph nodes are identified grossly. The mid right lobe contains one well delineated nodule that measures 1.3 cm SI x 1.2 cm ML x 1.0 cm AP. The cut surface of the nodule has a variegated brown tan color, solid and firm. A second nodule is identified on the inferior pole measuring 0.4 SI x 0.3 ML x 0.3 cm AP. The remainder of the thyroid tissue is brown in color homogeneous and grossly unremarkable. Sections of nodule and normal tissue are stored frozen. The remainder of the specimen is submitted as follows:

1A-J lobe, superior to inferior

1K isthmus

---