

LABORATORY MEDICINE PROGRAM

UUID: 59293B22-6D8E-4DC9-ABED-B4831E4FB34B
TCGA-EM-A3FP-01A-PR

Redacted



Surgical Pathology Consultation Report

Patient Name:

MRN:

DOB:

Gender:

HCN:

Ordering MD:

Copy To:

Service:

Visit #:

Location:

Facility:

Accession #:

Collected:

Received:

Reported:

Specimen(s) Received

1. Parathyroid: Right upper parathyroid QS
2. Parathyroid: Right inferior parathyroid/qs
3. Thyroid: Subtotal thyroid stitch marks upper pole right lobe

ICD-0-3

carcinoma, papillary, follicular
variant 8340/3

Site: thyroid, NOS C73.9 lw 11/22/11

Diagnosis

1. No pathological diagnosis: Parathyroid (right upper) biopsy
2. No pathological diagnosis: Parathyroid (right inferior) biopsy
3. Multifocal papillary carcinoma, dominant follicular variant, 2.2 cm, right; mild follicular nodular disease and chronic lymphocytic thyroiditis: Thyroid
No pathological diagnosis: Lymph nodes, 2, right perithyroidal
No pathological diagnosis: Lymph node, 1, isthmus
- Right hemithyroidectomy specimen

Synoptic Data

Procedure: Right hemithyroidectomy
Received: Fresh
Specimen Integrity: Intact
Specimen Size: Right lobe:
4.0 cm
3.8 cm
1.5 cm
Isthmus +/- pyramidal lobe:
4.5 cm
1.9 cm
1.0 cm
Specimen Weight: 19.3 g
Tumor Focality: Multifocal, ipsilateral
Multifocal, midline (isthmus)
----- DOMINANT TUMOR -----
Tumor Laterality: Right lobe
Tumor Size: Greatest dimension: 2.2cm
Additional dimension: 1.7 cm
Additional dimension: 1.6 cm
Histologic Type: Papillary carcinoma, follicular variant

Criteria	Yes	No
Diagnosis Discrepancy		X
Primary Tumor Site Discrepancy		X
HIPAA Discrepancy		X
Prior Malignancy History		X
Dual/Synchronous Primary Notes		X
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewed (Initials)	KMH lw 11/22/11 V12/11	
Date Reviewed		

Surgical Pathology Consultation Report

	Follicular architecture
	Classical cytomorphology
Histologic Grade:	Not applicable
Margins:	Margins uninvolved by carcinoma
Tumor Capsule:	Partially encapsulated
Tumor Capsular Invasion:	Present, extent minimal
Lymph-Vascular Invasion:	Not identified
Perineural Invasion:	Not identified
Extrathyroidal Extension:	Not identified
----- SECOND TUMOR -----	
Tumor Laterality:	Isthmus
Tumor Size:	Greatest dimension: 0.4cm
Histologic Type:	Papillary carcinoma, follicular variant
	Papillary carcinoma, microcarcinoma (occult, small or microscopic) variant
	Papillary carcinoma, oncocytic or oxyphilic variant
	Follicular architecture
	Classical cytomorphology
	Oncocytic or oxyphilic cytomorphology
Histologic Grade:	Not applicable
Margins:	Margins uninvolved by carcinoma
Tumor Capsule:	None
Tumor Capsular Invasion:	Not identified
Lymph-Vascular Invasion:	Not identified
Perineural Invasion:	Not identified
Extrathyroidal Extension:	Not identified
TNM Descriptors:	m (multiple primary tumors)
Primary Tumor (pT):	pT2: Tumor more than 2 cm, but not more than 4 cm, limited to thyroid
Regional Lymph Nodes (pN):	pN0: No regional lymph node metastasis
	Number of regional lymph nodes examined: 3
	Number of regional lymph nodes involved: 0
Distant Metastasis (pM):	Not applicable
Additional Pathologic Findings:	Adenomatoid nodule(s) or Nodular follicular disease
	Thyroiditis
	Parathyroid gland(s), within normal limits
	Other: additional 0.05 cm oncocytic follicular variant papillary microcarcinoma, right lobe

*Pathologic Staging is based on AJCC/UICC TNM

Electronically verified by:

Gross Description

1. The specimen labeled with the patient's name as "RT upper parathyroid QS" and consists of a fragment of tissue that measures 0.2 x 0.2 x 0.1 cm. The specimen is submitted in toto for frozen section.

1A resubmitted frozen tissue

2. The specimen labeled with the patient's name as "RT inferior parathyroid QS" and consists of a fragment of tissue that measures 0.2 x 0.2 x 0.1 cm. The specimen is submitted in toto for frozen section.

2A resubmitted frozen tissue

3. The specimen labeled with the patient's name and as "Subtotal thyroid stitch marks upper pole RT lobe" consists of a right hemithyroidectomy specimen that is received oriented with a suture and weighs 19.3 g. The right lobe measures 4.0 cm SI x 3.8 cm ML x 1.5 cm AP and the isthmus measures 4.5 cm SI x 1.9 cm ML x 1.0 cm AP. The external surfaces

Surgical Pathology Consultation Report

have fibrous adhesions and are painted with silver nitrate. No parathyroid glands or no lymph nodes are identified grossly. The right lobe contains 1 delineated, fleshy, yellow-pink nodule that measures 2.2 cm SI x 1.6 cm ML x 1.7 cm AP. The remainder of the thyroid tissue is red-brown with several other poorly defined nodules many of which have a translucent red appearance. Sections of the nodule and normal tissue are stored frozen. The remainder of the specimen is submitted as follows:

3A-J right lobe, superior to inferior (nodule in C-I)

3K-M isthmus

Quick Section Diagnosis

1A (right upper parathyroid) – parathyroid tissue present
Reported to OR13 a

2A (right inferior parathyroid) – parathyroid tissue present
Reported to OR13 at