

Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HIPAA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Noted		☒
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	RB	Date Reviewed: 6/12/12

UUID: 88D047C4-3144-4627-A7E8-2035D5C14852
 TCGA-EL-A3ZN-01A-PR
 Redacted


Surgical Pathology Report

Collected: _____ Received: _____ Case type: Surgical Case
 DOB: _____ Physician: _____ Accession: _____

DIAGNOSIS

(A) CONFIRM PARATHYROID:
Parathyroid tissue present.

ICD-O-3
 CQCF: carcinoma, papillary, thyroid 8260/3
 Path: carcinoma, papillary, follicular variant 8340/3
 Site: thyroid, NOS
 C73.9

(B) THYROID AND LEVEL 6 CONTENTS:
 PAPILLARY THYROID CARCINOMA, FOLLICULAR VARIANT
 Location: Left lobe
 Multi-focal: No
 Size = 1.3 cm
 Extrathyroidal extension: Present, extending into fibroadipose tissue
 Lymphovascular invasion: Present
 Resection Margins: Negative
 Lymph nodes:
 METASTATIC PAPILLARY THYROID CARCINOMA IN 2 OF 13 LYMPH NODES
 Location: Central, neck
 Extracapsular extension: Absent

hu
 6/17/12

(C) LEFT LEVELS 2A, 3, 4 AND 5:
 METASTATIC PAPILLARY THYROID CARCINOMA IN 4 OF 42 LYMPH NODES
 Extracapsular extension: Absent

GROSS DESCRIPTION

(A) CONFIRM PARATHYROID - A tan soft tissue fragment (0.2 x 0.1 x 0.1 cm). A touch prep is made and specimen is entirely submitted for frozen in A.
 *FS/DX: PARATHYROID GLAND.
 (B) THYROID AND LEVEL 6 CONTENTS - A total thyroidectomy consisting of right lobe (5.2 x 2.4 x 2.2 cm) and left lobe (4.8 x 2.2 x 1.6 cm) also in the container is a cystic hemorrhagic nodule (0.6 cm x diameter). The right lobe is unremarkable. The isthmus is unremarkable. The superior portion of the left lobe has a soft pink-tan multinodular focus (1.3 x 1.1 x 0.8 cm) which abuts the edge of the thyroid and is covered by loosely adherent soft tissue over the anterior surface. The remainder of the left lobe is unremarkable.
 SECTION CODE: B1-B3, right lobe representative from superior to inferior; B4, isthmus; B5-B7, superior left lobe with nodule; B8-B10, additional left lobe; B11, cystic nodule bisected; B12-B14, multiple lymph nodes from central adipose tissue in each cassette.
 (C) LEFT LEVELS 2A, 3, 4 AND 5 - Fibroadipose tissue (7.5 x 5.5 x 1.0 cm). Multiple lymph nodes are identified ranging from (0.1 x 0.1 x 0.1 to 1.5 x 1.0 x 0.7 cm).
 SECTION CODE: C1, one lymph node from level 2A; C2, one lymph node from level 3; C3-C5, lymph node from level 4 (C3, five lymph nodes; C4, four lymph nodes; C5, one lymph node, serially sectioned); C6-C13, lymph node from level 5 (C6-C12, each containing four lymph nodes; C13, two lymph nodes)

CLINICAL HISTORY

Surgical Pathology Report

DOB
Physician:

Collected:
Received:

Pathologist:
Case type: Surgical Case

Accession

Papillary thyroid carcinoma.

SNOMED CODES

T-C4200, M-80506, T-B6000, M-83403, T-C4200, M-80506,

"Some tests reported here may have been developed and performance characteristics determined by
approved by the U.S. Food and Drug Administration."

These tests have not been specifically cleared or

Entire report and diagnosis completed by:

Page 2 of 2

Surgical Pathology Report

File under: Pathology