

Collection Date:



# FINAL DIAGNOSIS:

## PART 1: THYROID, TOTAL THYROIDECTOMY (29 GRAMS) -

- A. PAPILLARY THYROID CARCINOMA, TWO FOCI (2.3 CM AND 0.4 CM), CONFINED TO THYROID.
- B. LARGEST FOCUS, LEFT LOBE (2.3 CM), ENCAPSULATED FOLLICULAR VARIANT WITH NO CAPSULAR OR ANGIOLYMPHATIC INVASION.
- C. PATHOLOGIC STAGE: pT2N0.
- D. ISTHMIC FOCUS (0.4 CM), MICROCARCINOMA WITH CYSTIC CHANGE, NO ANGIOLYMPHATIC INVASION (see comment).
- E. PATHOLOGIC STAGE: pT1aN0.
- F. THREE LYMPH NODES (ONE LEFT AND ONE RIGHT PERITHYROIDAL AND ONE ISTHMIC), NO TUMOR PRESENT (0/3).
- G. NODULAR THYROID HYPERPLASIA WITH CHRONIC LYMPHOCYTIC THYROIDITIS.

≥ 99% follicular variant per BS

## PART 2: PARATHYROID, 1/2 LEFT SUPERIOR, EXCISION (153 MG) - ENLARGED HYPERCELLULAR PARATHYROID.

## PART 3: PARATHYROID, 1/3 LEFT INFERIOR, EXCISION (14 MG) - SLIGHTLY ENLARGED MILDLY HYPERCELLULAR PARATHYROID.

## PART 4: LYMPH NODES, CENTRAL COMPARTMENT, SELECTIVE DISSECTION -

- A. FIVE LYMPH NODES, NO TUMOR PRESENT (0/5).
- B. THYROID TISSUE WITH CHRONIC LYMPHOCYTIC THYROIDITIS.
- C. SPECIMEN ENTIRELY SUBMITTED FOR HISTOLOGIC EVALUATION.

## PART 5: PARATHYROID, 1/4 RIGHT SUPERIOR, EXCISION (18 MG) - NORMOCYLLULAR PARATHYROID TISSUE.

ICD-0-3

carcinoma, papillary, follicular variant

8340/3

Site: thyroid, NOS C73.9

in 9/29/12

## COMMENT:

PART 1: The carcinoma is subcapsular, but shows no fibrosis. While technically there are two foci, these are different papillary carcinoma subtypes, and it can be argued that there is no intraglandular spread with respect to this microcarcinoma, which is cystic and of conventional histologic subtype.

PARTS 2, 3, AND 5: The intraoperative parathyroid hormone level decreased from 178 pg/ml to 12 pg/ml.

## CASE SYNOPSIS:

### SYNOPTIC DATA - PRIMARY THYROID TUMORS

SPECIMEN TYPE: Total Thyroidectomy  
 TUMOR SITE: Left Lobe  
 TUMOR FOCALITY: Multifocal  
 TUMOR SIZE (largest nodule): Greatest Dimension: 2.3 cm  
 HISTOLOGIC TYPE: Papillary carcinoma, follicular variant, non-encapsulated  
 TNM DESCRIPTORS: m  
 PRIMARY TUMOR (pT): pT2  
 REGIONAL LYMPH NODES (pN): pN0

Number of regional lymph nodes examined: 8  
 Number of regional lymph nodes involved: 0

DISTANT METASTASIS (pM): Not applicable  
 EXTRATHYROIDAL EXTENSION: Not identified  
 MARGINS: Margins uninvolved by carcinoma  
 LYMPH-VASCULAR INVASION: Not identified  
 ADDITIONAL PATHOLOGIC FINDINGS: Thyroiditis

### SYNOPTIC DATA - PRIMARY THYROID TUMORS

SPECIMEN TYPE: Total Thyroidectomy  
 TUMOR SITE: Isthmus  
 TUMOR FOCALITY: Multifocal  
 TUMOR SIZE (largest nodule): Greatest Dimension: 0.4 cm  
 HISTOLOGIC TYPE: Papillary carcinoma  
 TNM DESCRIPTORS: m  
 PRIMARY TUMOR (pT): pT1a  
 REGIONAL LYMPH NODES (pN): pN0

Number of regional lymph nodes examined: 8  
 Number of regional lymph nodes involved: 0

EXTRANODAL EXTENSION: Not identified  
 DISTANT METASTASIS (pM): Not applicable  
 EXTRATHYROIDAL EXTENSION: Not identified  
 MARGINS: Margins uninvolved by carcinoma  
 LYMPH-VASCULAR INVASION: Not identified  
 ADDITIONAL PATHOLOGIC FINDINGS: Thyroiditis

| Criteria                       | Yes                                 | No                                  |
|--------------------------------|-------------------------------------|-------------------------------------|
| Diagnosis Discrepancy          |                                     | <input checked="" type="checkbox"/> |
| Primary Tumor Site Discrepancy |                                     | <input checked="" type="checkbox"/> |
| HIPAA Discrepancy              |                                     | <input checked="" type="checkbox"/> |
| Prior Malignancy History       |                                     | <input checked="" type="checkbox"/> |
| Dual/Synchronous Primary Noted | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> |
| Case is (circle):              | QUALIFIED                           | DISQUALIFIED                        |
| Reviewer Initials: <i>BSH</i>  | Date Reviewed: <i>8/29/2012</i>     |                                     |

in 8/29/12