

LABORATORY MEDICINE PROGRAM

UUID: 44788DFC-8228-4228-A9DA-66AD8126C438

TCGA-EM-A309-01A-PR

Redacted



Surgical Pathology Consultation Report

Patient Name:

MRN:

DOB:

Gender:

HCN:

Ordering MD:

Copy To:

Service:

Visit #:

Location:

Facility:

Accession #

Collected:

Received:

Reported:

Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
IPAA Discrepancy		
Prior Malignancy History		
Just/Synchronous Primary Noted		
See it (circle):	QUALIFIED	DISQUALIFIED
Review (circle):		
Date Reviewed:	2/5/12	

Specimen(s) Received

1. Parathyroid: Right Lower
2. Thyroid ;Right hemithyroid stitch marks upper pole

Pw TSS on 2/20/13, TCGA tumor is
100% follicular variant. hu

Diagnosis

1. No pathological diagnosis: Parathyroid, right lower, biopsy.
2. Multifocal papillary carcinoma, dominant oncocytic follicular variant 4.5 cm, right; focal mild reactive C-cell hyperplasia: Thyroid
No pathological diagnosis: Lymph nodes, 2, isthmus
-Right hemithyroidectomy specimen. See Comment.

Comment

2. Although the clinical significance of focal mild C-cell hyperplasia is not well known in a hemithyroidectomy specimen, the features suggest a reactive phenomenon in this case. This case has been seen in consultation with Dr. [redacted] and he agrees with the interpretation and diagnosis.

Synoptic Data

ICD-0-3
CQEF: carcinoma, papillary, follicular variant 8340/3

8340/3

Path: carcinoma, papillary, follicular variant, and
oxyphilic 8342/3 - code to highest (8342/3)

Size: thyroid, NOS C73.9

hu
2/5/12

Procedure:

Received:

Specimen Integrity:

Specimen Size:

Right hemithyroidectomy

Fresh

Intact

Right lobe:

4.5 cm

3.2 cm

2.5 cm

Isthmus +/- pyramidal lobe:

2.9 cm

2.3 cm

0.8 cm

Specimen Weight:

27.7 g

Tumor Focality:

Multifocal, ipsilateral

Multifocal, midline (isthmus)

DOMINANT TUMOR

Tumor Laterality:

Right lobe

Tumor Size:

Greatest dimension: 4.5cm

Additional dimension: 3.2 cm

Additional dimension: 2.5 cm

Surgical Pathology Consultation Report

Histologic Type: Papillary carcinoma, follicular variant
Papillary carcinoma, oncocytic or oxyphilic variant
Follicular architecture
Oncocytic or oxyphilic cytomorphology

Histologic Grade: Not applicable

Margins: Margins uninvolved by carcinoma

Tumor Capsule: Totally encapsulated

Tumor Capsular Invasion: Present, extent minimal

Lymph-Vascular Invasion: Not identified

Perineural Invasion: Not identified

Extrathyroidal Extension: Not identified

----- SECOND TUMOR -----

Tumor Laterality: Isthmus

Tumor Size: Greatest dimension: 0.05cm

Histologic Type: Papillary carcinoma, follicular variant
Papillary carcinoma, microcarcinoma (occult, small or microscopic) variant
Follicular architecture
Classical cytomorphology

Histologic Grade: Not applicable

Margins: Margins uninvolved by carcinoma

Tumor Capsule: None

Tumor Capsular Invasion: Not identified

Lymph-Vascular Invasion: Not identified

Perineural Invasion: Not identified

Extrathyroidal Extension: Not identified

TNM Descriptors: m (multiple primary tumors)

Primary Tumor (pT): pT3: Tumor more than 4 cm, limited to thyroid or with minimal extrathyroidal extension
(eg, extension to sternothyroid muscle or perithyroid soft tissues)

Regional Lymph Nodes (pN): pN0: No regional lymph node metastasis
Number of regional lymph nodes examined: 2
Number of regional lymph nodes involved: 0

Distant Metastasis (pM): Not applicable

Additional Pathologic Findings: Thyroiditis, palpation
C-cell hyperplasia

Ancillary Studies: Type: IHC
Results: Calcitonin stains focal mild reactive C-cell hyperplasia.

*Pathologic Staging is based on AJCC/UICC TNM, 7th Edition

Electronically verified by:

Gross Description

1. The specimen labeled with the patient's name and as "RT Lower Parathyroid" contains a piece of tissue that measures 0.4 x 0.3 x 0.1 cm. It is frozen for intraoperative consultation.

1A frozen tissue resubmitted

2. The specimen labeled with the patient's name and as "RT Hemithyroid, Stitch Marks Upper Pole", consists of a right hemithyroidectomy specimen that is oriented with a suture and weighs 27.7 g. The lobe measures 4.5 cm SI x 3.2 cm ML x 2.5 cm AP and the isthmus measures 2.9 SI x 2.3 cm ML x 0.8 cm AP. The external surfaces have fibrous adhesions and are painted with Silver nitrate. No parathyroid glands and/or lymph nodes are identified grossly. The entire lobe contains a well delineated partly cystic containing brown fluid nodule(s) that measure 4.5 cm SI x 3.2 cm ML x 2.5 cm AP. The remainder of the thyroid tissue is grossly unremarkable. Sections of nodule and normal tissue are stored frozen. The remainder of the specimen is submitted in toto

Surgical Pathology Consultation Report

2A-H lobe, superior to inferior (D-E section bisected)
2I-L isthmus

Quick Section Diagnosis

1. Right lower parathyroid, qs: - Parathyroid tissue identified
OR called
