

ICD-0-3
 carcinoma, papillary, follicular variant
 8340/3
 Site: thyroid, NOS C23.9
 for 4/26/11

Surgical Pathology Consultation Report

Patient Name: _____
 MRN: _____ Service: _____ Accession #: _____
 DOB: _____ Visit #: _____ Collected: _____
 Gender: F Location: _____ Received: _____
 HCN: _____ Facility: _____ Reported: _____
 Ordering MD: _____
 Copy To: _____

Specimen(s) Received

1. Thyroid: right subtotal hemithyroid stitch upper pole

Diagnosis

Multifocal papillary carcinoma, dominant follicular variant, 2.2 cm: Thyroid
 No pathological diagnosis: Lymph node, isthmus
 - Right hemithyroidectomy specimen

Synoptic Data

Procedure: Right hemithyroidectomy
 Received: Fresh
 Specimen Integrity: Intact
 Specimen Size: Right lobe:
 3.9 cm
 2.3 cm
 1.6 cm
 Isthmus +/- pyramidal lobe:
 1.7 cm
 1.6 cm
 0.7 cm
 8.7 g
 Specimen Weight: 8.7 g
 Tumor Focality: Multifocal, ipsilateral
 ----- DOMINANT TUMOR -----
 Tumor Laterality: Right lobe
 Tumor Size: Greatest dimension: 2.2cm
 Additional dimension: 2.1 cm
 Additional dimension: 1.5 cm
 Histologic Type: Papillary carcinoma, follicular variant per TSS, follicular variant = 100%
 Papillary carcinoma, macrofollicular variant
 Follicular architecture
 Macrofollicular architecture
 Classical cytomorphology
 Histologic Grade: Not applicable
 Margins: Margins uninvolved by carcinoma
 Tumor Capsule: Partially encapsulated
 Tumor Capsular Invasion: Present, extent minimal
 Lymph-Vascular Invasion: Not identified

UUID: D48FB953-758B-4515-B244-09DCACD29441
 TCGA-EM-A22L-01A-PR

Redacted



Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
HPAA Discrepancy		
Prior Malignancy History		
Dual/Sync. Primary		
Case is (Circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	Date Reviewed:	

KMI 4/25/11 4/26/11 for

Perineural Invasion:	Not identified
Extrathyroidal Extension:	Not identified
----- SECOND TUMOR -----	
Tumor Laterality:	Right lobe
Tumor Size:	Greatest dimension: 0.1cm
Histologic Type:	Papillary carcinoma, follicular variant
	Papillary carcinoma, microcarcinoma (occult, small or microscopic) variant
	Follicular architecture
	Classical cytomorphology
Histologic Grade:	Not applicable
Margins:	Margins uninvolved by carcinoma
Tumor Capsule:	None
Tumor Capsular Invasion:	Not identified
Lymph-Vascular Invasion:	Not identified
Perineural Invasion:	Not identified
Extrathyroidal Extension:	Not identified
TNM Descriptors:	m (multiple primary tumors)
Primary Tumor (pT):	pT2: Tumor more than 2 cm, but not more than 4 cm, limited to thyroid
Regional Lymph Nodes (pN):	pN0: No regional lymph node metastasis
	Number of regional lymph nodes examined: 1
	Number of regional lymph nodes involved: 0
Distant Metastasis (pM):	Not applicable
Additional Pathologic Findings:	None Identified

*Pathologic Staging is based on AJCC/UICC TNM, 7th Edition

Comment

The dominant nodule exhibits multifocal transformation to papillary carcinoma and the extent of malignancy warrants classification of the entire nodule as papillary carcinoma. There is also a 0.1cm follicular variant papillary microcarcinoma in the right lobe. The overall morphological features are those of a low risk papillary carcinoma.

Electronically verified

by:

Gross Description

The specimen labeled with the patient's name and as "right subtotal thyroid stitch upper pole", consists of a right hemithyroidectomy specimen that weighs 8.7 g. The lobe measures 3.9 cm SI x 2.3 cm ML x 1.6 cm AP and the isthmus measures 1.7 SI x 1.6cm ML x 0.7cm AP. No parathyroid gland or lymph nodes are identified grossly. The mid superior lobe contains one well delineated nodule that measures 2.1 cm SI x 2.2cm ML x 1.5 cm AP. The cut surface of the nodule has a variegated red-tan color, and has a soft and flabby consistency. Sections of nodule and normal tissue are stored frozen. The remainder of the specimen is submitted as follows:

- 1A-F lobe, superior to inferior
- 1G isthmus