

UUID: 77194F16-66D9-4D4F-B111-0AC53989F6DC
TCGA-EM-A2CU-01A-PR Redacted



Surgical Pathology Consultation Report

Patient
Name:
MRN:
DOB:
Gender: F
HCN:
Ordering
MD:
Copy To:

Service:
Visit #:
Location:
Facility:

Accession #:
Collected:
Received:
Reported:

Specimen(s) Received

1. Soft-Tissue: Query right inferior parathyroid
2. Lymph node: Left paratracheal node
3. Thyroid: Total thyroid stitch left superior pole
4. Lymph node: Right paratracheal node

Criteria	Yes	No
Diagnosis Discrepancy		X
Primary Tumor Site Discrepancy		X
HIPAA Discrepancy		X
Prior Malignancy History		X
Dual/Synchronous Primary Noted		X
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewed Initials:	KMM	6/18/11

Diagnosis

1. No pathological diagnosis: Parathyroid (right inferior) biopsy
2. No pathological diagnosis: Lymph nodes, 4 (left paratracheal) biopsy
3. Papillary carcinoma, follicular variant, right, 4.0 cm; underlying focal chronic thyroiditis: Thyroid
Metastatic papillary thyroid carcinoma: Lymph nodes, 3 of 23 central compartment
No pathological diagnosis: Lymph nodes, 1, isthmic, 1 left perithyroidal
No pathological diagnosis: Parathyroids, 2, right perithyroidal and left central compartment
- total thyroidectomy specimen
4. Metastatic papillary thyroid carcinoma: Lymph node, 1 (right paratracheal) excisional biopsy

Synoptic Data

Procedure:

Received:

Specimen Integrity:

Specimen Size:

Total thyroidectomy with central compartment dissection

Fresh

Intact

Right lobe:

4.2 cm

3.3 cm

2.2 cm

Left lobe:

3.6 cm

2.3 cm

1.0 cm

1CD-0-3

carcinoma, papillary, follicular variant
8340/3

Site: thyroid, NOS C73.9 6/18/11

Isthmus +/- pyramidal lobe:
 1.2 cm
 1.2 cm
 0.6 cm
 Central compartment:
 4.8 cm
 1.6 cm
 1.2 cm
 Specimen Weight: 28.7 g
 Tumor Focality: Unifocal
 ----- DOMINANT TUMOR -----
 Tumor Laterality: ~~Right Lobes~~
 Tumor Size: Greatest dimension: 4.0cm
 Additional dimension: 2.8 cm
 Additional dimension: 1.7 cm
 Histologic Type: ~~Papillary carcinoma, follicular variant~~ *per TSS, follicular variant = 100%*
 Follicular architecture
 Classical cytomorphology
 Histologic Grade: Not applicable
 Margins: Margins uninvolved by carcinoma
 Tumor Capsule: None
 Tumor Capsular Invasion: Present, extent minimal
 Lymph-Vascular Invasion: Not identified
 Perineural Invasion: Not identified
 Extrathyroidal Extension: Not identified
 TNM Descriptors: Not applicable
 Primary Tumor (pT): pT2: Tumor more than 2 cm, but not more than 4 cm, limited to thyroid
 Regional Lymph Nodes (pN): pN1a: Nodal metastases to Level VI (pretracheal, paratracheal and prelaryngeal/Delphian) lymph nodes
 Number of regional lymph nodes examined: 30
 Number of regional lymph nodes involved: 4
 Lymph Node, Extranodal Extension Not identified
 Distant Metastasis (pM): Not applicable
 Additional Pathologic Findings: Thyroiditis, focal (nonspecific)

*Pathologic Staging is based on AJCC/UICC TNM, 7th Edition

verified by:

Electronically

Gross Description

1. The specimen labeled with the patient's name and as "Soft Tissue: Query right inferior parathyroid" consists of a piece of fibrous tissue that measures 0.1 x 0.1 cm. It is frozen for intraoperative consultation.

1A frozen section block resubmitted

2. The specimen labeled with the patient's name and as "Lymph Node: Left paratracheal node" consists of two fragments of red-yellow fatty tissue measuring 1.2 x 0.5 x 0.5 cm and 1.4 x 0.7 x 0.5 cm.

2A submitted in toto.

3. The specimen labeled with the patient's name and as "Thyroid: Total thyroid stitch left superior pole" consists of a thyroid gland that is oriented with a stitch and weighs 28.7 g. The right lobe measures 4.2 cm SI x 3.3 cm ML x 2.2 cm AP, the left lobe measures 3.6 cm SI x 2.3 cm ML x 1.0 cm AP and the isthmus measures 1.2 cm SI x 1.2 cm ML x 0.6 cm AP. The external surfaces have fibrous adhesions. No parathyroid glands are identified grossly. Attached to the inferior aspect of the left lobe is the central compartment contents that measure 4.8 x 1.6 x 1.2 cm and contain multiple lymph nodes ranging in size from 0.4 cm to 0.8 cm. The external surface is painted with Silver nitrate. The right lobe contains a well delineated nodule that measures 4.0 cm SI x 2.8 cm ML x 1.7 cm AP. The remainder of the thyroid tissue is unremarkable. Sections of right lobe nodule and normal tissue are stored frozen. The remainder of the specimen is submitted as follows:

3A-J right lobe, superior to inferior

3K-L isthmus

3M-R left lobe, superior to inferior

3S-T multiple lymph nodes each block

3U-V fatty tissue attached to the inferior aspect of the left lobe in toto.

4. The specimen labeled with the patient's name and as "Lymph Node: Right paratracheal node" consists of a nodule with attached fatty tissue measuring 3.2 x 2.6 x 1.7 cm received in 10% buffered formalin.

4A-E submitted in toto.

Quick Section Diagnosis

1A – parathyroid tissue. Reported to
