

a
Heterozygous SNVs per callable kb
1.0
0.9
0.8
0.7
0.6
AFR
ρ = 0.04
AMR
ρ = 0.24
EAS
ρ = 0.07
EUR
ρ = −0.08
SAS
ρ = 0.03
ρ: Spearman's, with raw depth
b
θVCF / θpsmcfa − 1 (%)
10
8
6
4
2
0
Bases missing in the called bins:
7.6% of each genome's
AFR
AMR
EAS
EUR
SAS
Superpopulation (52 genomes)
c
log10 first / next window Ne, split
2.0
1.5
1.0
0.5
0.0
−0.5
−0.5
0.0
0.5
1.0
1.5
2.0
log10 first / next window Ne, default
d
C psmc (psmcfa)
88.0 min (n = 52); 14.1x psmc-rs (psmcfa)
psmc-rs (psmcfa)
6.2 min (n = 2,504)
psmc-rs (VCF + mask)
5.9 min (n = 52)
psmc-rs (split first window)
5.6 min (n = 26)
psmc-rs bootstrap, per fit
7.8 min (3 runs of 101 fits, 32 threads: 24.7 min each)
0
10
20
30
40
50
60
70
80
Time per fit (min): median wall time on one thread; bootstrap: core time per fit
c: superpopulation and keys
AFR
AMR
EAS
EUR
SAS
An interval at psmc-rs's bound (λ = 10^4)
Mean of the other intervals
Range of the four split intervals
No change (diagonal)
NA12878 against GIAB HG001 v4.2.1:
1,564,410 het SNVs in 2,109 Mb;
precision 0.9992, recall 0.9966