

BEYOND THE FORD-CIRCLE LAW: PARABOLIC RENORMALIZATION AND THE SHAPE OF MANDELBROT SATELLITE BULBS (ABRIDGED VERSION)

LARS BENDIXEN

ABSTRACT. The main cardioid of the Mandelbrot set carries a “bulb” (a satellite component) at every rational internal angle p/q . To leading order the bulb at p/q is a round disc of diameter $2\sin(\pi p/q)/q^2$, which is the scaling of the Ford circles of number theory. We study how the bulbs deviate from this picture. A bulb is described by the multiplier of its attracting cycle. In the rescaled parameter $u = q^2\varepsilon$ this multiplier begins $1 - u$ for every bulb, which is the Ford law. The first coefficient that distinguishes bulbs, $\kappa(p/q)$, measures how far the bulb is from round. We prove that $\kappa(p/q) = (\iota_{p/q} - \frac{1}{2})/q$, where $\iota_{p/q}$ is the holomorphic index of the parabolic fixed point from which the bulb is born, and we give an efficient recursion for it. The rest of the work is numerical. It shows that κ is controlled by parabolic renormalization, which zooms in on the parabolic point and returns a new map, its horn map. For the bulbs at angles $1/q$, κ tends to a new constant $\kappa_0 \approx 0.02383 - 0.05230i$. This is the index of the horn map of $z + z^2$, and the two agree to 24 digits. The same mechanism, combined with an exactly computable phase, predicts the bulb sizes and the whole first-order correction in $1/q$. Repeating the renormalization along continued fractions gives constants that converge to a limit κ_* . At each rational tested, the bulbs approaching it from above and from below have different limits, given by the two horn maps of the parabolic point there; we conjecture that this holds at every rational. The bulbs of other components, and those of the family $z^3 + c$, follow the same principle with their own parabolic germs.

1. INTRODUCTION

Motivation. Pictures of the Mandelbrot set show the bulbs: discs along the edge of the main cardioid, one for each fraction p/q , shrinking as q grows. Their arrangement is arithmetic. Between two bulbs at Farey neighbours the largest sits at the mediant, and the sizes fall off like $1/q^2$, as do the radii of the Ford circles. This dictionary between the Mandelbrot set and number theory is well known, but it is a leading-order statement: the bulbs are discs of size about $1/q^2$. Looked at closely, the bulbs are not round, and each deviates from a disc in its own way. We ask whether that deviation is still arithmetic, and which arithmetic governs it. The answer is the continued fraction of p/q , read from its end by parabolic renormalization.

This makes the correction a quantitative test of near-parabolic renormalization [11], a tool that has so far mostly settled qualitative questions [3, 4]. Here it predicts one complex number per bulb, which the index formula computes exactly; for the bulbs at $1/q$ the two agree to 24 digits. The process produces constants that resemble Feigenbaum’s [5] in that they come from a renormalization operator, but they depend on the degree. Read as a function of the angle,

Date: September 2026.

2020 Mathematics Subject Classification. Primary 37F46; Secondary 37F25, 37F10.

Key words and phrases. Mandelbrot set, satellite components, Ford circles, holomorphic index, horn map, parabolic renormalization.

the correction jumps at every rational tested, as some continued-fraction functions of number theory do.

Setting. Let $c(\lambda) = \lambda/2 - \lambda^2/4$ parametrize the main cardioid of the Mandelbrot set by the multiplier $\lambda \in \mathbb{D}$ of its attracting fixed point; equivalently we use $f_\lambda(z) = \lambda z + z^2$. At $\lambda_0 = e^{2\pi i p/q}$ the satellite $W_{p/q}$ of period q is attached [6]. Its size is of order q^{-2} [10], and to first order it is a disc of diameter $2|c'(\lambda_0)|q^{-2} = 2\sin(\pi p/q)q^{-2}$ [8]. Up to the factor $\sin(\pi p/q)$ these are the Ford circles [9]. We study the first correction to this law.

Put $\lambda = \lambda_0 e^\varepsilon$, let $\rho_{p/q}(\varepsilon)$ be the multiplier of the q -cycle that bifurcates from 0, and write

$$(1) \quad R_{p/q}(u) := \rho_{p/q}(u/q^2) = \sum_{k \geq 0} r_k(p/q) u^k.$$

Up to a conformal change of variable, $W_{p/q} = \{|R_{p/q}| < 1\}$. We show that $r_0 = 1$ and $r_1 = -1$ for all p/q , so that to first order $W_{p/q}$ is the disc $|u - 1| < 1$; this is the Ford law. The first coefficient that depends on p/q is $\kappa(p/q) := r_2(p/q)$, and it governs the leading deviation of $W_{p/q}$ from a disc. The bulb size is measured by $G(p/q) = \frac{1}{2}|u_a|(1 + O(q^{-2}))$, where $R_{p/q}(u_a) = -1$ and $G = 1$ to first order.

Results. Section 2 contains the proved statements: the index formula for κ , and a normal form that computes it in $O(q^2)$ operations. The remaining sections are numerical, and each claim is stated as a conjecture together with its evidence. We find that, up to $O(q^{-2})$, the bulb is described by the near-parabolic renormalization of f_{λ_0} in the sense of Inou and Shishikura [11], composed with an exact phase. This picture yields five constants, collected in Table 1. The full version [1] contains proofs omitted here, degree-three details, error analysis and data; all code and data are in [15].

2. EXACT RESULTS

Write $f = f_{\lambda_0}$. The fixed point 0 has a single cycle of q petals, so $f^q(z) = z + az^{q+1} + O(z^{q+2})$ with $a \neq 0$. Let $\iota_{p/q} = \text{Res}_{z=0} dz/(z - f^q(z))$ be its holomorphic index [14].

Theorem 2.1 (Index formula). $\rho_{p/q}(\varepsilon) = 1 - q^2\varepsilon + q^3(\iota_{p/q} - \frac{1}{2})\varepsilon^2 + O(\varepsilon^3)$. Equivalently, $r_0 = 1$, $r_1 = -1$ and $\kappa(p/q) = (\iota_{p/q} - \frac{1}{2})/q$.

Sketch. For small $\varepsilon \neq 0$, f_λ^q has $q + 1$ simple fixed points near 0: the point 0, with multiplier $e^{q\varepsilon}$, and one q -cycle, with multiplier ρ . The sum of their indices converges to $\iota_{p/q}$, which gives

$$\frac{1}{1 - e^{q\varepsilon}} + \frac{q}{1 - \rho(\varepsilon)} \longrightarrow \iota_{p/q}.$$

The first term is $-1/(q\varepsilon) + \frac{1}{2} + O(\varepsilon)$. Boundedness of the sum forces $1 - \rho = q^2\varepsilon + O(\varepsilon^2)$, and the constant term gives the coefficient of ε^2 . $\square$

The proof uses only the multiplicity $q + 1$, so it applies to the satellites of every hyperbolic component of $z^d + c$, with the first-return map in place of f . Since f^q has coefficients in $\mathbb{Z}[\lambda_0]$, $\kappa(p/q) \in \mathbb{Q}(\zeta_q)$, and the values $\kappa(p'/q)$ form a Galois orbit; for instance $\kappa(\frac{1}{2}) = -\frac{3}{16}$ and $\kappa(\frac{1}{3}) = (-241 - 16\sqrt{3}i)/2646$.

Theorem 2.2 (Normal form). Let $\mathcal{N}(z) = \lambda_0 z(1 + b_1 z^q + b_2 z^{2q})$ be the resonant normal form of f to order $2q + 1$. Its coefficients satisfy an explicit triangular recursion of cost $O(q^2)$, and

$$a = qb_1, \quad \kappa(p/q) = \frac{1}{2} - \frac{1}{2q} - \frac{1}{2q^2} + \frac{b_2}{(qb_1)^2} = \iota(Q_{p/q}) - \frac{1}{2},$$

where $Q_{p/q}(W) = \mathcal{N}^q(W^{1/q})^q$ is the rotation quotient of $\mathcal{N}^q$.

constant	germ	value	bulbs	
κ_0	$z + z^2$	$0.02382588740220057 - 0.05230465911400367 i$	$3 \cdot 10^{-28}$	§3
$\kappa_0^{(3)}$	$v + v^2 + \frac{1}{3}v^3$	$0.1435992780952916 - 0.03031164789850254 i$	$1.3 \cdot 10^{-6}$ ($q = 1024$)	§7
$\kappa_{1/2}^+$	$-z + z^2$	$0.01840526161796478 - 0.04290372388023048 i$	$3 \cdot 10^{-10}$	§6
κ_*	$\mathcal{G}_* = \mathcal{P}(\bar{\mathcal{G}}_*)$	$0.011929183413 - 0.037455493752 i$	levels ≤ 3	§5
G_0	$e^u \mathcal{P}_0$	1.02641539	$4 \cdot 10^{-8}$	§4

TABLE 1. The constants. “Bulbs” is the difference between the horn-map value and the best extrapolated bulb limit ($\kappa_0^{(3)}$: error at $q = 1024$, decaying like q^{-2}). The extrapolation uncertainties are about $3 \cdot 10^{-25}$ for κ_0 , $2 \cdot 10^{-7}$ for $\kappa_{1/2}^+$ and 10^{-6} for G_0 .

Composing f symbolically q times costs $O(q^3)$ and suffers severe cancellation, since $|a(1/q)|$ grows like $(q/2\pi)^q$. The recursion is stable; in Arb ball arithmetic [2] it certifies $\kappa(1/q)$ for $q \leq 32768$.

3. THE HORN MAP OF $z + z^2$ AND κ_0

Let $E(Z) = \Phi_{\text{att}} \circ \Psi_{\text{rep}}(Z) = Z + \sum_{n \geq 1} a_n e^{2\pi i n Z}$ be the upper horn map of $f_1(z) = z + z^2$, with the standard normalization $a_0 = 0$ [13, 16]. Let

$$\mathcal{P}_0(W) = W \exp\left(2\pi i \sum_{n \geq 1} a_n W^n\right)$$

be its parabolic renormalization [11, 12].

Lemma 3.1. $\iota(\mathcal{P}_0) - \frac{1}{2} = a_2/(2\pi i a_1^2)$, independently of the normalization of the Fatou coordinates.

Proof. $\mathcal{P}_0(W) = W + AW^2 + BW^3 + \dots$ with $A = 2\pi i a_1$ and $B = 2\pi i a_2 + \frac{1}{2}A^2$, and $\iota = B/A^2$. A translation of both Fatou coordinates by τ sends $a_n \mapsto a_n e^{2\pi i n \tau}$, which leaves a_2/a_1^2 unchanged. $\square$

Numerical result 3.2. The horn map, computed from exact Fatou series at 60 and 80 digits, gives

$$\begin{aligned} \operatorname{Re} \kappa_0 &= 0.023825887402200569000729189645402, \\ \operatorname{Im} \kappa_0 &= -0.052304659114003666316522486207571, \end{aligned}$$

where $\kappa_0 := \iota(\mathcal{P}_0) - \frac{1}{2}$. Least-squares fits $\kappa(1/q) = \sum_{j \leq 10} \gamma_j q^{-j}$ over certified values $256 \leq q \leq 32768$ give γ_0 equal to this value to $3 \cdot 10^{-28}$, and $\gamma_1 = -i/(2\pi)$ to $2 \cdot 10^{-23}$. Other stable fits move γ_0 by at most $3 \cdot 10^{-25}$.

Conjecture 3.3. $\kappa(1/q) = \kappa_0 - \frac{i}{2\pi q} + O(q^{-2})$.

4. THE RENORMALIZATION PRINCIPLE

Near-parabolic renormalization $\mathcal{R}$ [11] assigns to $f_{e^{2\pi i \alpha}}$ a germ with multiplier $e^{-2\pi i/\alpha}$, and parabolic implosion [13, 16] identifies its limits with $\mathcal{P}_0$ times a phase. The phase is exact.

Proposition 4.1 (Kinematics). *If $e^{2\pi i \alpha} = \lambda_0 e^{u/q^2}$, then $e^{-2\pi i/\alpha} = e^{-2\pi i q/p} e^{\tilde{u}/p^2}$ with $\tilde{u} = u/(1 + u/(2\pi i p q))$.*




FIGURE 1. $\kappa(1/q) \rightarrow \kappa_0$ (degree two) and $\kappa^{(3)}(1/q) \rightarrow \kappa_0^{(3)}$ (degree three). After the universal correction $-i/(2\pi q)$ the error decays like q^{-2} .

Proof. $\alpha = (2\pi ipq + u)/(2\pi iq^2)$, and $2\pi ip^2(q/p - 1/\alpha) = \tilde{u}$. □

Fix p and r coprime, let $q = pN + r$, $\mu = e^{-2\pi ir/p}$, and let $\rho_{p,r}(u)$ be the multiplier of the p -cycle of $\mathcal{M}_u = \mu e^{u/p^2} \mathcal{P}_0$ that bifurcates from 0.

Conjecture 4.2 (Renormalization principle). $R_{p/q}(u) = \rho_{p,r}(\tilde{u}(u)) + O(q^{-2})$ coefficientwise, and uniformly on a neighbourhood of the arc from 0 to the point where $\rho_{p,r} = -1$.

Comparing coefficients gives $\kappa(p/q) = \kappa_{p,r} - i/(2\pi pq) + O(q^{-2})$, where

$$(2) \quad \kappa_{p,r} = \frac{\iota((\mu \mathcal{P}_0)^p) - \frac{1}{2}}{p},$$

by Theorem 2.1 applied to $\mu e^\varepsilon \mathcal{P}_0$. The first-order term involves no property of the germ.

Evidence. In all nine classes with $2 \leq p \leq 5$, (2) agrees with the extrapolated bulb limits to between $6 \cdot 10^{-9}$ and $2.6 \cdot 10^{-8}$, within the extrapolation uncertainty. The opposite phase fails.

The predicted bulb sizes agree with the bulbs to $9 \cdot 10^{-7}$ in all ten classes with $p \leq 5$. For $p = 1$ this gives diameters $2\pi G_0 q^{-3}(1 + O(q^{-1}))$ for the $1/q$ -bulbs at the cusp, with $G_0 = 1.02641539\dots$

For $2 \leq k \leq 8$, $(p, r) \in \{(1, 1), (2, 1), (3, 1), (3, 2)\}$ and $N = 64, \dots, 1024$, the residuals $|r_k(p/q) - [u^k] \rho_{p,r}(\tilde{u})|$ fall by a factor 3.8–4.3 per doubling of N , that is, like q^{-2} . Without $u \mapsto \tilde{u}$ the factor is only 1.1–2.0 (Figure 2). Numerically, then, the entire $O(1/q)$ part of $R_{p/q}$ is kinematic.

5. THE HIERARCHY AND κ_*

The renormalization $\mathcal{R}$ acts on continued fractions as the Gauss map, after conjugation by $z \mapsto \bar{z}$. Therefore, when the first k partial quotients are large, each renormalization should be close to $\mathcal{P}$ applied to the complex conjugate of the previous one. Normalize germs to




FIGURE 2. Residuals of r_2, r_3 of $R_{1/q}$ against the renormalized family, with (solid) and without (dashed) the Möbius substitution $u \mapsto \tilde{u}$.

$v + v^2 + O(v^3)$, write $\bar{\mathcal{G}}(W) = \overline{\mathcal{G}(\bar{W})}$, and put

$$\mathcal{G}_1 = \mathcal{P}(f_1), \quad \mathcal{G}_{k+1} = \mathcal{P}(\bar{\mathcal{G}}_k), \quad \kappa_k = \iota(\mathcal{G}_k) - \frac{1}{2}.$$

Numerical result 5.1. Two independent settings agree to $2 \cdot 10^{-14}$ for $k \leq 7$. The differences $\kappa_{k+1} - \kappa_k$ contract antilinearly, $\Delta \mapsto \nu \bar{\Delta}$ with $|\nu| = 0.0419$. This is the signature of the antilinear derivative of $\mathcal{G} \mapsto \mathcal{P}(\bar{\mathcal{G}})$. Extrapolation gives

$$\kappa_* = \lim_k \kappa_k = 0.011929183413 - 0.037455493752 i.$$

Without the conjugation our code reproduces the fixed point of Lanford and Yampolsky [12]: cubic coefficient $0.5142768 - 0.0346276 i$, eigenvalue ratio $-0.01764 + 0.04038 i$. That operator, however, does not govern the bulbs.

Conjecture 5.2. $\lim_{N_k \rightarrow \infty} \cdots \lim_{N_1 \rightarrow \infty} \kappa([0; N_1, \dots, N_k])$ equals κ_k for k odd and $\bar{\kappa}_k$ for k even.

This holds numerically for $k \leq 3$. At $k = 2$ the bounded- p route gives $\bar{\kappa}_2$ to $1.2 \cdot 10^{-8}$, and the diagonal $[0; N, N]$ gives it to $3 \cdot 10^{-10}$. At $k = 3$ the diagonal $[0; N, N, N]$, with $q \leq 64080$, gives κ_3 to 10^{-5} , while $|\kappa_3 - \kappa_2| = 7.8 \cdot 10^{-4}$.

6. LIMITS AT PARABOLIC ROOTS

Let $p/q = [0; a_1, \dots, a_\ell]$, using either of its two continued-fraction expansions. Let τ be a fixed tail with $[0; \tau] = r_\tau/p_\tau$, where $r_\tau = 0$ and $p_\tau = 1$ for $\tau = \emptyset$. As $N \rightarrow \infty$ the satellites $[0; a_1, \dots, a_\ell, N, \tau]$ approach the root of $W_{p/q}$, from above if ℓ is even and from below if ℓ is odd. There the cusp germ is replaced by the q -petal germ $f_{e^{2\pi i p/q}}(z) = e^{2\pi i p/q} z + z^2$.

The construction of Section 3 extends to $g = f_{e^{2\pi i p/q}}^q$. There is a formal Fatou coordinate

$$\Phi = -z^{-q}/(qa) + \sum s_k z^k + \beta \log z,$$

where $\beta = \text{résit}(g) = \frac{q+1}{2} - \iota(g)$; this follows from a residue computation. The orbits from a repelling petal above or below an escape band land in different attracting petals, and are carried back by f^m with $m \equiv -jp^{-1} \pmod{q}$. This defines an upper and a lower horn map, with germs $\mathcal{P}_{p/q}^\pm$ normalized tangent to the identity. Put $\kappa_{p/q}^\pm = \iota(\mathcal{P}_{p/q}^\pm) - \frac{1}{2}$. Then $\kappa_{0/1}^\pm = \kappa_0, \overline{\kappa_0}$ and $\kappa_{p/q}^- = \overline{\kappa_{(q-p)/q}^+}$, but for $q \geq 3$ the constants $\kappa_{p/q}^+$ and $\kappa_{p/q}^-$ at the same p/q are not conjugate.

Conjecture 6.1 (Root limits). Let $\mu_\tau = e^{-2\pi i r_\tau/p_\tau}$. From above,

$$\lim_{N \rightarrow \infty} \kappa([0; a_1, \dots, a_\ell, N, \tau]) = \frac{\iota((\mu_\tau \mathcal{P}_{p/q}^+)^{p_\tau}) - \frac{1}{2}}{p_\tau}.$$

From below the same formula holds with $\overline{\mu_\tau} \mathcal{P}_{p/q}^-$ in place of $\mu_\tau \mathcal{P}_{p/q}^+$. In particular κ has the distinct one-sided limits $\kappa_{p/q}^\pm$ at every rational.

The conjugate phase from below is forced by $\kappa(1-x) = \overline{\kappa(x)}$. Table 2 tests the conjecture in fifteen cases. Eight have $\tau = \emptyset$, at $\frac{1}{3}, \frac{1}{4}, \frac{2}{5}, \frac{3}{7}$ from both sides. Seven have tails with $p_\tau = 2, 3$, at $\frac{1}{3}, \frac{1}{2}$ and $\frac{1}{4}$; four of these approach from below with a non-real phase. All agree to $6 \cdot 10^{-11}$. At $\frac{1}{3}$, for instance, $\kappa_{1/3}^- = 0.0158408 + 0.0401752i$ and $\kappa_{1/3}^+ = 0.0165812 - 0.0395047i$.

Other components. For a hyperbolic component W , call the first-return map of the parabolic cycle at its root the *root germ* of W . We conjecture (the root principle) that the limits of κ and of G along the satellites of W at angles $[0; N, \tau]$ and $1 - [0; N, \tau]$ are given by the rule above, with the upper and lower horn germs of the root germ. We tested this for the following components:

- the period-two disc: its root germ is $f_{e^{2\pi i/2}}$, and its constant $\kappa_{1/2}^+$ is known to 32 digits, and the best bulb fits agree with it to $2 \cdot 10^{-9}$ (uncertainty $2 \cdot 10^{-7}$);
- the period-three disc $W_{1/3}$: both sides follow $\mathcal{P}_{1/3}^\pm$, for κ with the tails $\tau = \emptyset, 2, 3, (1, 2)$ and for G with $\tau = \emptyset, 2, 3$. The conjugate phase from below is confirmed directly: the unconjugated one misses by $1.8 \cdot 10^{-2}$. The other three candidate germs miss by 10^{-3} or more;
- the real period-three component, whose root is a cusp with root germ normalized to $v + v^2 - \frac{2}{49}v^3 + \dots$ (exactly): agreement to $4 \cdot 10^{-10}$, with a kinematic first-order term;
- a period-six component at the $\frac{1}{3}$ -root of the period-two disc.

At that last root the root germ is not formally conjugate to $f_{e^{2\pi i/3}}$. Its root index differs by $6 \cdot 10^{-2}$, yet its horn constants differ from $\kappa_{1/3}^\pm$ by only about $1.5 \cdot 10^{-4}$, and the satellites follow its own constants.

7. DEGREE THREE AND OUTLOOK

For $z^3 + c$, near the cusp of the main component, the germ is $v + v^2 + \frac{1}{3}v^3$. Its horn map gives $\kappa_0^{(3)} = 0.1435992780952916 - 0.0303116478985025i$, and $\kappa^{(3)}(1/q) + i/(2\pi q) \rightarrow \kappa_0^{(3)}$ like q^{-2} . The universal first-order term is unchanged and the constant is not. Here each bulb has two points of multiplier -1 , and both of their sizes are predicted. Integer-relation searches (PSLQ) found no closed form for $\kappa_0, \kappa_0^{(3)}$ or $\kappa_{1/2}^+$ in a small basis.

Open problems.

- (1) Prove Conjectures 3.3 and 4.2 via continuity of near-parabolic renormalization.
- (2) Prove that the fixed point $\mathcal{G}_* = \mathcal{P}(\tilde{\mathcal{G}}_*)$ exists and describe its spectrum.
- (3) Prove Conjecture 6.1, and explain the jumps $\kappa_{p/q}^+ - \kappa_{p/q}^-$. They are nearly imaginary and depend mainly on q .

p/q	side	τ	horn map of $e^{2\pi ip/q}z + z^2$	bulbs, extrapolated	$ \Delta $
1/3	below	$\emptyset$	$0.015840773234 + 0.040175221242i$	$0.015840773241 + 0.040175221243i$	$6.5 \cdot 10^{-12}$
1/3	above	$\emptyset$	$0.016581162594 - 0.039504714589i$	$0.016581162601 - 0.039504714589i$	$6.4 \cdot 10^{-12}$
1/4	below	$\emptyset$	$0.014455842876 + 0.039068895117i$	$0.014455842881 + 0.039068895118i$	$5.3 \cdot 10^{-12}$
1/4	above	$\emptyset$	$0.015760467204 - 0.037801851383i$	$0.015760467209 - 0.037801851382i$	$5.3 \cdot 10^{-12}$
2/5	below	$\emptyset$	$0.014434584508 + 0.037505178012i$	$0.014434584513 + 0.037505178012i$	$4.6 \cdot 10^{-12}$
2/5	above	$\emptyset$	$0.014296912442 - 0.037686119109i$	$0.014296912446 - 0.037686119110i$	$4.6 \cdot 10^{-12}$
3/7	above	$\emptyset$	$0.013301284920 - 0.037190044710i$	$0.013301284861 - 0.037190044712i$	$5.9 \cdot 10^{-11}$
3/7	below	$\emptyset$	$0.013998687363 + 0.036367863348i$	$0.013998687363 + 0.036367863334i$	$1.4 \cdot 10^{-11}$
1/3	above	2	$0.046849904480 - 0.018242741860i$	$0.046849904482 - 0.018242741860i$	$2.7 \cdot 10^{-12}$
1/3	below	2	$0.046510029333 + 0.018510459943i$	$0.046510029336 + 0.018510459943i$	$2.8 \cdot 10^{-12}$
1/3	above	3	$0.052913142048 - 0.001728643502i$	$0.052913142050 - 0.001728643502i$	$1.7 \cdot 10^{-12}$
1/2	below	3	$0.053307011907 + 0.002605955801i$	$0.053307011909 + 0.002605955801i$	$2.5 \cdot 10^{-12}$
1/3	below	3	$0.052713652883 + 0.001878998752i$	$0.052713652885 + 0.001878998752i$	$1.7 \cdot 10^{-12}$
1/1	below	3	$0.054529447219 + 0.005017189409i$	$0.054529447226 + 0.005017189408i$	$6.7 \cdot 10^{-12}$
1/3	below	(1, 2)	$0.053167372528 + 0.020247303769i$	$0.053167372529 + 0.020247303768i$	$1.6 \cdot 10^{-12}$

TABLE 2. One-sided limits of κ at rationals: the tail rule of Conjecture 6.1 against the bulbs $[0; a_1, \dots, a_\ell, N, \tau]$, extrapolated from $N = 128, \dots, 2048$ (degree four against degree three: at most $1.3 \cdot 10^{-9}$).

(4) Relate κ_0 to the Écalé–Voronin invariants of $z + z^2$ [7, 17].

REFERENCES

- [1] L. Bendixen, *Beyond the Ford-circle law: parabolic renormalization and the shape of Mandelbrot satellite bulbs*, full version, 2026; in [15], [paper/kappa0.pdf](#).
- [2] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), 1281–1292.
- [3] X. Buff and A. Chéritat, *The Brjuno function continuously estimates the size of quadratic Siegel disks*, Ann. of Math. (2) **164** (2006), 265–312.
- [4] X. Buff and A. Chéritat, *Quadratic Julia sets with positive area*, Ann. of Math. (2) **176** (2012), 673–746.
- [5] M. J. Feigenbaum, *Quantitative universality for a class of nonlinear transformations*, J. Stat. Phys. **19** (1978), 25–52.
- [6] A. Douady and J. H. Hubbard, *Étude dynamique des polynômes complexes*, Parties I et II, Publ. Math. d’Orsay 84-02, 85-04, 1984–1985.
- [7] J. Écalé, *Les fonctions résurgentes*, Tomes I–III, Publ. Math. d’Orsay, 1981–1985.
- [8] A. C. Fowler and M. J. McGuinness, *The size of Mandelbrot bulbs*, Chaos Solitons Fractals X **3** (2019), 100019.
- [9] L. R. Ford, *Fractions*, Amer. Math. Monthly **45** (1938), 586–601.
- [10] J. Guckenheimer and R. McGehee, *A proof of the Mandelbrot N^2 conjecture*, Report No. 15, Institut Mittag-Leffler, 1984.
- [11] H. Inou and M. Shishikura, *The renormalization for parabolic fixed points and their perturbation*, preprint; <https://www.math.kyoto-u.ac.jp/~mitsu/pararenorm/>.
- [12] O. E. Lanford III and M. Yampolsky, *Fixed Point of the Parabolic Renormalization Operator*, SpringerBriefs in Mathematics, Springer, 2014.
- [13] P. Lavaurs, *Systèmes dynamiques holomorphes: explosion de points périodiques paraboliques*, Thèse, Orsay, 1989.
- [14] J. Milnor, *Dynamics in One Complex Variable*, 3rd ed., Princeton University Press, 2006.
- [15] *mandelbrot-bulbs-and-ford-circles-research*, code and data, <https://github.com/larsbx/mandelbrot-bulbs-and-ford-circles-research>.
- [16] M. Shishikura, *Bifurcation of parabolic fixed points*, in: The Mandelbrot Set, Theme and Variations, LMS Lecture Note Ser. 274, Cambridge Univ. Press, 2000, 325–363.
- [17] S. M. Voronin, *Analytic classification of germs of conformal mappings $(\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ with identity linear part*, Funct. Anal. Appl. **15** (1981), 1–13.