



Fungal isolate composition by genus
Share of all isolates. Grey = Incertae sedis (unknown or unresolved at this rank)


Lauraceae leaves
Incertae sedis: 11.4%



Ficus leaves
Incertae sedis: 20.5%



Ficus wood
Incertae sedis: 20.4%



genus

- | | | |
|-------------------|-------------------|-------------------|
| Diaporthe | Clonostachys | Filobasidium |
| Nigrospora | Cytospora | Fusarium |
| Colletotrichum | Didymella | Hypomontagnella |
| Aspergillus | Epicoccum | Linnemannia |
| Pseudocercospora | Nodulisporium | Metapochonia |
| Penicillium | Paraphaeosphaeria | Mucor |
| Pestalotiopsis | Podila | Myxospora |
| Annulohyphoxylon | Tolypocladium | Phaeosphaeria |
| Neopestalotiopsis | Trichoderma | Pseudogymnoascus |
| Xylaria | Amycosphaerella | Pyrenopezizomyces |
| Cladosporium | Acremonium | Rhynchogastrea |
| Phaeoacremonium | Beauveria | Saitozyma |
| Nemania | Cercophora | Thelonectria |
| Neofabraea | Coccomyces | Incertae sedis |
| Neofusicoccum | Debaryomyces | |