


Fungal isolate composition by order

Share of all isolates. Grey = Incertae sedis (unknown or unresolved at this rank)


Lauraceae leaves

Incertae sedis: 0.0%




Ficus leaves

Incertae sedis: 5.3%



Ficus wood

Incertae sedis: 2.4%



- order
- Diaporthales
 - leotiomyceta; sordariomyceta | Sordariomycetes | Xylariomycetidae | Amphisphaeriales
 - Xylariales
 - Eurotiales
 - Glomerellales
 - leotiomyceta; dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales
 - leotiomyceta; sordariomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales
 - Pleosporales
 - Cladosporiales
 - Togniniales
 - Helotiales
 - Botryosphaeriales
 - Coniochaetales
 - Mortierellomycota | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales
 - Tremellales
 - leotiomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales
 - Cystofilobasidiales
 - Mortierellomycotina | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales
 - Russulales
 - Sordariales
 - dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales
 - Chaetothyriales
 - Filobasidiales
 - Marthamycetales
 - Mucorales
 - Serinales
 - Thelebolales
 - leotiomyceta | Sordariomycetes | Xylariomycetidae | Amphisphaeriales
 - leotiomyceta; sordariomycet | Leotiomycetes | Incertae sedis | Rhytismatales
 - leotiomyceta; sordariomyceta | Leotiomycetes | Incertae sedis | Rhytismatales
 - Incertae sedis