



Fungal isolate composition by species
Share of all isolates. Grey = Incertae sedis (unknown or unresolved at this rank)


Lauraceae leaves
Incertae sedis: 65.8%



Ficus leaves
Incertae sedis: 53.8%



Ficus wood
Incertae sedis: 65.3%



species

- | | | | |
|--------------------------|---------------------------|----------------------|-------------------|
| aff. quetri | cf. musae 1 | aff. colombiense | cf. passiflorae 1 |
| crousii | cf. spathulatum | aff. godetiae | cf. passiflorae 2 |
| arboricola | cf. vesicularifera | aff. neglecta | cf. rubens |
| cf. brassicicola | humilis | aff. passiflorae 1 | cf. solani |
| cf. gorlenkoana | italicum | aff. passiflorae 2 | cf. suchlasporia |
| gorlenkoana | macarangae | aff. passiflorae 3 | cf. tabebuiae |
| austroafricanus | aff. diversiseta | aff. passiflorae 5 | hyalinus |
| hainanensis | aff. eucalypti | aff. passiflorae 6 | kochmanii |
| cf. substygium | aff. sacchari-officinarum | aff. passiflorae 7 | kongii |
| cf. godetiae | aff. verrucosum 1 | aff. passiflorae 8 | marywardiae |
| cf. mexicanum | africana | aff. spongiphila | mycoparasiticum |
| aff. brasiliensis | basitruncata | aff. terebenthifolii | podzolica |
| cf. diversiseta | bialowiezense | aff. tomentosae | pseudobassiana |
| cf. sacchari-officinarum | cf. aurantiaca gen 2 | aff. verrucosum 2 | pyriformis |
| creber | cf. farinosa | aff. viridistroma | roseopurpureum |
| aff. abortiva | cymbidicola | cf. complexa | stemmata |
| aff. adscendens 2 | guangdongensis | cf. hongkongensis | sydowii |
| aff. passiflorae 4 | rubi | cf. hunteri | uniguttulatum |
| aff. prosopidis | vancouverens | cf. musae 2 | verhagenii |
| aff. sydowii | aff. adscendens 1 | cf. ningerense | wenshanense |
| cf. aurantiaca gen 1 | aff. araucanorum | cf. passiflorae | Incertae sedis |