


Fungal isolate composition by family

Share of all isolates. Grey = Incertae sedis (unknown or unresolved at this rank)

Lauraceae leaves

Incertae sedis: 7.8%




family

- Diaporthaceae
- Apiosporaceae
- Aspergillaceae
- Glomerellaceae
- Xylariaceae
- Sporocadaceae

Ficus leaves


Incertae sedis: 12.1%



- Didymosphaeriaceae
- Botryosphaeriaceae
- Coniochaetaceae
- Cytosporaceae
- Hypocreaceae
- Mortierellomycota | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales | Mortierellaceae
- leotiomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales | Bionectriaceae
- Lasiosphaeriaceae
- Nectriaceae
- Stereaceae
- dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales | Mycosphaerellaceae
- Clavicipitaceae
- Cordycipitaceae

Ficus wood

Incertae sedis: 9.0%



- Debaryomycetaceae
- Filobasidiaceae
- Mortierellomycotina | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales | Mortierellaceae
- Mucoraceae
- Pleosporineae
- Rhynchogastremaceae
- Stachybotryaceae
- Thelebolaceae
- Trimorphomycetaceae
- leotiomyceta; sordariomycet | Leotiomycetes | Incertae sedis | Rhytismatales | Rhytismataceae
- leotiomyceta; sordariomyceta | Leotiomycetes | Incertae sedis | Rhytismatales | Rhytismataceae
- leotiomyceta; sordariomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales | Bionectriaceae
- Incertae sedis