


Fungal isolate composition by order

Share of all isolates. Grey = Incertae sedis (unknown or unresolved at this rank)


Lauraceae leaves

Incertae sedis: 0.0%




Ficus leaves

Incertae sedis: 5.3%



Ficus wood

Incertae sedis: 2.4%



order

- Diaporthales
- leotiomyceta; sordariomyceta | Sordariomycetes | Xylariomycetidae | Amphisphaeriales
- Xylariales
- Eurotiales
- Glomerellales
- leotiomyceta; dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales
- leotiomyceta; sordariomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales
- Pleosporales
- Cladosporiales
- Togniniales
- Helotiales
- Botryosphaeriales
- Coniochaetales
- Mortierellomycota | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales
- Tremellales
- leotiomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales
- Cystofilobasidiales
- Mortierellomycotina | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales
- Russulales
- Sordariales
- dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales
- Chaetothyriales
- Filobasidiales
- Marthamycetales
- Mucorales
- Serinales
- Thelebolales
- leotiomyceta | Sordariomycetes | Xylariomycetidae | Amphisphaeriales
- leotiomyceta; sordariomycet | Leotiomycetes | Incertae sedis | Rhytismatales
- leotiomyceta; sordariomyceta | Leotiomycetes | Incertae sedis | Rhytismatales
- Incertae sedis