
NMDS with top 15 taxa − substrate_x_orientation
Bray−Curtis | stress 0.109 (weak ties) / 0.2078 (tie−aware) | 39% of sample pairs share no taxa
UNRELIABLE ORDINATION − tie−aware stress 0.2078 exceeds 0.20; read the statistics, not this map
Ficus leaves − E
Ficus leaves − S
Ficus wood − E
Ficus wood − S
Lauraceae leaves − E
Lauraceae leaves − NW
Ficus leaves − N
Ficus leaves − W
Ficus wood − N
Ficus wood − W
Lauraceae leaves − N
sub_ori
Ficus leaves − W
Lauraceae leaves − N
Neopestalotiopsis sichuanensis or N. phyllostachydis or N. fidjiensis
Diaporthe ocoteae
Nigrospora hainanensis
Nigrospora gorlenkoana
Diaporthe phaseolorum
Diaporthe passiflorae
Diaporthe crousi
Diaporthe sp. 7
Diaporthe sp. 6
Annulohypoxylon stygium
Colletotrichum arboricola
Diaporthe passiflorae or D. malorum
Colletotrichum brassicicola
Pseudocercospora quetri
Pestalotiopsis telopae or P. kenyana or P. australasiae (4 spp.)
NMDS2
NMDS1
1
0
−1
−2
−2
−1
0
1
2
3