


Fungal isolate composition by family

Unknown/missing taxa pooled as Incertae sedis


Ficus leaves

Lauraceae leaves




family

- Incertae sedis
- Diaporthaceae
- Apiosporaceae
- Aspergillaceae
- Glomerellaceae
- Xylariaceae
- Sporocadaceae
- leotiomyceta; dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales | Mycosphaerellaceae
- Hypoxylaceae
- Ophiocordycipitaceae
- Cladosporiaceae
- Togniniaceae
- Didymellaceae



- Dermateaceae
- Didymosphaeriaceae
- Botryosphaeriaceae
- Coniochaetaceae
- Cytosporaceae
- Hypocreaceae
- Mortierellomycota | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales | Mortierellaceae
- leotiomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales | Bionectriaceae
- Lasiosphaeriaceae
- Nectriaceae
- Stereaceae
- dothideomyceta | Dothideomycetes | Dothideomycetidae | Mycosphaerellales | Mycosphaerellaceae
- Clavicipitaceae

Ficus wood



- Cordycipitaceae
- Debaryomycetaceae
- Filobasidiaceae
- Mortierellomycota | Incertae sedis | Mortierellomycetes | Incertae sedis | Mortierellales | Mortierellaceae
- Mucoraceae
- Pleosporineae
- Rhynchogastremaceae
- Stachybotryaceae
- Thelebolaceae
- Trimorphomycetaceae
- leotiomyceta; sordariomycet | Leotiomycetes | Incertae sedis | Rhytismatales | Rhytismataceae
- leotiomyceta; sordariomyceta | Leotiomycetes | Incertae sedis | Rhytismatales | Rhytismataceae
- leotiomyceta; sordariomyceta | Sordariomycetes | Hypocreomycetidae | Hypocreales | Bionectriaceae