

# McIntoshery in Geometry

Lashi Bandara

Institut für Mathematik  
Universität Potsdam

15 February 2018

# Message from Andreas Rosén







$$\int_0^{\infty} \| \nabla (\epsilon t \Delta u) \|^2 \frac{dt}{t} \simeq \| u \|^2 \Leftrightarrow T \text{ F.c. (McIntosh)}$$

SA Ingram 1946  
V.MATTES 1946  
SA Ingram 1946

# Part 1 - Spectral flows and harmonic analysis

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

Topologies on  $\mathcal{CF}^{\text{sa}}$ :

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

Topologies on  $\mathcal{CF}^{\text{sa}}$ :

$$\rho_G(T, S) = \left\| \frac{T+i}{T-i} - \frac{S+i}{S-i} \right\|, \quad (\text{Graph metric})$$

$$\rho_R(T, S) = \left\| \frac{T}{\sqrt{1+T^2}} - \frac{S}{\sqrt{1+S^2}} \right\|. \quad (\text{Riesz metric})$$

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

Topologies on  $\mathcal{CF}^{\text{sa}}$ :

$$\rho_G(T, S) = \left\| \frac{T+i}{T-i} - \frac{S+i}{S-i} \right\|, \quad (\text{Graph metric})$$

$$\rho_R(T, S) = \left\| \frac{T}{\sqrt{1+T^2}} - \frac{S}{\sqrt{1+S^2}} \right\|. \quad (\text{Riesz metric})$$

Function  $f: [0, 1] \rightarrow \mathcal{CF}^{\text{sa}}$  continuous. Spectral flow  $\text{SF}(f)$  - “net number of eigenvalues crossing 0”.

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

Topologies on  $\mathcal{CF}^{\text{sa}}$ :

$$\rho_G(T, S) = \left\| \frac{T+i}{T-i} - \frac{S+i}{S-i} \right\|, \quad (\text{Graph metric})$$

$$\rho_R(T, S) = \left\| \frac{T}{\sqrt{1+T^2}} - \frac{S}{\sqrt{1+S^2}} \right\|. \quad (\text{Riesz metric})$$

Function  $f: [0, 1] \rightarrow \mathcal{CF}^{\text{sa}}$  continuous. Spectral flow  $\text{SF}(f)$  - “net number of eigenvalues crossing 0”.

- ★ Connections to the study of particle physics.

$\mathcal{H}$  separable Hilbert space,  $\mathcal{CF}^{\text{sa}}$  set of closed unbounded self-adjoint Fredholm operators on  $\mathcal{H}$ .

Topologies on  $\mathcal{CF}^{\text{sa}}$ :

$$\rho_G(T, S) = \left\| \frac{T+i}{T-i} - \frac{S+i}{S-i} \right\|, \quad (\text{Graph metric})$$

$$\rho_R(T, S) = \left\| \frac{T}{\sqrt{1+T^2}} - \frac{S}{\sqrt{1+S^2}} \right\|. \quad (\text{Riesz metric})$$

Function  $f: [0, 1] \rightarrow \mathcal{CF}^{\text{sa}}$  continuous. Spectral flow  $\text{SF}(f)$  - “net number of eigenvalues crossing 0”.

- ★ Connections to the study of particle physics.
- ★ Due to Atiyah and Singer in 1969 in [AS69] on the index theory of skew-adjoint Fredholm operators using topological language.

# Topological and geometric assumptions

Topological assumptions:

## Topological and geometric assumptions

Topological assumptions:

- ©  $\mathcal{M}$  smooth manifold (potentially non-compact),

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

Geometric assumptions:

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

Geometric assumptions:

- $\mathcal{M}$  carries a Riemannian structure via metric  $g$ ,

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

Geometric assumptions:

- ◊  $\mathcal{M}$  carries a Riemannian structure via metric  $g$ ,
- ◊  $(\mathcal{M}, g)$  is metric complete,

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

Geometric assumptions:

- ◊  $\mathcal{M}$  carries a Riemannian structure via metric  $g$ ,
- ◊  $(\mathcal{M}, g)$  is metric complete,
- ◊  $g$  has curvature bounds: there exists  $C_R < \infty$  such that  $|\text{Ric}_g| \leq C_R$  and  $|\nabla \text{Ric}_g| \leq C_R$ .

# Topological and geometric assumptions

Topological assumptions:

- ◎  $\mathcal{M}$  smooth manifold (potentially non-compact),
- ◎  $\Sigma = \partial\mathcal{M}$  smooth *compact* boundary (possibly  $= \emptyset$ ),
- ◎  $\mathcal{M}$  admits a Spin structure.

Geometric assumptions:

- ◊  $\mathcal{M}$  carries a Riemannian structure via metric  $g$ ,
- ◊  $(\mathcal{M}, g)$  is metric complete,
- ◊  $g$  has curvature bounds: there exists  $C_R < \infty$  such that  $|\text{Ric}_g| \leq C_R$  and  $|\nabla \text{Ric}_g| \leq C_R$ .

Obtain Spin bundle  $\Delta\mathcal{M} = P_{\text{Spin}}(\mathcal{M}) \times_{\eta} \Delta\mathbb{R}^n$ , and Atiyah-Singer Dirac operator  $D_g : C^\infty(\Delta\mathcal{M}) \rightarrow C^\infty(\Delta\mathcal{M})$ .

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1}|u|_g \leq |u|_h \leq C|u|_g, u \in T_x M \right\} \right).$$

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1}|u|_g \leq |u|_h \leq C|u|_g, u \in T_x M \right\} \right).$$

$\mathcal{M}$  without boundary and let  $h$  be another metric that is at least  $C^{0,1}$  such that  $\rho_M(g, h) < \infty$ .

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1} |u|_g \leq |u|_h \leq C |u|_g, u \in T_x M \right\} \right).$$

$\mathcal{M}$  without boundary and let  $h$  be another metric that is at least  $C^{0,1}$  such that  $\rho_M(g, h) < \infty$ .

$\not D_g$  and  $\not D_h$  - Atiyah-Singer Dirac operator of  $g$  and  $h$ .

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1} |u|_g \leq |u|_h \leq C |u|_g, u \in T_x M \right\} \right).$$

$M$  without boundary and let  $h$  be another metric that is at least  $C^{0,1}$  such that  $\rho_M(g, h) < \infty$ .

$\not D_g$  and  $\not D_h$  - Atiyah-Singer Dirac operator of  $g$  and  $h$ .

Via abstract theorem of Lesch in [Les05]:

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1}|u|_g \leq |u|_h \leq C|u|_g, u \in T_x M \right\} \right).$$

$M$  without boundary and let  $h$  be another metric that is at least  $C^{0,1}$  such that  $\rho_M(g, h) < \infty$ .

$\not D_g$  and  $\not D_h$  - Atiyah-Singer Dirac operator of  $g$  and  $h$ .

Via abstract theorem of Lesch in [Les05]:

$$\left\| \frac{\not D_g}{\sqrt{1 + \not D_g^2}} - \frac{\not D_h}{\sqrt{1 + \not D_h^2}} \right\|_{L^2 \rightarrow L^2} \lesssim \|\not D_g - \not D_h\|_{H^1 \rightarrow L^2}$$

## An example of a perturbation theorem

Another metric  $h$ , the distance to  $g$ :

$$\rho_M(g, h) = \log \left( \inf \left\{ C \geq 1 : C^{-1}|u|_g \leq |u|_h \leq C|u|_g, u \in T_x M \right\} \right).$$

$M$  without boundary and let  $h$  be another metric that is at least  $C^{0,1}$  such that  $\rho_M(g, h) < \infty$ .

$\not D_g$  and  $\not D_h$  - Atiyah-Singer Dirac operator of  $g$  and  $h$ .

Via abstract theorem of Lesch in [Les05]:

$$\left\| \frac{\not D_g}{\sqrt{1 + \not D_g^2}} - \frac{\not D_h}{\sqrt{1 + \not D_h^2}} \right\|_{L^2 \rightarrow L^2} \lesssim \|\not D_g - \not D_h\|_{H^1 \rightarrow L^2}$$

Continuity - requires  $\rho_M(g, h) \leq \varepsilon$  and  $\|\nabla^g h\| \leq \varepsilon$ .

# Metric perturbations

Theorem (B.-McIntosh-Rosén (2016) in [BMR18])

*Suppose  $(\mathcal{M}, g)$  be without boundary and there is a  $\kappa > 0$  with  $\text{inj}(\mathcal{M}, g) > \kappa$ . Fix  $C > 0$  and let  $h$  be a  $C^{0,1}$  metric with  $\rho_M(g, h) \leq 1$  and  $|\nabla^g h| \leq C$ .*

# Metric perturbations

Theorem (B.-McIntosh-Rosén (2016) in [BMR18])

Suppose  $(\mathcal{M}, g)$  be without boundary and there is a  $\kappa > 0$  with  $\text{inj}(\mathcal{M}, g) > \kappa$ . Fix  $C > 0$  and let  $h$  be a  $C^{0,1}$  metric with  $\rho_M(g, h) \leq 1$  and  $|\nabla^g h| \leq C$ . Then, we have the perturbation estimate

$$\left\| \frac{\mathbb{D}_g}{\sqrt{1 + \mathbb{D}_g^2}} - \frac{\mathbb{D}_h}{\sqrt{1 + \mathbb{D}_h^2}} \right\|_{L^2 \rightarrow L^2} \lesssim \rho_M(g, h),$$

where the implicit constant depends on  $C$ ,  $\dim \mathcal{M}$  and  $C_R$  and  $\kappa$ .

## Example

$\mathcal{M} = \mathbb{R}^2$  and  $g = \delta_{\mathbb{R}^2}$ , Euclidean metric.

## Example

$\mathcal{M} = \mathbb{R}^2$  and  $g = \delta_{\mathbb{R}^2}$ , Euclidean metric. For  $\varepsilon > 0$ ,

$$g_\varepsilon(x) = dx^1 \otimes dx^1 + \left(1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right)\right) dx^2 \otimes dx^2.$$

## Example

$\mathcal{M} = \mathbb{R}^2$  and  $g = \delta_{\mathbb{R}^2}$ , Euclidean metric. For  $\varepsilon > 0$ ,

$$g_\varepsilon(x) = dx^1 \otimes dx^1 + \left(1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right)\right) dx^2 \otimes dx^2.$$

As a matrix:

$$g_\varepsilon(x) = \begin{pmatrix} 1 & 0 \\ 0 & 1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right) \end{pmatrix}.$$

## Example

$\mathcal{M} = \mathbb{R}^2$  and  $g = \delta_{\mathbb{R}^2}$ , Euclidean metric. For  $\varepsilon > 0$ ,

$$g_\varepsilon(x) = dx^1 \otimes dx^1 + \left(1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right)\right) dx^2 \otimes dx^2.$$

As a matrix:

$$g_\varepsilon(x) = \begin{pmatrix} 1 & 0 \\ 0 & 1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right) \end{pmatrix}.$$

Have

$$\rho_{\mathcal{M}}(g, g_\varepsilon) \leq \varepsilon$$

## Example

$\mathcal{M} = \mathbb{R}^2$  and  $g = \delta_{\mathbb{R}^2}$ , Euclidean metric. For  $\varepsilon > 0$ ,

$$g_\varepsilon(x) = dx^1 \otimes dx^1 + \left(1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right)\right) dx^2 \otimes dx^2.$$

As a matrix:

$$g_\varepsilon(x) = \begin{pmatrix} 1 & 0 \\ 0 & 1 + \varepsilon \sin\left(\frac{|x|}{\varepsilon}\right) \end{pmatrix}.$$

Have

$$\rho_{\mathcal{M}}(g, g_\varepsilon) \leq \varepsilon$$

but

$$|\nabla^g g_\varepsilon(0)| = 1.$$

# Boundary conditions

Assume  $\Sigma \neq \emptyset$

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

*Local boundary condition* is  $\mathcal{B} = H^{\frac{1}{2}}(\mathcal{E})$ ,

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

*Local boundary condition* is  $\mathcal{B} = H^{\frac{1}{2}}(\mathcal{E})$ , where  $\mathcal{E}$  smooth subbundle of  $\mathcal{V}|_{\Sigma}$ .

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

*Local boundary condition* is  $\mathcal{B} = H^{\frac{1}{2}}(\mathcal{E})$ , where  $\mathcal{E}$  smooth subbundle of  $\mathcal{V}|_{\Sigma}$ .

$\pi_{\mathcal{E}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  and  $\pi_{\tilde{\mathcal{E}}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  orthogonal projectors.

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

*Local boundary condition* is  $\mathcal{B} = H^{\frac{1}{2}}(\mathcal{E})$ , where  $\mathcal{E}$  smooth subbundle of  $\mathcal{V}|_{\Sigma}$ .

$\pi_{\mathcal{E}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  and  $\pi_{\tilde{\mathcal{E}}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  orthogonal projectors. The *gap* between  $\mathcal{E}$  and  $\tilde{\mathcal{E}}$

$$\hat{\delta}_{\infty}(\mathcal{B}, \tilde{\mathcal{B}}) = \sup_{x \in \mathcal{M}} |\pi_{\mathcal{E}}(x) - \pi_{\tilde{\mathcal{E}}}(x)|.$$

## Boundary conditions

Assume  $\Sigma \neq \emptyset$ ,  $(\mathcal{V}, h)$  Riemannian vector bundle, and  $D$  first-order, elliptic, symmetric differential operator on  $(\mathcal{V}, h)$ .

*Local boundary condition* is  $\mathcal{B} = H^{\frac{1}{2}}(\mathcal{E})$ , where  $\mathcal{E}$  smooth subbundle of  $\mathcal{V}|_{\Sigma}$ .

$\pi_{\mathcal{E}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  and  $\pi_{\tilde{\mathcal{E}}} : \mathcal{V}|_{\Sigma} \rightarrow \mathcal{E}$  orthogonal projectors. The *gap* between  $\mathcal{E}$  and  $\tilde{\mathcal{E}}$

$$\hat{\delta}_{\infty}(\mathcal{B}, \tilde{\mathcal{B}}) = \sup_{x \in \mathcal{M}} |\pi_{\mathcal{E}}(x) - \pi_{\tilde{\mathcal{E}}}(x)|.$$

$D_{\mathcal{B}} = D$  with domain

$$\mathcal{D}(D_{\mathcal{B}}) = \{u \in \mathcal{D}(D_{\max}) : \mathcal{R} u \in \mathcal{B}\},$$

where  $\mathcal{R}$  is the trace map.

# Boundary value perturbations

Theorem (B.-Rosén (2017))

Let  $(\mathcal{M}, g)$  have  $\Sigma \neq \emptyset$  with a precompact neighbourhood  $Z$  of  $\Sigma$  and a  $\kappa > 0$  with  $\text{inj}(\mathcal{M} \setminus Z, g) > \kappa$ .

# Boundary value perturbations

Theorem (B.-Rosén (2017))

Let  $(\mathcal{M}, g)$  have  $\Sigma \neq \emptyset$  with a precompact neighbourhood  $Z$  of  $\Sigma$  and a  $\kappa > 0$  with  $\text{inj}(\mathcal{M} \setminus Z, g) > \kappa$ .

Fix  $C_B > 0$  and let  $\mathcal{B}$  and  $\tilde{\mathcal{B}}$  be two local self-adjoint  $\mathbb{D}$ -elliptic boundary conditions which satisfy:

# Boundary value perturbations

## Theorem (B.-Rosén (2017))

Let  $(\mathcal{M}, g)$  have  $\Sigma \neq \emptyset$  with a precompact neighbourhood  $Z$  of  $\Sigma$  and a  $\kappa > 0$  with  $\text{inj}(\mathcal{M} \setminus Z, g) > \kappa$ .

Fix  $C_B > 0$  and let  $\mathcal{B}$  and  $\tilde{\mathcal{B}}$  be two local self-adjoint  $\mathbb{D}$ -elliptic boundary conditions which satisfy:

$$(B1) \quad \|\mathcal{B}\|_{\text{Lip}} + \|\tilde{\mathcal{B}}\|_{\text{Lip}} \leq C_B, \text{ and}$$

# Boundary value perturbations

Theorem (B.-Rosén (2017))

Let  $(\mathcal{M}, g)$  have  $\Sigma \neq \emptyset$  with a precompact neighbourhood  $Z$  of  $\Sigma$  and a  $\kappa > 0$  with  $\text{inj}(\mathcal{M} \setminus Z, g) > \kappa$ .

Fix  $C_B > 0$  and let  $\mathcal{B}$  and  $\tilde{\mathcal{B}}$  be two local self-adjoint  $\mathbb{D}$ -elliptic boundary conditions which satisfy:

(B1)  $\|\mathcal{B}\|_{\text{Lip}} + \|\tilde{\mathcal{B}}\|_{\text{Lip}} \leq C_B$ , and

(B2)  $C_j \geq 1$  satisfying:

$$C_j^{-1} \|u\|_{\mathbb{D}^j, Z} \leq \|u\|_{H^j(Z, \Delta \mathcal{M})} \leq C_j \|u\|_{\mathbb{D}^j, Z}.$$

# Boundary value perturbations

Theorem (B.-Rosén (2017))

Let  $(\mathcal{M}, g)$  have  $\Sigma \neq \emptyset$  with a precompact neighbourhood  $Z$  of  $\Sigma$  and a  $\kappa > 0$  with  $\text{inj}(\mathcal{M} \setminus Z, g) > \kappa$ .

Fix  $C_B > 0$  and let  $\mathcal{B}$  and  $\tilde{\mathcal{B}}$  be two local self-adjoint  $\mathbb{D}$ -elliptic boundary conditions which satisfy:

(B1)  $\|\mathcal{B}\|_{\text{Lip}} + \|\tilde{\mathcal{B}}\|_{\text{Lip}} \leq C_B$ , and

(B2)  $C_j \geq 1$  satisfying:

$$C_j^{-1} \|u\|_{\mathbb{D}^j, Z} \leq \|u\|_{H^j(Z, \Delta \mathcal{M})} \leq C_j \|u\|_{\mathbb{D}^j, Z}.$$

Here,  $\|\mathcal{B}\|_{\text{Lip}} = \sup_{x \in \Sigma} (|\pi_{\mathcal{E}}(x)| + |\nabla \pi_{\mathcal{E}}(x)|)$ .

## Theorem (Cont...)

*Then, we have the perturbation estimate*

$$\left\| \frac{\not{D}_{\mathcal{B}}}{\sqrt{1 + \not{D}_{\mathcal{B}}^2}} - \frac{\not{D}_{\tilde{\mathcal{B}}}}{\sqrt{1 + \not{D}_{\tilde{\mathcal{B}}}^2}} \right\|_{L^2 \rightarrow L^2} \lesssim \hat{\delta}_\infty(\mathcal{B}, \tilde{\mathcal{B}}),$$

*where the implicit constant depends on  $\dim \mathcal{M}$ ,  $C_R$ ,  $\kappa$ ,  $Z$  and the constants in (B1)-(B2).*

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional.

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional. Then  $\Delta \mathcal{M} = \Delta^+ \mathcal{M} \overset{\perp}{\oplus} \Delta^- \mathcal{M}$  and

$$\not{D} = \begin{pmatrix} 0 & \not{D}^- \\ \not{D}^+ & 0 \end{pmatrix}.$$

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional. Then  $\Delta \mathcal{M} = \Delta^+ \mathcal{M} \overset{\perp}{\oplus} \Delta^- \mathcal{M}$  and

$$\mathbb{D} = \begin{pmatrix} 0 & \mathbb{D}^- \\ \mathbb{D}^+ & 0 \end{pmatrix}.$$

Let  $B \in \text{End}(\Delta^+ \Sigma)$  smooth and invertible,

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional. Then  $\Delta \mathcal{M} = \Delta^+ \mathcal{M} \overset{\perp}{\oplus} \Delta^- \mathcal{M}$  and

$$\not{D} = \begin{pmatrix} 0 & \not{D}^- \\ \not{D}^+ & 0 \end{pmatrix}.$$

Let  $B \in \text{End}(\Delta^+ \Sigma)$  smooth and invertible, define  $\Delta_B$  with fibres

$$\Delta_{B,x} \Sigma = \left\{ (\psi, \vec{n}(x) \cdot B(x)\psi) : \psi \in \Delta_x^+ \Sigma \right\}$$

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional. Then  $\Delta \mathcal{M} = \Delta^+ \mathcal{M} \overset{\perp}{\oplus} \Delta^- \mathcal{M}$  and

$$\not{D} = \begin{pmatrix} 0 & \not{D}^- \\ \not{D}^+ & 0 \end{pmatrix}.$$

Let  $B \in \text{End}(\Delta^+ \Sigma)$  smooth and invertible, define  $\Delta_B$  with fibres

$$\Delta_{B,x} \Sigma = \left\{ (\psi, \vec{n}(x) \cdot B(x)\psi) : \psi \in \Delta_x^+ \Sigma \right\}$$

Important class of local boundary conditions:

$$\mathcal{B}_B = H^{\frac{1}{2}}(\Delta_B \Sigma).$$

## Local boundary conditions in even dimensions

Assume  $\mathcal{M}$  even dimensional. Then  $\Delta \mathcal{M} = \Delta^+ \mathcal{M} \overset{\perp}{\oplus} \Delta^- \mathcal{M}$  and

$$\not{D} = \begin{pmatrix} 0 & \not{D}^- \\ \not{D}^+ & 0 \end{pmatrix}.$$

Let  $B \in \text{End}(\Delta^+ \Sigma)$  smooth and invertible, define  $\Delta_B$  with fibres

$$\Delta_{B,x} \Sigma = \left\{ (\psi, \vec{n}(x) \cdot B(x)\psi) : \psi \in \Delta_x^+ \Sigma \right\}$$

Important class of local boundary conditions:

$$\mathcal{B}_B = H^{\frac{1}{2}}(\Delta_B \Sigma).$$

$\not{D}$ -elliptic and self-adjoint whenever  $B(x)^* = B(x)$ .

# Part 2 - Boundary value problems for first-order elliptic operators

# Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

## Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

## Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

Induced Dirac operator

## Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

Induced Dirac operator

$$\not{D}_{\mathcal{W}} = \begin{pmatrix} \not{D} & \not{\sigma}^* \\ \not{\sigma} & \not{D}_{RS} \end{pmatrix}.$$

## Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

Induced Dirac operator

$$\not{D}_{\mathcal{W}} = \begin{pmatrix} \not{D} & \not{\sigma}^* \\ \not{\sigma} & \not{D}_{\text{RS}} \end{pmatrix}.$$

Natural questions:

# Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

Induced Dirac operator

$$\not{D}_{\mathcal{W}} = \begin{pmatrix} \not{D} & \not{\sigma}^* \\ \not{\sigma} & \not{D}_{\text{RS}} \end{pmatrix}.$$

Natural questions:

- ★ Boundary conditions - ellipticity, self-adjointness, regularity of eigensections?

# Motivation

$\mathcal{M}$  Riemannian Spin manifold, smooth compact boundary  $\Sigma$ ,

$$\mathcal{W} := T\mathcal{M} \otimes \Delta \mathcal{M} \cong \Delta \mathcal{M} \overset{\perp}{\oplus} \Delta_{\frac{3}{2}} \mathcal{M}.$$

Induced Dirac operator

$$\not{D}_{\mathcal{W}} = \begin{pmatrix} \not{D} & \not{\sigma}^* \\ \not{\sigma} & \not{D}_{\text{RS}} \end{pmatrix}.$$

Natural questions:

- ★ Boundary conditions - ellipticity, self-adjointness, regularity of eigensections?
- ★ Fredholmness and index theorems?

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathcal{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_{\Sigma}$ .

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathbb{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_\Sigma$ .

Principal symbol:

$$\sigma_{A_{RS}}(x, \xi) = \sigma_{\mathbb{D}_{RS}}(x, \tau(x))^{-1} \circ \sigma_{\mathbb{D}_{RS}}(x, \xi).$$

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathbb{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_\Sigma$ .

Principal symbol:

$$\sigma_{A_{RS}}(x, \xi) = \sigma_{\mathbb{D}_{RS}}(x, \tau(x))^{-1} \circ \sigma_{\mathbb{D}_{RS}}(x, \xi).$$

Fundamental assumption in the Bär-Ballmann framework [BB12]:

$A_{RS}$  is a symmetric operator.

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathcal{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_\Sigma$ .

Principal symbol:

$$\sigma_{A_{RS}}(x, \xi) = \sigma_{\mathcal{D}_{RS}}(x, \tau(x))^{-1} \circ \sigma_{\mathcal{D}_{RS}}(x, \xi).$$

Fundamental assumption in the Bär-Ballmann framework [BB12]:

$A_{RS}$  is a symmetric operator.

Boundary value problems for  $\mathcal{D}_{RS}$  “live” in:

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathcal{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_\Sigma$ .

Principal symbol:

$$\sigma_{A_{RS}}(x, \xi) = \sigma_{\mathcal{D}_{RS}}(x, \tau(x))^{-1} \circ \sigma_{\mathcal{D}_{RS}}(x, \xi).$$

Fundamental assumption in the Bär-Ballmann framework [BB12]:

$A_{RS}$  is a symmetric operator.

Boundary value problems for  $\mathcal{D}_{RS}$  “live” in:

$$\check{H}(A_{RS}) := \chi_{(-\infty, 0]}(A_{RS}) H^{\frac{1}{2}}(\Delta_{\frac{3}{2}} \Sigma) \bigoplus \chi_{(0, \infty)}(A_{RS}) H^{-\frac{1}{2}}(\Delta_{\frac{3}{2}} \Sigma),$$

## Adapted operator

Boundary adapted operator  $A_{RS}$  to  $\mathcal{D}_{RS}$  on  $\Delta_{\frac{3}{2}} \Sigma := \Delta_{\frac{3}{2}} \mathcal{M}|_\Sigma$ .

Principal symbol:

$$\sigma_{A_{RS}}(x, \xi) = \sigma_{\mathcal{D}_{RS}}(x, \tau(x))^{-1} \circ \sigma_{\mathcal{D}_{RS}}(x, \xi).$$

Fundamental assumption in the Bär-Ballmann framework [BB12]:

$A_{RS}$  is a symmetric operator.

Boundary value problems for  $\mathcal{D}_{RS}$  “live” in:

$$\check{H}(A_{RS}) := \chi_{(-\infty, 0]}(A_{RS}) H^{\frac{1}{2}}(\Delta_{\frac{3}{2}} \Sigma) \bigoplus \chi_{(0, \infty)}(A_{RS}) H^{-\frac{1}{2}}(\Delta_{\frac{3}{2}} \Sigma),$$

STOP Rarita-Schwinger  $\mathcal{D}_{RS}$  does not give rise to a symmetric  $A_{RS}$ .

## General setup

(A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;
- (A3)  $T$  is an interior vectorfield along  $\partial M$  and  $\tau$  the associated co-vectorfield;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;
- (A3)  $T$  is an interior vectorfield along  $\partial M$  and  $\tau$  the associated co-vectorfield;
- (A4)  $(\mathcal{E}, h^{\mathcal{E}})$ ,  $(\mathcal{F}, h^{\mathcal{F}}) \rightarrow \mathcal{M}$  are Hermitian vector bundles over  $\mathcal{M}$ ;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;
- (A3)  $T$  is an interior vectorfield along  $\partial M$  and  $\tau$  the associated co-vectorfield;
- (A4)  $(\mathcal{E}, h^{\mathcal{E}})$ ,  $(\mathcal{F}, h^{\mathcal{F}}) \rightarrow \mathcal{M}$  are Hermitian vector bundles over  $\mathcal{M}$ ;
- (A5)  $D : C^\infty(\mathcal{E}) \rightarrow C^\infty(\mathcal{F})$  is a first-order elliptic differential operator;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;
- (A3)  $T$  is an interior vectorfield along  $\partial M$  and  $\tau$  the associated co-vectorfield;
- (A4)  $(\mathcal{E}, h^{\mathcal{E}})$ ,  $(\mathcal{F}, h^{\mathcal{F}}) \rightarrow \mathcal{M}$  are Hermitian vector bundles over  $\mathcal{M}$ ;
- (A5)  $D : C^\infty(\mathcal{E}) \rightarrow C^\infty(\mathcal{F})$  is a first-order elliptic differential operator;
- (A6)  $D$  and  $D^*$  are complete;

## General setup

- (A1)  $\mathcal{M}$  is a manifold with compact boundary  $\Sigma = \partial\mathcal{M}$ ;
- (A2)  $\mu$  is a smooth volume measure on  $\mathcal{M}$  and  $\nu$  is the induced smooth volume measure on  $\Sigma$ ;
- (A3)  $T$  is an interior vectorfield along  $\partial M$  and  $\tau$  the associated co-vectorfield;
- (A4)  $(\mathcal{E}, h^{\mathcal{E}})$ ,  $(\mathcal{F}, h^{\mathcal{F}}) \rightarrow \mathcal{M}$  are Hermitian vector bundles over  $\mathcal{M}$ ;
- (A5)  $D : C^\infty(\mathcal{E}) \rightarrow C^\infty(\mathcal{F})$  is a first-order elliptic differential operator;
- (A6)  $D$  and  $D^*$  are complete;
- (A7)  $A$  and  $\tilde{A}$  are adapted differential operators on  $\mathcal{E}_\Sigma = \mathcal{E}|_{\partial M}$  and  $\mathcal{F}_\Sigma = \mathcal{F}|_{\partial M}$  respectively to the operators  $D$  and  $D^*$  respectively.

## Good coordinates

From [BB12], neighbourhood  $U$  of  $\Sigma$  such that there exists  $T > 0$  with

$$U \cong Z_{[0, T]} := [0, T] \times \Sigma$$

## Good coordinates

From [BB12], neighbourhood  $U$  of  $\Sigma$  such that there exists  $T > 0$  with

$$U \cong Z_{[0, T]} := [0, T] \times \Sigma$$

and

$$D = \sigma_t(\partial_t + A + R_t) : C^\infty(Z_{[0, T]}, \mathcal{E}) \rightarrow C^\infty(Z_{[0, T]}, \mathcal{F}),$$

## Good coordinates

From [BB12], neighbourhood  $U$  of  $\Sigma$  such that there exists  $T > 0$  with

$$U \cong Z_{[0, T]} := [0, T] \times \Sigma$$

and

$$D = \sigma_t(\partial_t + A + R_t) : C^\infty(Z_{[0, T]}, \mathcal{E}) \rightarrow C^\infty(Z_{[0, T]}, \mathcal{F}),$$

$\sigma_t = \sigma_D(x, dt(x))$  and  $R_t$  PDO of order 1

## Good coordinates

From [BB12], neighbourhood  $U$  of  $\Sigma$  such that there exists  $T > 0$  with

$$U \cong Z_{[0, T]} := [0, T] \times \Sigma$$

and

$$D = \sigma_t(\partial_t + A + R_t) : C^\infty(Z_{[0, T]}, \mathcal{E}) \rightarrow C^\infty(Z_{[0, T]}, \mathcal{F}),$$

$\sigma_t = \sigma_D(x, dt(x))$  and  $R_t$  PDO of order 1,

$$\|R_t u\|_{L^2(\Sigma)} \lesssim t \|A u\|_{L^2(\Sigma)} + \|u\|_{L^2(\Sigma)}.$$

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;
  - (ii)  $\exists C > 0$  such that for  $\zeta \in \mathbb{C} \setminus (S_\omega^o \cup B_R)$ , we have  $|\zeta| \| (A - \zeta I) \| \leq C$ ;

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;
  - (ii)  $\exists C > 0$  such that for  $\zeta \in \mathbb{C} \setminus (S_\omega^o \cup B_R)$ , we have  $|\zeta| \|A - \zeta I\| \leq C$ ;
  - (iii)  $\text{spec}(A)$  is a discrete set with no finite accumulation points.

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;
  - (ii)  $\exists C > 0$  such that for  $\zeta \in \mathbb{C} \setminus (S_\omega^o \cup B_R)$ , we have  $|\zeta| \|(\mathbf{A} - \zeta I)\| \leq C$ ;
  - (iii)  $\text{spec}(A)$  is a discrete set with no finite accumulation points.
- For  $r \in R$  such that  $l_r = \{\text{Re } \zeta = r\} \cap \text{spec}(A) = \emptyset$ ,  $A_r = A - r$  is an invertible  $\omega_r$  bi-sectorial operator.

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;
  - (ii)  $\exists C > 0$  such that for  $\zeta \in \mathbb{C} \setminus (S_\omega^o \cup B_R)$ , we have  $|\zeta| \| (A - \zeta I) \| \leq C$ ;
  - (iii)  $\text{spec}(A)$  is a discrete set with no finite accumulation points.
- For  $r \in R$  such that  $l_r = \{\text{Re } \zeta = r\} \cap \text{spec}(A) = \emptyset$ ,  $A_r = A - r$  is an invertible  $\omega_r$  bi-sectorial operator.
- Theorem of Grubb in [Gru12] (c.f. also Seeley in [See67]): spectral projectors  $\chi^\pm(A_r)$  are PDOs of order zero.

## Bi-sectorial boundary defining operators

- Via ellipticity,  $\text{spec}(\sigma_A(x, \xi)) \cap i\mathbb{R} = \emptyset, \forall (x, \xi) \in \Sigma \times T^*\Sigma$ .
- Theorem of Shubin in [Shu01]: the  $L^2$  spectrum of the operator  $A$  satisfies the following:
  - (i)  $\exists R > 0$  and  $\omega \in [0, \pi/2)$  such that  $\text{spec}(A) \subset S_\omega^o \cup B_R$ ;
  - (ii)  $\exists C > 0$  such that for  $\zeta \in \mathbb{C} \setminus (S_\omega^o \cup B_R)$ , we have  $|\zeta| \| (A - \zeta I) \| \leq C$ ;
  - (iii)  $\text{spec}(A)$  is a discrete set with no finite accumulation points.
- For  $r \in R$  such that  $l_r = \{\text{Re } \zeta = r\} \cap \text{spec}(A) = \emptyset$ ,  $A_r = A - r$  is an invertible  $\omega_r$  bi-sectorial operator.
- Theorem of Grubb in [Gru12] (c.f. also Seeley in [See67]): spectral projectors  $\chi^\pm(A_r)$  are PDOs of order zero.
- Define

$$\check{H}(A) = \chi^-(A) H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \quad \bigoplus \quad \chi^+(A) H^{-\frac{1}{2}}(\mathcal{E}_\Sigma).$$

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .
- $|A_r|$  is invertible  $\omega_r$ -sectorial.

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .
- $|A_r|$  is invertible  $\omega_r$ -sectorial.
- $\Psi$ DO differential calculus:  $\mathcal{D}(|A_r|) = \mathcal{D}(|A_r|^*)$ .

## McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .
- $|A_r|$  is invertible  $\omega_r$ -sectorial.
- $\Psi\text{DO}$  differential calculus:  $\mathcal{D}(|A_r|) = \mathcal{D}(|A_r|^*)$ .
- Theorem of Auscher-McIntosh-Nahmod from [AMN97]:  $|A_r|$  has an  $H^\infty$  functional calculus.

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .
- $|A_r|$  is invertible  $\omega_r$ -sectorial.
- $\Psi\text{DO}$  differential calculus:  $\mathcal{D}(|A_r|) = \mathcal{D}(|A_r|^*)$ .
- Theorem of Auscher-McIntosh-Nahmod from [AMN97]:  $|A_r|$  has an  $H^\infty$  functional calculus.
- Define  $H_D^1(\mathcal{E}) = \mathcal{D}(D_{\max}) \cap H_{\text{loc}}^1(\mathcal{E})$  with

# McIntoshery

- Let  $\text{sgn}(A_r) = \chi^+(A_r) - \chi^-(A_r)$ .
- Define  $|A_r| = A_r \text{sgn}(A_r)$ .
- $|A_r|$  is invertible  $\omega_r$ -sectorial.
- $\Psi\text{DO}$  differential calculus:  $\mathcal{D}(|A_r|) = \mathcal{D}(|A_r|^*)$ .
- Theorem of Auscher-McIntosh-Nahmod from [AMN97]:  $|A_r|$  has an  $H^\infty$  functional calculus.
- Define  $H_D^1(\mathcal{E}) = \mathcal{D}(D_{\max}) \cap H_{\text{loc}}^1(\mathcal{E})$  with

$$\|u\|_{H_D^1}^2 := \|\eta u\|_{H^1}^2 + \|Du\|^2 + \|u\|,$$

where  $\eta$  is a compactly supported cutoff near the boundary.

## Lemma

For  $\theta \in (\omega_r, \pi/2)$  fixed, there exists an inner product  $\langle \cdot, \cdot \rangle_{N,\theta}$  such that  $|A_r|$  is  $m$ - $\theta$ -accretive and for which the estimate

$$\begin{aligned} \|(\partial_t + A)u\|_{L^2(Z_{[0,\infty)})}^2 &\simeq \|u'\|_{L^2(Z_{[0,\infty)})}^2 + \|Au\|_{L^2(Z_{[0,\infty)})}^2 \\ &\quad - \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r) u_0, u_0 \rangle_{N,\theta} - r \|u_0\|_{N,\theta}^2, \end{aligned}$$

holds for  $u \in C_c^\infty(Z_{[0,\infty)}; E)$  where  $u_0 = u|_\Sigma$ .

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.
- (ii) Define  $\|\cdot\|_{N,\theta}^2 := \|\chi^+(A_r) \cdot\|_\theta^2 + \|\chi^-(A_r) \cdot\|_\theta^2$ .

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.
- (ii) Define  $\|\cdot\|_{N,\theta}^2 := \|\chi^+(A_r) \cdot\|_\theta^2 + \|\chi^-(A_r) \cdot\|_\theta^2$ .
- (iii)  $|A_r|$  is still  $m$ - $\theta$ -accretive with respect to  $\langle \cdot, \cdot \rangle_{N,\theta}$ .

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.
- (ii) Define  $\|\cdot\|_{N,\theta}^2 := \|\chi^+(A_r) \cdot\|_\theta^2 + \|\chi^-(A_r) \cdot\|_\theta^2$ .
- (iii)  $|A_r|$  is still  $m$ - $\theta$ -accretive with respect to  $\langle \cdot, \cdot \rangle_{N,\theta}$ .
- (iv)  $\chi^\pm(A_r)$  are self-adjoint in  $\langle \cdot, \cdot \rangle_{N,\theta}$  as is  $\operatorname{sgn}(A_r)$ .

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.
- (ii) Define  $\|\cdot\|_{N,\theta}^2 := \|\chi^+(A_r) \cdot\|_\theta^2 + \|\chi^-(A_r) \cdot\|_\theta^2$ .
- (iii)  $|A_r|$  is still  $m$ - $\theta$ -accretive with respect to  $\langle \cdot, \cdot \rangle_{N,\theta}$ .
- (iv)  $\chi^\pm(A_r)$  are self-adjoint in  $\langle \cdot, \cdot \rangle_{N,\theta}$  as is  $\operatorname{sgn}(A_r)$ .
- (v)  $a(u, v) = \langle |A_r|u, v \rangle_{N,\theta}$  is  $m$ - $\theta$ -accretive form.

For any equivalent norm  $\|\cdot\|_\star \simeq \|\cdot\|_{L^2(\mathcal{E}_\Sigma)}$ ,

$$\begin{aligned}\|(\partial_t + A)u\|_\star^2 &= \|u'\|_\star^2 + \|Au\|_\star^2 + \operatorname{Re} \langle |A_r| \operatorname{sgn}(A_r)u, u \rangle'_\star + r \langle u, u \rangle'_\star \\ &\quad + \operatorname{Re}(\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_\star - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_\star)\end{aligned}$$

- (i) Via Similarity Theorem due to Callier–Grabowski–Le Merdy (c.f. [Haa06] by Haase), there exists  $\langle \cdot, \cdot \rangle_\theta$  so that  $|A_r|$  is  $m$ - $\theta$ -accretive.
- (ii) Define  $\|\cdot\|_{N,\theta}^2 := \|\chi^+(A_r) \cdot\|_\theta^2 + \|\chi^-(A_r) \cdot\|_\theta^2$ .
- (iii)  $|A_r|$  is still  $m$ - $\theta$ -accretive with respect to  $\langle \cdot, \cdot \rangle_{N,\theta}$ .
- (iv)  $\chi^\pm(A_r)$  are self-adjoint in  $\langle \cdot, \cdot \rangle_{N,\theta}$  as is  $\operatorname{sgn}(A_r)$ .
- (v)  $a(u, v) = \langle |A_r|u, v \rangle_{N,\theta}$  is  $m$ - $\theta$ -accretive form.

$$\begin{aligned}\langle u', |A_r| \operatorname{sgn}(A_r)u \rangle_{N,\theta} - \langle |A_r| \operatorname{sgn}(A_r)u', u \rangle_{N,\theta} \\ = a(\operatorname{sgn}(A_r)u', u)^{\text{conj}} - a(u, \operatorname{sgn}(A_r)u') \in \operatorname{Im} \mathbb{R}\end{aligned}$$

## Theorem (Bär-B. (2018))

*The following hold:*

## Theorem (Bär-B. (2018))

*The following hold:*

- (i)  $C_c^\infty(\mathcal{E})$  is dense in  $\mathcal{D}(D_{\max})$  as well as  $\mathcal{D}((D^*)_{\max})$  with respect to the corresponding graph norms;

## Theorem (Bär-B. (2018))

*The following hold:*

- (i)  $C_c^\infty(\mathcal{E})$  is dense in  $\mathcal{D}(D_{\max})$  as well as  $\mathcal{D}((D^*)_{\max})$  with respect to the corresponding graph norms;
- (ii) the trace maps  $C_c^\infty(\mathcal{E}) \rightarrow C_c^\infty(\mathcal{E}_\Sigma)$  and  $C_c^\infty(\mathcal{F}) \rightarrow C_c^\infty(\mathcal{F}_\Sigma)$  given by  $u \mapsto u|_\Sigma$  extends uniquely to surjective bounded linear maps  $\mathcal{D}(D_{\max}) \rightarrow \check{H}(A)$  and  $\mathcal{D}((D^*)_{\max}) \rightarrow \check{H}(\tilde{A})$ ;

## Theorem (Bär-B. (2018))

*The following hold:*

- (i)  $C_c^\infty(\mathcal{E})$  is dense in  $\mathcal{D}(D_{\max})$  as well as  $\mathcal{D}((D^*)_{\max})$  with respect to the corresponding graph norms;
- (ii) the trace maps  $C_c^\infty(\mathcal{E}) \rightarrow C_c^\infty(\mathcal{E}_\Sigma)$  and  $C_c^\infty(\mathcal{F}) \rightarrow C_c^\infty(\mathcal{F}_\Sigma)$  given by  $u \mapsto u|_\Sigma$  extends uniquely to surjective bounded linear maps  $\mathcal{D}(D_{\max}) \rightarrow \check{H}(A)$  and  $\mathcal{D}((D^*)_{\max}) \rightarrow \check{H}(\tilde{A})$ ;
- (iii) The spaces

$$H_D^1 = \left\{ u \in \mathcal{D}(D_{\max}) : u|_\Sigma \in H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \right\}$$

$$H_{D^*}^1 = \left\{ u \in \mathcal{D}((D^*)_{\max}) : u|_\Sigma \in H^{\frac{1}{2}}(\mathcal{F}_\Sigma) \right\}; \quad \text{and}$$

## Theorem (Bär-B. (2018))

*The following hold:*

- (i)  $C_c^\infty(\mathcal{E})$  is dense in  $\mathcal{D}(D_{\max})$  as well as  $\mathcal{D}((D^*)_{\max})$  with respect to the corresponding graph norms;
- (ii) the trace maps  $C_c^\infty(\mathcal{E}) \rightarrow C_c^\infty(\mathcal{E}_\Sigma)$  and  $C_c^\infty(\mathcal{F}) \rightarrow C_c^\infty(\mathcal{F}_\Sigma)$  given by  $u \mapsto u|_\Sigma$  extends uniquely to surjective bounded linear maps  $\mathcal{D}(D_{\max}) \rightarrow \check{H}(A)$  and  $\mathcal{D}((D^*)_{\max}) \rightarrow \check{H}(\tilde{A})$ ;
- (iii) The spaces

$$H_D^1 = \left\{ u \in \mathcal{D}(D_{\max}) : u|_\Sigma \in H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \right\}$$

$$H_{D^*}^1 = \left\{ u \in \mathcal{D}((D^*)_{\max}) : u|_\Sigma \in H^{\frac{1}{2}}(\mathcal{F}_\Sigma) \right\}; \quad \text{and}$$

- (iv) for all  $u \in \mathcal{D}(D_{\max})$  and  $v \in \mathcal{D}((D^*)_{\max})$ ,

$$\langle D_{\max} u, v \rangle_{L^2(\mathcal{F})} - \langle u, (D^*)_{\max} v \rangle_{L^2(\mathcal{E})} = - \langle \sigma_0 u|_\Sigma, v|_\Sigma \rangle_{L^2(\mathcal{F}_\Sigma)}$$

## Higher regularity

Banach-valued Cauchy problem:  $f \in L^2(Z_{[0,\rho]}, E)$ ,

$$\partial_t W(t; f) + |A_r| W(t; f) = f(t), \quad \lim_{t \rightarrow 0} W(t; f) = 0.$$

## Higher regularity

Banach-valued Cauchy problem:  $f \in L^2(Z_{[0,\rho]}, E)$ ,

$$\partial_t W(t; f) + |A_r| W(t; f) = f(t), \quad \lim_{t \rightarrow 0} W(t; f) = 0.$$

Solution given by:

$$W(t; f) = \int_0^t e^{-(t-s)|A_r|} f(s) \, ds.$$

## Higher regularity

Banach-valued Cauchy problem:  $f \in L^2(Z_{[0,\rho]}, E)$ ,

$$\partial_t W(t; f) + |A_r| W(t; f) = f(t), \quad \lim_{t \rightarrow 0} W(t; f) = 0.$$

Solution given by:

$$W(t; f) = \int_0^t e^{-(t-s)|A_r|} f(s) \, ds.$$

Define:

$$\begin{aligned} S_{0,r} u(t) &= \int_0^t e^{-(t-s)|A_r|} \sigma_0^{-1} \chi^+(A_r) u(s) \, ds \\ &\quad - \int_t^\rho e^{-(s-t)|A_r|} \sigma_0^{-1} \chi^-(A_r) u(s) \, ds \end{aligned}$$

**Let**  $(C_\rho u)(s) = u(\rho - s)$ ,

(i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .
- (iii)  $D_{0,r}S_{0,r} = I$ , where  $D_{0,r} = \sigma_0(\partial_t + A_r)$ .

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .
- (iii)  $D_{0,r}S_{0,r} = I$ , where  $D_{0,r} = \sigma_0(\partial_t + A_r)$ .
- (iv)  $S_{0,r} : H^k(Z_{[0,\rho]}, E) \rightarrow H^{k+1}(Z_{[0,\rho]}, E)$  bounded.

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .
- (iii)  $D_{0,r}S_{0,r} = I$ , where  $D_{0,r} = \sigma_0(\partial_t + A_r)$ .
- (iv)  $S_{0,r} : H^k(Z_{[0,\rho]}, E) \rightarrow H^{k+1}(Z_{[0,\rho]}, E)$  bounded.
- (v)  $D_{0,r} : H^{k+1}(Z_{[0,\rho]}, E; B_0) \rightarrow H^k(Z_{[0,\rho]}, E)$  isomorphism, where  
 $B_0 = \chi^-(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \oplus \chi^+(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma)$

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .
- (iii)  $D_{0,r}S_{0,r} = I$ , where  $D_{0,r} = \sigma_0(\partial_t + A_r)$ .
- (iv)  $S_{0,r} : H^k(Z_{[0,\rho]}, E) \rightarrow H^{k+1}(Z_{[0,\rho]}, E)$  bounded.
- (v)  $D_{0,r} : H^{k+1}(Z_{[0,\rho]}, E; B_0) \rightarrow H^k(Z_{[0,\rho]}, E)$  isomorphism, where  
 $B_0 = \chi^-(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \oplus \chi^+(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma)$
- (vi)  $(I - S_{0,r}D_{0,r})u = e^{-t|A_r|}(\chi^+(A_r)u(0))$  whenever  
 $\chi^-(A_r)(u(\rho)) = 0$ ,

Let  $(C_\rho u)(s) = u(\rho - s)$ ,

- (i)  $S_{0,r}u(t) = W(t; \sigma_0^{-1}\chi^+(A_r)u) - W(\rho - t; \sigma_0^{-1}\chi^-(A_r)C_\rho u)$
- (ii)  $\chi^+(A_r)(S_{0,r}u)(0) = \chi^-(A_r)(S_{0,r}u)(\rho) = 0$ .
- (iii)  $D_{0,r}S_{0,r} = I$ , where  $D_{0,r} = \sigma_0(\partial_t + A_r)$ .
- (iv)  $S_{0,r} : H^k(Z_{[0,\rho]}, E) \rightarrow H^{k+1}(Z_{[0,\rho]}, E)$  bounded.
- (v)  $D_{0,r} : H^{k+1}(Z_{[0,\rho]}, E; B_0) \rightarrow H^k(Z_{[0,\rho]}, E)$  isomorphism, where  
 $B_0 = \chi^-(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma) \oplus \chi^+(A_r)H^{\frac{1}{2}}(\mathcal{E}_\Sigma)$
- (vi)  $(I - S_{0,r}D_{0,r})u = e^{-t|A_r|}(\chi^+(A_r)u(0))$  whenever  
 $\chi^-(A_r)(u(\rho)) = 0$ ,

Key estimate:

$$\int_0^\rho \|\partial_t W(t; f)\|_{L^2(\mathcal{E}_\Sigma)}^2 dt + \int_0^\rho \||A_r| W(t; f)\|_{L^2(\mathcal{E}_\Sigma)}^2 \lesssim \int_0^\rho \|f(t)\|_{L^2(\mathcal{E}_\Sigma)}^2.$$

## Theorem

*The following holds:*

$$\begin{aligned}\mathcal{D}(D_{\max}) \cap H_{loc}^{k+1}(\mathcal{E}) \\ = \{u \in \mathcal{D}(D_{\max}) : Du \in H_{loc}^k(\mathcal{F}) \text{ and} \\ \chi^+(A_r)(u|_{\Sigma}) \in H^{k+\frac{1}{2}}(\mathcal{E}_{\Sigma})\}.\end{aligned}$$

## Future/further directions

- ✿ Non-compact spatial  $\Sigma$ :

## Future/further directions

- ✿ Non-compact spatial  $\Sigma$ :  
    bounded geometry on  $\Sigma$ ;

## Future/further directions

- ✿ Non-compact spatial  $\Sigma$ :
  - bounded geometry on  $\Sigma$ ;
  - decay conditions at spatial infinity;

## Future/further directions

- ✿ Non-compact spatial  $\Sigma$ :
  - bounded geometry on  $\Sigma$ ;
  - decay conditions at spatial infinity;
- ✿ BVPs when  $\mathcal{M}$  is Lorentzian,  $\Sigma$  Riemannian and spatially compact,

## Future/further directions

- ✿ Non-compact spatial  $\Sigma$ :
  - bounded geometry on  $\Sigma$ ;
  - decay conditions at spatial infinity;
- ✿ BVPs when  $\mathcal{M}$  is Lorentzian,  $\Sigma$  Riemannian and spatially compact,
- ✿  $\Sigma$  is just a Lipschitz boundary.

# References I

- [AMN97] Pascal Auscher, Alan McIntosh, and Andrea Nahmod.  
**Holomorphic functional calculi of operators, quadratic estimates and interpolation.**  
*Indiana Univ. Math. J.*, 46(2):375–403, 1997.
- [AS69] M. F. Atiyah and I. M. Singer.  
**Index theory for skew-adjoint Fredholm operators.**  
*Inst. Hautes Études Sci. Publ. Math.*, (37):5–26, 1969.
- [BB12] Christian Bär and Werner Ballmann.  
**Boundary value problems for elliptic differential operators of first order.**  
17:1–78, 2012.
- [BMR18] Lashi Bandara, Alan McIntosh, and Andreas Rosén.  
**Riesz continuity of the atiyah–singer dirac operator under perturbations of the metric.**  
*Mathematische Annalen*, 370(1):863–915, Feb 2018.
- [Gru12] Gerd Grubb.  
**The sectorial projection defined from logarithms.**  
*Math. Scand.*, 111(1):118–126, 2012.

## References II

- [Haa06] Markus Haase.  
*The functional calculus for sectorial operators*, volume 169 of *Operator Theory: Advances and Applications*.  
Birkhäuser Verlag, Basel, 2006.
- [Les05] Matthias Lesch.  
**The uniqueness of the spectral flow on spaces of unbounded self-adjoint Fredholm operators.**  
In *Spectral geometry of manifolds with boundary and decomposition of manifolds*, volume 366 of *Contemp. Math.*, pages 193–224. Amer. Math. Soc., Providence, RI, 2005.
- [See67] R. T. Seeley.  
**Complex powers of an elliptic operator.**  
pages 288–307, 1967.
- [Shu01] M. A. Shubin.  
*Pseudodifferential operators and spectral theory*.  
Springer-Verlag, Berlin, second edition, 2001.  
Translated from the 1978 Russian original by Stig I. Andersson.