

The world of rough metrics

Lashi Bandara

Institut für Mathematik
Universität Potsdam

4 December 2020



Divergence form equations

Let $x \mapsto A(x) = (a_{ij}(x))$ symmetric, measurable matrix function.

Suppose $\kappa > 0$ and $\Lambda < \infty$ such that $x \in \mathcal{L}$ a.e.,

$$\kappa|u|_{\mathbb{R}^n}^2 \leq A(x)u \cdot u \leq \Lambda|u|_{\mathbb{R}^n}^2. \quad (1)$$

Then,

$$L_A = d^* A d = - \sum_{i,j=1}^n \partial_j(a_{ij}) \partial_i$$

non-negative self-adjoint operator on $L^2(\mathbb{R}^n, \mathcal{L})$.

- Lax-Milgram theorem circa 1954,
- *a priori estimates* of De Giorgi-Moser-Nash circa 1957:
if $u \in \text{dom}(L_A)$ such that $L_A u = 0 \implies u$ Hölder continuous.

Geometric perspective

Let g be smooth metric tensor on $\mathbb{R}^n \implies x \mapsto B(x)$ such that $g_x(u, v) = \delta(B(x)u, v) = B(x)u \cdot v$.

If g satisfies: $\exists C \geq 1$ s.t.

$$C^{-1}|u|_{\delta} \leq |u|_g \leq C|u|_{\delta},$$

then B satisfies (1).

Measure: $\mu_g = \sqrt{\det B} \mathcal{L}$.

Laplacian: $\Delta_g = d_g^* d$.

For $u \in \text{dom}(\Delta_g)$ and $v \in C_c^\infty(\mathbb{R}^n)$:

$$\begin{aligned}
 \langle \Delta_g u, v \rangle_{L^2(\mathbb{R}^n, \mu_g)} &= \int_{\mathbb{R}^n} g(du, \overline{dv}) \, d\mu_g \\
 &= \int_{\mathbb{R}^n} (Bdu) \cdot \overline{v} \, (\det B)^{\frac{1}{2}} \, d\mathcal{L} \\
 &= \int_{\mathbb{R}^n} ((\det B)^{\frac{1}{2}} Bdu) \cdot \overline{dv} \, d\mathcal{L} \\
 &= \int_{\mathbb{R}^n} d^{*,\delta}((\det B)^{\frac{1}{2}} B)du \, \overline{v} \, (\det B)^{-\frac{1}{2}} \, d\mu_g \\
 &= \left\langle (\det B)^{-\frac{1}{2}} d^{*,\delta}((\det B)^{\frac{1}{2}} B)du, v \right\rangle_{L^2(\mathbb{R}^n, \mu_g)} .
 \end{aligned}$$

I.e.

$$\Delta_g = (\det B)^{-\frac{1}{2}} d^{*,\delta}((\det B)^{\frac{1}{2}} B)d.$$

Kato's square root problem

(\mathcal{M}, g) Riemannian manifold.

Operator $L_{B, B_0} := d_g^* B d + B_0$, with \mathbb{C} -valued coefficients.

Assume: $\kappa_1 \leq B_0(x) \leq \kappa_2$, $B \in L^\infty(\mathcal{M}; \text{End}(T^*\mathcal{M}))$ and $x - \mu - \text{a.e.}$,

$$\text{Re } g_x(B(x)u, u) \geq \kappa |u|_{g_x}^2.$$

On $\mathcal{H} := L^2(\mathcal{M}) \oplus (L^2(\mathcal{M}) \oplus L^2(\mathcal{M}; T^*\mathcal{M}))$:

$$\Pi_g(B, B_0) = \begin{pmatrix} 0 & S^* X \\ S & 0 \end{pmatrix}, \quad \Pi_g(B, B_0)^2 = \begin{pmatrix} L_{B, B_0} & 0 \\ 0 & S S^* X \end{pmatrix}.$$

where $S : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M}) \oplus L^2(\mathcal{M}; T^*\mathcal{M})$ given by $Su = (u, du)$.

N.B. $\Pi_g(B, B_0)$ *not* self-adjoint - only ω -bisectorial for $\omega < \pi/2$.

Key estimate:

$$\|f(\Pi_g(B, B_0))\|_{L^2 \rightarrow L^2} \lesssim \|f\|_\infty. \quad (2)$$

for f bounded on a sector containing the spectrum of L_{B, B_0} , and holomorphic in the interior of the sector. I.e., $f(\zeta) = e^{-\zeta}$.

Implies: $\text{dom}(\sqrt{L_{B, B_0}}) = H^1(\mathcal{M}, g)$.

(Kato square root problem c.f [AHL⁺02, AKM06, BM16])

If h another metric on \mathcal{M} s.t. $\exists C \geq 1, C^{-1}|u|_g \leq |u|_h \leq C|u|_g$:

$$(2) \text{ holds for } g \iff (2) \text{ holds for } h.$$

Idea: $\Pi_h(B, B_0) = \Pi_g(\tilde{B}, \tilde{B}_0)$.

Why?

$$\|e^{-t(\Delta_g + I)} - e^{-t(\Delta_h + I)}\| \lesssim \|g - h\|_{L^\infty}.$$

Fix \mathcal{M} a smooth manifold. I.e. a smooth differentiable structure for $\mathcal{M} \rightsquigarrow$ exterior derivative d .

A measure structure for free:

- $A \subset \mathcal{M}$ *measurable* if for all charts (ψ, U) , $\psi(A \cap U) \subset \mathbb{R}^n$ \mathcal{L} -measurable.
- $A \subset \mathcal{M}$ *null measure* if for all charts (ψ, U) , $\mathcal{L}(\psi(A \cap U)) = 0$.

A measurable $\iff \mu_g$ -measurable for any smooth g .

A -null measure $\iff \mu_g(A) = 0$.

For any $\mathcal{V} \rightarrow \mathcal{M}$ vector bundle, we can talk about $\Gamma_R(\mathcal{V})$ - measurable sections of \mathcal{V} without a metric on \mathcal{M} .

Definition (Rough metric)

Let $g \in \Gamma_R(\text{Sym } T^*\mathcal{M} \otimes T^*\mathcal{M})$ such that: $\forall x \in \mathcal{M} \exists (U, \psi)$ chart around x and $\exists C \geq 1$ such that

$$C^{-1}|u|_{(\psi^*\delta)(y)} \leq |u|_{g(y)} \leq C|u|_{(\psi^*\delta)(y)},$$

$y - \text{a.e.} \in U$ and where δ is the Euclidean metric.

Say g is a *rough metric*.

Chart (U, ψ) satisfies the *local comparability condition*.

- Well-defined induced Radon measure via locally comparable charts:

$$d\mu_g(x) = \sqrt{\det g(x)} \, d\psi^*\mathcal{L}.$$

- A μ_g -measurable $\iff A$ measurable.
- $\mu_g(A) = 0 \iff A$ null-measure.

Lebesgue and Sobolev Spaces

Tensor bundle: $\mathcal{T}^{(p,q)}\mathcal{M} := (\otimes_{i=0}^p T^*\mathcal{M}) \otimes (\otimes_{i=0}^q T\mathcal{M})$.

Metric g extends to $\mathcal{T}^{(p,q)}\mathcal{M}$.

$$u \in L^p(\mathcal{T}^{(p,q)}\mathcal{M}, g) \text{ for } p \in (1, \infty) \iff \int_{\mathcal{M}} |u(x)|_{g(x)}^p d\mu_g(x) < \infty.$$

Similarly $u \in L^\infty(\mathcal{T}^{(p,q)}\mathcal{M}, g)$ if $\exists C < \infty$ such that $|u(x)|_{g(x)} \leq C$ x - a.e..

Operator $d_p = d : C^\infty \cap L^p(\mathcal{M}, g) \rightarrow C^\infty \cap L^p(\mathcal{M}; T^*\mathcal{M}, g)$ *closable* in $L^p(\mathcal{M}, g)$.

Define:

$$W^{1,p}(\mathcal{M}, g) := \text{dom}(\overline{d_p}), \quad W_0^{1,p}(\mathcal{M}, g) := \overline{C_c^\infty(\mathcal{M})}^{\|\cdot\|_{W^{1,p}}}.$$

See [Ban16].

Laplacian

Note: $H^1(\mathcal{M}, g) := W^{1,2}(\mathcal{M}, g)$ is a Hilbert space.

Define: $\mathcal{E}[u, v] := \langle du, dv \rangle_{L^2(\mathcal{M}, g)}$.

$\mathcal{W} \subset H^1(\mathcal{M}, g)$ closed subspace such that $H_0^1(\mathcal{M}, g) \subset \mathcal{W}$.

$\mathcal{E}_{\mathcal{W}} = \mathcal{E}$ with $\text{dom}(\mathcal{E}_{\mathcal{W}}) = \mathcal{W} \rightsquigarrow \Delta_{g, \mathcal{W}} := d_{\mathcal{W}}^{*,g} d_{\mathcal{W}}$.

Satisfies: $\text{dom}(\sqrt{\Delta_{g, \mathcal{W}}}) = \mathcal{W}$.

$\mathcal{W} = H^1(\mathcal{M}, g) \rightsquigarrow \Delta_N$ “Neumann Laplacian”,

$\mathcal{W} = H_0^1(\mathcal{M}, g) \rightsquigarrow \Delta_D$ “Dirichlet Laplacian”.

\mathcal{M} compact $\partial\mathcal{M} = \emptyset$: $W^{1,p}(\mathcal{M}, g) = W_0^{1,p}(\mathcal{M}, g) = W^{1,p}(\mathcal{M})$.

$H^1(\mathcal{M}, g) = H_0^1(\mathcal{M}, g) \iff \Delta_N = \Delta_D$.

☠ In general $\text{dom}(\Delta_D) \neq H^2(\mathcal{M})$. ☠

Examples

1. \mathcal{M} any smooth manifold, g a C^∞ complete Riemannian metric, $\exists \eta \in \mathbb{R}$ s.t. $\text{Ric}(g) \geq \eta g$. Then,

$$\Delta_N = \Delta_D = d_2^{*,g} \overline{d_2} = -\overline{\text{tr } \nabla^{T^* \mathcal{M}} d_2} = -\overline{\text{tr } \nabla^{T \mathcal{M}} \nabla} := \Delta,$$


and $\text{dom}(\Delta) = H_0^2(\mathcal{M}, g) = H^2(\mathcal{M}, g) = \{u \in L^2(\mathcal{M}) : u \in H^1(\mathcal{M}, g), \nabla^g du \in L^2(\mathcal{M}; \mathcal{T}^{(2,0)} \mathcal{M})\}$.

2. \mathcal{M} any smooth manifold, g a C^0 Riemannian metric $\implies g$ rough metric.
3. $\mathcal{M} = \Omega \subset \mathbb{R}^n$, bounded smooth domain, $g = \delta$. $\Delta_D = -\sum_{j=1}^n \partial_j^2$ with Dirichlet BCs, $\Delta_N = -\sum_{j=1}^n \partial_j^2$ with Neumann BCs.
4. $f : \mathbb{R}^n \rightarrow \mathbb{R}$ Lipschitz. Pullback metric $f^* \delta_{n+1}(u, v) = df u \cdot df v$ rough metric on \mathbb{R}^n .
5. \mathcal{M}, \mathcal{N} smooth manifolds, $\varphi : \mathcal{M} \rightarrow \mathcal{N}$ Diffeomorphism. If h a C^0 Riemannian metric on \mathcal{N} , $\varphi^* h$ rough metric on \mathcal{M} .

6. \mathcal{M} smooth, (ψ, U) chart such that $\psi(U) = B_1(0)$. For $\alpha \in (0, \pi]$, let

$$\varphi(x) = \left(\psi(x), \cot\left(\frac{\alpha}{2}\right) (1 - |\psi(x)|_{\mathbb{R}^n}) \right).$$


Suppose $g|_{\mathcal{M} \setminus U} \in C^\infty$ and $g|_U = \psi^* \delta_{n+1}$.



Then, g is a rough metric and $g = dr^2 + \sin^2(\alpha)r^2 dy^2$ in polar coordinates around x .

7. $\square^n = \partial[-1, 1]^{n+1}$ Euclidean cube.

$\varphi : \square^n \rightarrow S^n \subset \mathbb{R}^{n+1}$, radial projection $\varphi(x) := \frac{x}{|x|}$.



$d_{\square^n}(x, y) \simeq d_{S^n}(\varphi(x), \varphi(y)) \rightsquigarrow \varphi^{-1} : S^n \rightarrow \square^n$ Lipeomorphism.

$(S^n, (\varphi^{-1})^*(\delta|_{\square^n}))$ isometric to $\square^n \subset \mathbb{R}^{n+1}$.

$\exists B \in \mathbf{\Gamma}_R(\text{Sym End } (T^*S^n))$ such that

$(\varphi^{-1})^*(\delta|_{\square^n})_x(u, v) = g_{S^n_x}(Bu, v)$, x a.e.

$$\Delta_{\square^n} = d_{\square^n}^* d_{\square^n} = \varphi^*(\det B)^{-\frac{1}{2}} d^{*, S^n} ((\det B)^{\frac{1}{2}} B) d^{S^n} (\varphi^{-1})^*.$$

Weyl Asymptotics

Theorem (B.-Nursultanov-Rowlett 2018 [BNR20])

\mathcal{M} compact with smooth boundary $\partial\mathcal{M}$. $\mathcal{W} \subset H^1(\mathcal{M}, g)$ closed subspace, $H_0^1(\mathcal{M}, g) \subset \mathcal{W}$. Then,

- (i) $\Delta_{g, \mathcal{W}}$ has discrete non-negative spectrum with finite dimensional eigenspaces, and
- (ii) Letting $N(\lambda, \Delta_{g, \mathcal{W}})$ be the number of eigenvalues $\leq \lambda$,

$$\lim_{\lambda \rightarrow \infty} \frac{N(\lambda, \Delta_{g, \mathcal{W}})}{\lambda^{\frac{n}{2}}} = \frac{\omega_n}{(2\pi)^n} \mu_g(\mathcal{M}).$$

- $\text{dom}(\Delta_{g,\mathcal{W}}) \subset \text{dom}(\sqrt{\Delta_{g,\mathcal{W}}}) = \mathcal{W} \subset H^1(\mathcal{M}, g) = H^1(\mathcal{M})$.
- $\Delta_{g,\mathcal{W}}$ self-adjoint \implies
 $(\imath + \Delta_{g,\mathcal{W}})^{-1} : L^2(\mathcal{M}) \rightarrow \text{dom}(\Delta_{g,\mathcal{W}}) \subset H^1(\mathcal{M}) \xrightarrow{\text{compact}} L^2(\mathcal{M})$.
- Lack of distance: how to do domain monotonicity?
- Cover \mathcal{M} *almost-everywhere* by *mutually disjoint* Lipschitz domains in locally comparable charts (ψ, U) .
- $\exists B_\psi \in L^\infty(\text{Sym End}(T^*U))$ s.t. for all $v \in C_c^\infty(\mathring{U})$

$$\langle \Delta_{g,\mathcal{W}} u, v \rangle_{L^2(\mathcal{M},g)} = \left\langle B_\psi d^{\mathbb{R}^n} \psi^* u, d^{\mathbb{R}^n} v^* \right\rangle_{L^2(U; \sqrt{\det B_\psi} \, d\mathcal{L})}.$$

- Results of Birman-Solomjak [BS72] yield asymptotics in (ψ, U) for Dirichlet and Neumann problems of induced operator in $\varphi(U)$.
- Patch (carefully).

Heat equation

Setting so far: \mathcal{M} manifold, d from differentiable structure, g rough metric $\rightsquigarrow \mu_g$ and L^p , $W^{1,p}$. *No distance.*

$\mathcal{W} \subset H^1(\mathcal{M}, g)$ closed subspace, $H_0^1(\mathcal{M}, g) \subset \mathcal{W}$, and $C^\infty(\mathcal{M}) \cap \mathcal{W}$ dense in \mathcal{W} .

$d_{\mathcal{W}} = d$ with $\text{dom}(d_{\mathcal{W}}) = \mathcal{W} \rightsquigarrow$ Laplacian: $\Delta_{g, \mathcal{W}} := d_{\mathcal{W}}^{*, g} d_{\mathcal{W}}$.

$u \in C^1((0, \infty), \text{dom}(\Delta_{g, \mathcal{W}}))$ solution to the $\Delta_{g, \mathcal{W}}$ -heat equation with initial condition $u_0 \in L^2(\mathcal{M}, g)$ if:

- (i) $\partial_t u(\cdot, t) = \Delta_{g, \mathcal{W}} u(\cdot, t) \quad \forall t \in (0, \infty)$
- (ii) $\lim_{t \rightarrow 0} u(\cdot, t) = u_0$ in $L^2(\mathcal{M}, g)$.

Borel functional calculus \rightsquigarrow every such solution u uniquely given by:

$$u(\cdot, t) = e^{-t\Delta_{g, \mathcal{W}}} u_0.$$

Heat kernels

$(t, x, y) \mapsto \rho_t^{g, \mathcal{W}}(x, y) : (0, \infty) \times \mathcal{M} \times \mathcal{M}$ separably measurable, almost-everywhere symmetric in (x, y) is a heat kernel if:

- (i) $\lim_{t \rightarrow 0} \rho_t^{g, cW}(\cdot, y) = \delta_y$ (delta mass at y),
- (ii) if u solution to the heat equation with initial data $u_0 \in L^2(\mathcal{M}, g)$,

$$u(t, x) = \int_{\mathcal{M}} \rho_t^{g, \mathcal{W}}(x, y) u_0(y) d\mu_g(y).$$

Idea:

1. Show that for a.e. $x \in \mathcal{M}$, $\exists C_t < \infty$ s.t. for $v \in L^2(\mathcal{M}, g)$,

$$|(e^{-t\Delta_{g, \mathcal{W}}} v)(x)| \leq C_t \|v\|_{L^2}. \quad (3)$$

2. Implies $(v \mapsto e^{-t\Delta_{g, \mathcal{W}}} v)(x) \in L^2(\mathcal{M}, g)^*$. Riesz Representation theorem: $\exists a_{t,x} \in L^2(\mathcal{M}, g)$ such that $(e^{-t\Delta_{g, \mathcal{W}}} v)(x) = \langle a_{t,x}, v \rangle_{L^2}$.
3. Write $\rho_t^{g, \mathcal{W}}(x, y) := \langle a_{\frac{t}{2}, x}, a_{\frac{t}{2}, y} \rangle_{L^2}$.
4. Beurling-Deny condition $\|\sqrt{\Delta_{g, \mathcal{W}}} u\|_{L^2} \leq \|\sqrt{\Delta_{g, \mathcal{W}}} u\|_{L^2} \implies \rho_t^{g, \mathcal{W}} \geq 0$.

\mathcal{M} compact boundaryless $\implies \mathcal{W} = H_0^1(\mathcal{M}, g) = H^1(\mathcal{M}, g) \rightsquigarrow$
unique Δ_g .

Fix h smooth auxiliary metric. $\exists B \in L^\infty(\mathcal{M}; \text{Sym End}(T^*\mathcal{M}), h)$ s.t.
 $g(u, v) = h(Bu, v)$. Then,

$$\Delta_g = -\theta^{-1} d^{*,h}(B\theta) \bar{d}.$$

$\exists \eta \in \mathbb{R}$ s.t. $\text{Ric}(h) \geq \eta h$. Saloff-Coste in [SC92] \implies parabolic
Harnack estimates for $u \geq 0$ satisfying

$$\partial_t u = -\theta^{-1} d^{*,h}(B\theta) \bar{d} u = \Delta_g u.$$

Implies (3), and $\exists \alpha > 0$ s.t.

$$(t, x, y) \mapsto \rho_t^g(x, y) \in C^\omega((0, \infty); C^\alpha(\mathcal{M} \times \mathcal{M})).$$

See [Ban17].

Example: $g = \varphi^* h$, h smooth, $\varphi : \mathcal{M} \rightarrow \mathcal{M}$ Lipeomorphism.

$$(x, y) \mapsto \rho_t^g(x, y) = \rho_t^h(\varphi(x), \varphi(y)) \in C^{0,1}(\mathcal{M} \times \mathcal{M}).$$

Theorem (B.-Bryan 2019 [BB20])

For \mathcal{M} smooth manifold, g rough metric, \mathcal{W} subspace as before, there exists a unique heat kernel $\rho_t^{g, \mathcal{W}}$ satisfying:

- (i) $\rho_t^{g, \mathcal{W}} > 0$ for $t > 0$,
- (ii) $\forall K \in \mathcal{M}, \forall 0 < t_1 < t_2, \exists \alpha(K, t_1, t_2)$ such that

$$(t, x, y) \mapsto \rho_t^{g, \mathcal{W}} C^{\omega} \left([t_1, t_2]; C^{\alpha(K, t_1, t_2)}(K \times K) \right).$$

1. Cover \mathcal{M} by locally comparable charts. Inside (U, ψ) :
 $\exists B_{\psi} \in L^{\infty}(U; \text{Sym End}(T^* \mathcal{M}))$ such that $\forall v \in C_c^{\infty}(U)$,

$$\langle \Delta_{g, \mathcal{W}} u, v \rangle_{L^2(\mathcal{M}, g)} = \left\langle B_{\psi} d^{\mathbb{R}^n}(\psi^* u), d^{\mathbb{R}^n}(\psi^* v) \right\rangle_{L^2(\psi(U), \sqrt{\det B_{\psi}} \mathcal{L})}.$$

2. Parabolic Harnack estimates in *weighted* Sobolev spaces
 $H^1(\psi(U), \sqrt{\det B_{\psi}} \mathcal{L}) \implies (3)$ and regularity.

Varadhan asymptotics

Important paper: [Nor97] by Norris. Setting: \mathcal{M} Lipschitz.

Abstract methods for existence of $\rho_t^{g, \mathcal{W}}$ when $\mathcal{W} = H_0^1(\mathcal{M}, g)$ or $H^1(\mathcal{M}, g)$.

Distance:

$$\mathbf{d}_g(x, y) := \sup \{ f(x) - f(y) : f \in C^{0,1}(\mathcal{M}), |df(x)|_g \leq 1 \text{ x a.e.} \}.$$

Important theorem:

$$\lim_{t \rightarrow 0} 4t \log \rho_t^{g, \mathcal{W}}(x, y) = -\mathbf{d}_g^2(x, y).$$

$\rightsquigarrow (\mathcal{M}, \mathbf{d}_g, \mu_g)$ measure metric space, infinitesimally Hilbertian.

Question: Synthetic curvature properties in terms of g ?

Strum's example of non-uniqueness

Let $\mathcal{M} = \mathbb{R}^n$. In [Stu97] by Sturm, shows
 $\exists A \in L^\infty(\mathbb{R}^n; \text{Sym Mat}(n))$ s.t. for x a.e.

$$\frac{1}{2}|u|_{\mathbb{R}^n}^2 \leq A(x)u \cdot u < |u|_{\mathbb{R}^n}^2$$

- $g(u, v) := A(x)u \cdot v$ rough metric on \mathbb{R}^n ,
- $d\mu_g(x) = \sqrt{\det A(x)} d\mathcal{L} < d\mathcal{L}$.

But

$$\mathbf{d}_g(x, y) = |x - y| \implies \mathcal{H}^{\mathbf{d}_g} \neq \mu_g.$$

Cannot happen for $A \in C^0 \cap L^\infty(\mathbb{R}^n; \text{Sym Mat}(n))$.

Future outlook

Current works in “progress”:

- Study of Weyl asymptotics on \mathcal{M} with boundary, rough metric, for certain *Robin* boundary conditions (with Medet Nursultanov and Julie Rowlett).
- (\mathcal{M}, g) automatically RCD when \mathcal{M} compact. Synthetic bound in terms of g ? (with Chiara Rigoni).

Questions:

1. $(\mathcal{M}, g) \rightsquigarrow (\mathcal{M}, \mathbf{d}_g, \mu_g)$. Synthetic curvature properties?
2. Notions of convergence for $(\mathcal{M}_i, g_i) \rightarrow (\mathcal{M}_\infty, g_\infty)$?

3. Given g, h , suppose $\exists C(g, h) \geq 1$ such that for x a.e.,

$$C(g, h)^{-1} |u|_{g(x)} \leq |u|_{h(x)} \leq C(g, h) |u|_{g(x)}.$$

Recall $Su = (u, du)$ and

$$\Pi_g(B, B_0) = \begin{pmatrix} 0 & S^*X \\ S & 0 \end{pmatrix}, \quad \Pi_g(B, B_0)^2 = \begin{pmatrix} L_{B, B_0} & 0 \\ 0 & SS^*X \end{pmatrix}.$$

$\Pi_g(B, B_0)$ first-order factorisation of $L_{B, B_0} = d_2^{*,g} B \overline{d_2}$.

$$\|f(\Pi_g(B, B_0))\| \lesssim \|f\|_\infty \iff \|f(\Pi_h(B, B_0))\| \lesssim \|f\|_\infty.$$

Holy grail:

$$\|f(\Pi_{g, B, B_0})\| \lesssim \|f\|_\infty \stackrel{?}{\implies} \text{curvature bound on } (\mathcal{M}, \mathbf{d}_g, \mu_g).$$



- [AHL⁺02] Pascal Auscher, Steve Hofmann, Michael Lacey, Alan McIntosh, and Philippe Tchamitchian.
The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n .
Ann. of Math. (2), 156(2):633–654, 2002.
- [AKM06] Andreas Axelsson, Stephen Keith, and Alan McIntosh.
Quadratic estimates and functional calculi of perturbed Dirac operators.
Invent. Math., 163(3):455–497, 2006.
- [Ban16] Lashi Bandara.
Rough metrics on manifolds and quadratic estimates.
Mathematische Zeitschrift, 283(3-4):1245–1281, 2016.
- [Ban17] Lashi Bandara.
Continuity of solutions to space-varying pointwise linear elliptic equations.
Publicacions Matemàtiques, 61(1):239–258, 2017.
- [BB20] Lashi Bandara and Paul Bryan.
Heat kernels and regularity for rough metrics on smooth manifolds.
Mathematische Zeitschrift, to appear, 2020.
- [BM16] Lashi Bandara and Alan McIntosh.
The Kato Square Root Problem on Vector Bundles with Generalised Bounded Geometry.
Journal of Geometric Analysis, 26(1):428–462, 2016.
- [BNR20] Lashi Bandara, Medet Nursultanov, and Julie Rowlett.
Eigenvalue asymptotics for weighted Laplace equations on rough Riemannian manifolds with boundary.
Annali della Scuola Normale Superiore di Pisa. Classe di Scienze, to appear, 2020.
- [BS72] M. Sh. Birman and M. Z. Solomjak.
Spectral asymptotics of nonsmooth elliptic operators. I, II.
Trudy Moskov. Mat. Obšč., 27:3–52; *ibid.* 28 (1973), 3–34, 1972.
- [Nor97] James R. Norris.
Heat kernel asymptotics and the distance function in Lipschitz Riemannian manifolds.
Acta Math., 179(1):79–103, 1997.
- [SC92] Laurent Saloff-Coste.
Uniformly elliptic operators on Riemannian manifolds.
J. Differential Geom., 36(2):417–450, 1992.
- [Stu97] Karl-Theodor Sturm.
Is a diffusion process determined by its intrinsic metric?
Chaos Solitons Fractals, 8(11):1855–1860, 1997.