

Geometry and the Kato square root problem

Lashi Bandara

Centre for Mathematics and its Applications
Australian National University

7 June 2013

Geometric Analysis Seminar
University of Wollongong

Outline

- Brief overview of the Kato square root problem on \mathbb{R}^n .

Outline

- Brief overview of the Kato square root problem on \mathbb{R}^n .
- A motivating application to hyperbolic PDE.

Outline

- Brief overview of the Kato square root problem on \mathbb{R}^n .
- A motivating application to hyperbolic PDE.
- Recent progress on the Kato square root problem on smooth manifolds by McIntosh and B.

Outline

- Brief overview of the Kato square root problem on \mathbb{R}^n .
- A motivating application to hyperbolic PDE.
- Recent progress on the Kato square root problem on smooth manifolds by McIntosh and B.
- Recent progress on subelliptic Kato square root problems on Lie groups by ter Elst, McIntosh, and B.

Outline

- Brief overview of the Kato square root problem on \mathbb{R}^n .
- A motivating application to hyperbolic PDE.
- Recent progress on the Kato square root problem on smooth manifolds by McIntosh and B.
- Recent progress on subelliptic Kato square root problems on Lie groups by ter Elst, McIntosh, and B.
- Kato square root problem on smooth manifolds with non-smooth metrics, connection to geometric flows and PDEs.

The Kato square root problem

Let $A \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^N))$ and $a \in L^\infty(\mathbb{R}^n)$.

The Kato square root problem

Let $A \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^N))$ and $a \in L^\infty(\mathbb{R}^n)$. Suppose that there exists $\kappa_1, \kappa_2 > 0$ such that for all $u \in W^{1,2}(\mathbb{R}^n)$,

$$\operatorname{Re} a(x) \geq \kappa_1 \quad \text{and} \quad \operatorname{Re} \langle A \nabla u, \nabla u \rangle \geq \kappa_2 \|u\|^2.$$

The Kato square root problem

Let $A \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^N))$ and $a \in L^\infty(\mathbb{R}^n)$. Suppose that there exists $\kappa_1, \kappa_2 > 0$ such that for all $u \in W^{1,2}(\mathbb{R}^n)$,

$$\operatorname{Re} a(x) \geq \kappa_1 \quad \text{and} \quad \operatorname{Re} \langle A \nabla u, \nabla u \rangle \geq \kappa_2 \|u\|^2.$$

The Kato square root problem on \mathbb{R}^n is the statement that

$$\begin{aligned} \mathcal{D}(\sqrt{-a \operatorname{div} A \nabla}) &= W^{1,2}(\mathbb{R}^n) \\ \|\sqrt{-a \operatorname{div} A \nabla} u\| &\simeq \|\nabla u\|. \end{aligned} \tag{K1}$$

The Kato square root problem

Let $A \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^N))$ and $a \in L^\infty(\mathbb{R}^n)$. Suppose that there exists $\kappa_1, \kappa_2 > 0$ such that for all $u \in W^{1,2}(\mathbb{R}^n)$,

$$\operatorname{Re} a(x) \geq \kappa_1 \quad \text{and} \quad \operatorname{Re} \langle A \nabla u, \nabla u \rangle \geq \kappa_2 \|u\|^2.$$

The Kato square root problem on \mathbb{R}^n is the statement that

$$\begin{aligned} \mathcal{D}(\sqrt{-a \operatorname{div} A \nabla}) &= W^{1,2}(\mathbb{R}^n) \\ \|\sqrt{-a \operatorname{div} A \nabla} u\| &\simeq \|\nabla u\|. \end{aligned} \tag{K1}$$

This was answered in the positive in 2002 by Pascal Auscher, Steve Hofmann, Michael Lacey, Alan McIntosh and Phillipe Tchamitchian in [AHLMcT].

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} .

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$.

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$. In general, this is not true by a counterexample of McIntosh in 1972 in [Mc72].

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$. In general, this is not true by a counterexample of McIntosh in 1972 in [Mc72].
- A second related question is the following.

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$. In general, this is not true by a counterexample of McIntosh in 1972 in [Mc72].
- A second related question is the following. Suppose that J_t is a family of *closed, densely-defined, Hermitian* forms on \mathcal{H} with domain \mathcal{W} and $L(t)$ the associated self-adjoint operators to J_t with domain \mathcal{W} .

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$. In general, this is not true by a counterexample of McIntosh in 1972 in [Mc72].
- A second related question is the following. Suppose that J_t is a family of *closed, densely-defined, Hermitian* forms on \mathcal{H} with domain \mathcal{W} and $L(t)$ the associated self-adjoint operators to J_t with domain \mathcal{W} . If $t \mapsto J_t$ extends to holomorphic family (for small z), then is $\partial_t \sqrt{L(t)} : \mathcal{V} \rightarrow \mathcal{H}$ a bounded operator?

- If further $A = A^*$, (K1) is a trivial consequence of the Lax-Milgram Theorem.
- Solution to (K1) implies that $\mathcal{D}(\sqrt{-\operatorname{div} A \nabla}) = \mathcal{D}(\sqrt{-\operatorname{div} A^* \nabla})$.
We can ask a more abstract question for *accretive* operators L on a Hilbert space \mathcal{H} . There, the question is whether $\mathcal{D}(\sqrt{L^*}) = \mathcal{D}(\sqrt{L})$. In general, this is not true by a counterexample of McIntosh in 1972 in [Mc72].
- A second related question is the following. Suppose that J_t is a family of *closed, densely-defined, Hermitian* forms on \mathcal{H} with domain \mathcal{W} and $L(t)$ the associated self-adjoint operators to J_t with domain \mathcal{W} . If $t \mapsto J_t$ extends to holomorphic family (for small z), then is $\partial_t \sqrt{L(t)} : \mathcal{V} \rightarrow \mathcal{H}$ a bounded operator?
- Counterexample to this second question by McIntosh in 1982 in [Mc82].

Motivations from PDE

For $k = 1, 2$, let $L_k = -\operatorname{div} A_k \nabla$ where $A_k \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^n))$ non-negative self-adjoint and L_k uniformly elliptic.

Motivations from PDE

For $k = 1, 2$, let $L_k = -\operatorname{div} A_k \nabla$ where $A_k \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^n))$ non-negative self-adjoint and L_k uniformly elliptic.

As aforementioned, $\mathcal{D}(\sqrt{L_k}) = W^{1,2}(\mathbb{R}^n)$ and $\|\sqrt{L_k}u\| \simeq \|\nabla u\|$ for $u \in W^{1,2}(\mathbb{R}^n)$.

Motivations from PDE

For $k = 1, 2$, let $L_k = -\operatorname{div} A_k \nabla$ where $A_k \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^n))$ non-negative self-adjoint and L_k uniformly elliptic.

As aforementioned, $\mathcal{D}(\sqrt{L_k}) = W^{1,2}(\mathbb{R}^n)$ and $\|\sqrt{L_k}u\| \simeq \|\nabla u\|$ for $u \in W^{1,2}(\mathbb{R}^n)$.

Let u_k be solutions to the wave equation with respect to L_k with the same initial data.

Motivations from PDE

For $k = 1, 2$, let $L_k = -\operatorname{div} A_k \nabla$ where $A_k \in L^\infty(\mathbb{R}^n, \mathcal{L}(\mathbb{C}^n))$ non-negative self-adjoint and L_k uniformly elliptic.

As aforementioned, $\mathcal{D}(\sqrt{L_k}) = W^{1,2}(\mathbb{R}^n)$ and $\|\sqrt{L_k}u\| \simeq \|\nabla u\|$ for $u \in W^{1,2}(\mathbb{R}^n)$.

Let u_k be solutions to the wave equation with respect to L_k with the same initial data. That is,

$$\begin{aligned}\partial_t^2 u_k + L_k u_k &= 0 \\ \partial_t u_k|_{t=0} &= g \in L^2(\mathbb{R}^n) \\ u_k(0) &= f \in W^{1,2}(\mathbb{R}^n).\end{aligned}$$

Suppose there exists a $C > 0$

$$\|\sqrt{L_1}u - \sqrt{L_2}u\| \leq C\|A_1 - A_2\|_\infty\|\nabla u\|. \quad (\text{P})$$

Suppose there exists a $C > 0$

$$\|\sqrt{L_1}u - \sqrt{L_2}u\| \leq C\|A_1 - A_2\|_\infty \|\nabla u\|. \quad (\text{P})$$

Then, whenever $t > 0$, the following estimate holds:

$$\begin{aligned} \|u_1(t) - u_2(t)\| + \left\| \int_0^t \nabla(u_1(s) - u_2(s)) \, ds \right\| \\ \leq Ct\|A_1 - A_2\|_\infty (\|\nabla f\| + \|g\|). \end{aligned}$$

Suppose there exists a $C > 0$

$$\|\sqrt{L_1}u - \sqrt{L_2}u\| \leq C\|A_1 - A_2\|_\infty \|\nabla u\|. \quad (\text{P})$$

Then, whenever $t > 0$, the following estimate holds:

$$\begin{aligned} \|u_1(t) - u_2(t)\| + \left\| \int_0^t \nabla(u_1(s) - u_2(s)) \, ds \right\| \\ \leq Ct\|A_1 - A_2\|_\infty (\|\nabla f\| + \|g\|). \end{aligned}$$

See [Aus].

The estimate (P) is related to the second question of Kato.

The estimate (P) is related to the second question of Kato.

By solving the Kato square root problem (K1) for *complex* coefficients A , we are able to automatically obtain (P) from (K1).

Kato square root problem on manifolds

Let \mathcal{M} be a smooth, complete Riemannian manifold with metric g , Levi-Civita connection ∇ , and volume measure μ .

Kato square root problem on manifolds

Let \mathcal{M} be a smooth, complete Riemannian manifold with metric g , Levi-Civita connection ∇ , and volume measure μ .

Write $\operatorname{div} = -\nabla^*$ in L^2 and let $S = (\operatorname{I}, \nabla)$.

Kato square root problem on manifolds

Let \mathcal{M} be a smooth, complete Riemannian manifold with metric g , Levi-Civita connection ∇ , and volume measure μ .

Write $\operatorname{div} = -\nabla^*$ in L^2 and let $S = (\mathbf{I}, \nabla)$.

Assume $a \in L^\infty(\mathcal{M})$ and $A = (A_{ij}) \in L^\infty(\mathcal{M}, \mathcal{L}(L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M})))$.

Kato square root problem on manifolds

Let \mathcal{M} be a smooth, complete Riemannian manifold with metric g , Levi-Civita connection ∇ , and volume measure μ .

Write $\operatorname{div} = -\nabla^*$ in L^2 and let $S = (I, \nabla)$.

Assume $a \in L^\infty(\mathcal{M})$ and $A = (A_{ij}) \in L^\infty(\mathcal{M}, \mathcal{L}(L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M})))$.

Consider the following second order differential operator

$L_A : \mathcal{D}(L_A) \subset L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$ defined by

$$L_A u = a S^* A S u = -a \operatorname{div}(A_{11} \nabla u) - a \operatorname{div}(A_{10} u) + a A_{01} \nabla u + a A_{00} u.$$

The main theorem on manifolds

Theorem (B.-Mc, 2012)

Let \mathcal{M} be a smooth, complete Riemannian manifold with $|\text{Ric}| \leq C$ and $\text{inj}(\mathcal{M}) \geq \kappa > 0$. Suppose the following ellipticity condition holds: there exists $\kappa_1, \kappa_2 > 0$ such that

$$\begin{aligned}\text{Re} \langle av, v \rangle &\geq \kappa_1 \|v\|^2 \\ \text{Re} \langle ASu, Su \rangle &\geq \kappa_2 \|u\|_{W^{1,2}}^2\end{aligned}$$

for $v \in L^2(\mathcal{M})$ and $u \in W^{1,2}(\mathcal{M})$. Then, $\mathcal{D}(\sqrt{L_A}) = \mathcal{D}(\nabla) = W^{1,2}(\mathcal{M})$ and $\|\sqrt{L_A}u\| \simeq \|\nabla u\| + \|u\| = \|u\|_{W^{1,2}}$ for all $u \in W^{1,2}(\mathcal{M})$.

Lipschitz estimates

Since we allow the coefficients a and A to be *complex*, we obtain the following stability result as a consequence:

Theorem (B.-Mc, 2012)

Let \mathcal{M} be a smooth, complete Riemannian manifold with $|\text{Ric}| \leq C$ and $\text{inj}(\mathcal{M}) \geq \kappa > 0$. Suppose that there exist $\kappa_1, \kappa_2 > 0$ such that $\text{Re} \langle av, v \rangle \geq \kappa_1 \|v\|^2$ and $\text{Re} \langle ASu, Su \rangle \geq \kappa_2 \|u\|_{W^{1,2}}^2$ for $v \in L^2(\mathcal{M})$ and $u \in W^{1,2}(\mathcal{M})$. Then for every $\eta_i < \kappa_i$, whenever $\|\tilde{a}\|_\infty \leq \eta_1$, $\|\tilde{A}\|_\infty \leq \eta_2$, the estimate

$$\|\sqrt{L_A} u - \sqrt{L_{A+\tilde{A}}} u\| \lesssim (\|\tilde{a}\|_\infty + \|\tilde{A}\|_\infty) \|u\|_{W^{1,2}}$$

holds for all $u \in W^{1,2}(\mathcal{M})$. The implicit constant depends in particular on A, a and η_i .

The Hodge-Dirac operator

Let $\Omega(\mathcal{M})$ denote the algebra of differential forms over \mathcal{M} under the exterior product \wedge .

The Hodge-Dirac operator

Let $\Omega(\mathcal{M})$ denote the algebra of differential forms over \mathcal{M} under the exterior product \wedge .

Let d be the exterior derivative as an operator on $L^2(\Omega(\mathcal{M}))$ and d^* its adjoint, both of which are *nilpotent* operators.

The Hodge-Dirac operator

Let $\Omega(\mathcal{M})$ denote the algebra of differential forms over \mathcal{M} under the exterior product \wedge .

Let d be the exterior derivative as an operator on $L^2(\Omega(\mathcal{M}))$ and d^* its adjoint, both of which are *nilpotent* operators.

The Hodge-Dirac operator is then the self-adjoint operator $D = d + d^*$.

The Hodge-Dirac operator

Let $\Omega(\mathcal{M})$ denote the algebra of differential forms over \mathcal{M} under the exterior product \wedge .

Let d be the exterior derivative as an operator on $L^2(\Omega(\mathcal{M}))$ and d^* its adjoint, both of which are *nilpotent* operators.

The Hodge-Dirac operator is then the self-adjoint operator $D = d + d^*$.
The Hodge-Laplacian is then $D^2 = d d^* + d^* d$.

The Hodge-Dirac operator

Let $\Omega(\mathcal{M})$ denote the algebra of differential forms over \mathcal{M} under the exterior product \wedge .

Let d be the exterior derivative as an operator on $L^2(\Omega(\mathcal{M}))$ and d^* its adjoint, both of which are *nilpotent* operators.

The Hodge-Dirac operator is then the self-adjoint operator $D = d + d^*$. The Hodge-Laplacian is then $D^2 = d d^* + d^* d$.

For an invertible $A \in L^\infty(\mathcal{L}(\Omega(\mathcal{M})))$, we consider perturbing D to obtain the operator $D_A = d + A^{-1}d^*A$.

Curvature endomorphism for forms

Let $\{\theta^i\}$ be an orthonormal frame at x for $\Omega^1(\mathcal{M}) = T^*\mathcal{M}$.

Curvature endomorphism for forms

Let $\{\theta^i\}$ be an orthonormal frame at x for $\Omega^1(\mathcal{M}) = T^*\mathcal{M}$.

Denote the components of the curvature tensor in this frame by Rm_{ijkl} .

Curvature endomorphism for forms

Let $\{\theta^i\}$ be an orthonormal frame at x for $\Omega^1(\mathcal{M}) = T^*\mathcal{M}$.

Denote the components of the curvature tensor in this frame by Rm_{ijkl} .
The curvature endomorphism is then the operator

$$R\omega = Rm_{ijkl} \theta^i \wedge (\theta^j \lrcorner (\theta^k \wedge (\theta^l \lrcorner \omega)))$$

for $\omega \in \Omega_x(\mathcal{M})$.

Curvature endomorphism for forms

Let $\{\theta^i\}$ be an orthonormal frame at x for $\Omega^1(\mathcal{M}) = T^*\mathcal{M}$.

Denote the components of the curvature tensor in this frame by Rm_{ijkl} .
The curvature endomorphism is then the operator

$$\mathbf{R}\omega = \text{Rm}_{ijkl} \theta^i \wedge (\theta^j \lrcorner (\theta^k \wedge (\theta^l \lrcorner \omega)))$$

for $\omega \in \Omega_x(\mathcal{M})$.

This can be seen as an extension of Ricci curvature for forms, since $g(\mathbf{R}\omega, \eta) = \text{Ric}(\omega^b, \eta^b)$ whenever $\omega, \eta \in \Omega_x^1(\mathcal{M})$ and where $b : T^*\mathcal{M} \rightarrow T\mathcal{M}$ is the flat isomorphism through the metric g .

Curvature endomorphism for forms

Let $\{\theta^i\}$ be an orthonormal frame at x for $\Omega^1(\mathcal{M}) = T^*\mathcal{M}$.

Denote the components of the curvature tensor in this frame by Rm_{ijkl} .
The curvature endomorphism is then the operator

$$R\omega = \text{Rm}_{ijkl} \theta^i \wedge (\theta^j \lrcorner (\theta^k \wedge (\theta^l \lrcorner \omega)))$$

for $\omega \in \Omega_x(\mathcal{M})$.

This can be seen as an extension of Ricci curvature for forms, since $g(R\omega, \eta) = \text{Ric}(\omega^b, \eta^b)$ whenever $\omega, \eta \in \Omega_x^1(\mathcal{M})$ and where $b : T^*\mathcal{M} \rightarrow T\mathcal{M}$ is the flat isomorphism through the metric g .

The Weitzenböck formula then asserts that $D^2 = \text{tr}_{12} \nabla^2 + R$.

Theorem (B., 2012)

Let \mathcal{M} be a smooth, complete Riemannian manifold and let $\beta \in \mathbb{C} \setminus \{0\}$. Suppose there exist $\eta, \kappa > 0$ such that $|\text{Ric}| \leq \eta$ and $\text{inj}(\mathcal{M}) \geq \kappa$. Furthermore, suppose there is a $\zeta \in \mathbb{R}$ satisfying $g(\text{R} u, u) \geq \zeta |u|^2$, for $u \in \Omega_x(\mathcal{M})$ and $A \in L^\infty(\mathcal{L}(\Omega(\mathcal{M})))$ and $\kappa_1 > 0$ satisfying

$$\text{Re} \langle Au, u \rangle \geq \kappa_1 \|u\|^2.$$

Then, $\mathcal{D}(\sqrt{D_A^2 + |\beta|^2}) = \mathcal{D}(D_A) = \mathcal{D}(d) \cap \mathcal{D}(d^* A)$ and $\|\sqrt{D_A^2 + |\beta|^2} u\| \simeq \|D_A u\| + \|u\|$.

Lie groups

Let \mathcal{G} be a Lie group of dimension n with Lie algebra \mathfrak{g} and equipped with the left-invariant Haar measure μ .

Lie groups

Let \mathcal{G} be a Lie group of dimension n with Lie algebra \mathfrak{g} and equipped with the left-invariant Haar measure μ .

We say that a linearly independent $\mathfrak{a} = \{a_1, \dots, a_k\} \subset \mathfrak{g}$ is an *algebraic basis* if we can recover a basis for \mathfrak{g} through multi-commutation.

Lie groups

Let \mathcal{G} be a Lie group of dimension n with Lie algebra \mathfrak{g} and equipped with the left-invariant Haar measure μ .

We say that a linearly independent $\mathfrak{a} = \{a_1, \dots, a_k\} \subset \mathfrak{g}$ is an *algebraic basis* if we can recover a basis for \mathfrak{g} through multi-commutation.

Let A_i denote the right-translation of a_i and $A^i = A_i^*$. Let $\text{span}\{A_1, \dots, A_k\} = \mathcal{A} \subset T\mathcal{G}$ be the bundle obtained through the right-translation of \mathfrak{a} and $\mathcal{A}^* = \{A^1, \dots, A^k\}$ the dual of \mathcal{A} .

Subelliptic distance

Theorem of Carathéodory-Chow tells us that for any two points $x, y \in \mathcal{G}$, we can find an absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathcal{G}$ such that

$$\dot{\gamma}(t) = \sum_i \dot{\gamma}^i(t) A_i(\gamma(t)) \in \mathcal{A}.$$

Subelliptic distance

Theorem of Carathéodory-Chow tells us that for any two points $x, y \in \mathcal{G}$, we can find an absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathcal{G}$ such that

$$\dot{\gamma}(t) = \sum_i \dot{\gamma}^i(t) A_i(\gamma(t)) \in \mathcal{A}.$$

The length of such a curve then is given by

$$\ell(\gamma) = \int_0^1 \left(\sum_i |\dot{\gamma}^i(t)|^2 \right)^{\frac{1}{2}} dt$$

Subelliptic distance

Theorem of Carathéodory-Chow tells us that for any two points $x, y \in \mathcal{G}$, we can find an absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathcal{G}$ such that

$$\dot{\gamma}(t) = \sum_i \dot{\gamma}^i(t) A_i(\gamma(t)) \in \mathcal{A}.$$

The length of such a curve then is given by

$$\ell(\gamma) = \int_0^1 \left(\sum_i |\dot{\gamma}^i(t)|^2 \right)^{\frac{1}{2}} dt$$

Define distance $d(x, y)$ as the infimum over the length of all such curves.

Subelliptic distance

Theorem of Carathéodory-Chow tells us that for any two points $x, y \in \mathcal{G}$, we can find an absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathcal{G}$ such that

$$\dot{\gamma}(t) = \sum_i \dot{\gamma}^i(t) A_i(\gamma(t)) \in \mathcal{A}.$$

The length of such a curve then is given by

$$\ell(\gamma) = \int_0^1 \left(\sum_i |\dot{\gamma}^i(t)|^2 \right)^{\frac{1}{2}} dt$$

Define distance $d(x, y)$ as the infimum over the length of all such curves.

The measure μ is Borel-regular with respect to d .

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Each vector field A_i is a skew-adjoint differential operator.

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Each vector field A_i is a skew-adjoint differential operator. We consider it as a unbounded operator on $L^2(\mathcal{G})$ with domain $\mathcal{D}(A_i)$.

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Each vector field A_i is a skew-adjoint differential operator. We consider it as a unbounded operator on $L^2(\mathcal{G})$ with domain $\mathcal{D}(A_i)$.

By also considering ∇ as a closed, densely-defined operator on $L^2(\mathcal{M})$, we obtain the first-order Sobolev space $W^{1,2}(\mathcal{G})' = \mathcal{D}(\nabla) = \cap_{i=1}^k \mathcal{D}(A_i)$.

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Each vector field A_i is a skew-adjoint differential operator. We consider it as a unbounded operator on $L^2(\mathcal{G})$ with domain $\mathcal{D}(A_i)$.

By also considering ∇ as a closed, densely-defined operator on $L^2(\mathcal{M})$, we obtain the first-order Sobolev space $W^{1,2}(\mathcal{G})' = \mathcal{D}(\nabla) = \cap_{i=1}^k \mathcal{D}(A_i)$.

We write the divergence as $\operatorname{div} = -\nabla^*$.

Subelliptic operators

For $f \in C^\infty(\mathcal{G})$, define

$$\nabla f = A_i f A^i.$$

This defines an *sub-connection* on $C^\infty(\mathcal{M})$.

Each vector field A_i is a skew-adjoint differential operator. We consider it as a unbounded operator on $L^2(\mathcal{G})$ with domain $\mathcal{D}(A_i)$.

By also considering ∇ as a closed, densely-defined operator on $L^2(\mathcal{M})$, we obtain the first-order Sobolev space $W^{1,2}(\mathcal{G})' = \mathcal{D}(\nabla) = \cap_{i=1}^k \mathcal{D}(A_i)$.

We write the divergence as $\operatorname{div} = -\nabla^*$. Then, the subelliptic Laplacian associated to \mathcal{A} is

$$\Delta = -\operatorname{div} \nabla = -\sum_{i=1}^k A_i^2.$$

Nilpotent Lie groups

The Lie group \mathcal{G} is *nilpotent* if the inductively defined sequence $\mathfrak{g}_1 = [\mathfrak{g}, \mathfrak{g}]$, $\mathfrak{g}_2 = [\mathfrak{g}_1, \mathfrak{g}]$, \dots is eventually zero.

Nilpotent Lie groups

The Lie group \mathcal{G} is *nilpotent* if the inductively defined sequence $\mathfrak{g}_1 = [\mathfrak{g}, \mathfrak{g}]$, $\mathfrak{g}_2 = [\mathfrak{g}_1, \mathfrak{g}]$, \dots is eventually zero.

Theorem (B.-E.-Mc., 2012)

Let (\mathcal{G}, d, μ) be a connected, nilpotent Lie group with \mathfrak{a} an algebraic basis, d the associated sub-elliptic distance, and μ the left Haar measure. Suppose that $a, A \in L^\infty$ and that there exist $\kappa_1, \kappa_2 > 0$ satisfying

$$\operatorname{Re} \langle av, v \rangle \geq \kappa_1 \|v\|^2, \quad \text{and} \quad \operatorname{Re} \langle A \nabla u, \nabla u \rangle \geq \kappa_2 \|\nabla u\|^2.$$

for every $v \in L^2(\mathcal{G})$ and $u \in W^{1,2}(\mathcal{G})'$. Then, $\mathcal{D}(\sqrt{-a \operatorname{div} A \nabla}) = W^{1,2}(\mathcal{G})'$ and $\|\sqrt{-a \operatorname{div} A \nabla} u\| \simeq \|\nabla u\|$ for $u \in W^{1,2}(\mathcal{G})'$.

General Lie groups

Let $S = (I, \nabla)$ as in the manifold case.

General Lie groups

Let $S = (I, \nabla)$ as in the manifold case.

Theorem (B.-E.-Mc., 2012)

Let (\mathcal{G}, d, μ) be a connected Lie group, \mathfrak{a} an algebraic basis, d the associated sub-elliptic distance, and μ the left Haar measure. Let $a, A \in L^\infty$ such that

$$\operatorname{Re} \langle av, v \rangle \geq \kappa_1 \|v\|^2, \quad \text{and} \quad \operatorname{Re} \langle ASu, Su \rangle \geq \kappa_2 \|u\|_{W^{1,2}'}$$

*for every $v \in L^2(\mathcal{G})$ and $u \in W^{1,2}(\mathcal{G})'$. Then, $\mathcal{D}(\sqrt{aS^*AS}) = W^{1,2}(\mathcal{G})'$ with $\|\sqrt{aS^*AS}u\| \simeq \|u\|_{W^{1,2}'} = \|u\| + \|\nabla u\|$.*

Operator theory

We adapt the framework due to Axelsson (Rosén), Keith, McIntosh in [AKMc].

Operator theory

We adapt the framework due to Axelsson (Rosén), Keith, McIntosh in [AKMc].

Let \mathcal{H} be a Hilbert space and $\Gamma : \mathcal{H} \rightarrow \mathcal{H}$ a closed, densely-defined, nilpotent operator.

Operator theory

We adapt the framework due to Axelsson (Rosén), Keith, McIntosh in [AKMc].

Let \mathcal{H} be a Hilbert space and $\Gamma : \mathcal{H} \rightarrow \mathcal{H}$ a closed, densely-defined, nilpotent operator.

Suppose that $B_1, B_2 \in \mathcal{L}(\mathcal{H})$ such that there exist $\kappa_1, \kappa_2 > 0$ satisfying

$$\operatorname{Re} \langle B_1 u, u \rangle \geq \kappa_1 \|u\|^2 \quad \text{and} \quad \operatorname{Re} \langle B_2 v, v \rangle \geq \kappa_2 \|v\|^2$$

for $u \in \mathcal{R}(\Gamma^*)$ and $v \in \mathcal{R}(\Gamma)$.

Operator theory

We adapt the framework due to Axelsson (Rosén), Keith, McIntosh in [AKMc].

Let \mathcal{H} be a Hilbert space and $\Gamma : \mathcal{H} \rightarrow \mathcal{H}$ a closed, densely-defined, nilpotent operator.

Suppose that $B_1, B_2 \in \mathcal{L}(\mathcal{H})$ such that there exist $\kappa_1, \kappa_2 > 0$ satisfying

$$\operatorname{Re} \langle B_1 u, u \rangle \geq \kappa_1 \|u\|^2 \quad \text{and} \quad \operatorname{Re} \langle B_2 v, v \rangle \geq \kappa_2 \|v\|^2$$

for $u \in \mathcal{R}(\Gamma^*)$ and $v \in \mathcal{R}(\Gamma)$.

Furthermore, suppose the operators B_1, B_2 satisfy $B_1 B_2 \mathcal{R}(\Gamma) \subset \mathcal{N}(\Gamma)$ and $B_2 B_1 \mathcal{R}(\Gamma^*) \subset \mathcal{N}(\Gamma^*)$.

Operator theory

We adapt the framework due to Axelsson (Rosén), Keith, McIntosh in [AKMc].

Let \mathcal{H} be a Hilbert space and $\Gamma : \mathcal{H} \rightarrow \mathcal{H}$ a closed, densely-defined, nilpotent operator.

Suppose that $B_1, B_2 \in \mathcal{L}(\mathcal{H})$ such that there exist $\kappa_1, \kappa_2 > 0$ satisfying

$$\operatorname{Re} \langle B_1 u, u \rangle \geq \kappa_1 \|u\|^2 \quad \text{and} \quad \operatorname{Re} \langle B_2 v, v \rangle \geq \kappa_2 \|v\|^2$$

for $u \in \mathcal{R}(\Gamma^*)$ and $v \in \mathcal{R}(\Gamma)$.

Furthermore, suppose the operators B_1, B_2 satisfy $B_1 B_2 \mathcal{R}(\Gamma) \subset \mathcal{N}(\Gamma)$ and $B_2 B_1 \mathcal{R}(\Gamma^*) \subset \mathcal{N}(\Gamma^*)$.

The primary operator we consider is $\Pi_B = \Gamma + B_1 \Gamma^* B_2$.

If the quadratic estimates

$$\int_0^\infty \|t\Pi_B(1+t^2\Pi_B^2)^{-1}u\|^2 \simeq \|u\| \quad (\text{Q})$$

hold for every $u \in \overline{\mathcal{R}(\Pi_B)}$, then, \mathcal{H} decomposes into the spectral subspaces of Π_B as $\mathcal{H} = \mathcal{N}(\Pi_B) \oplus E_+ \oplus E_-$ and

$$\begin{aligned} \mathcal{D}(\sqrt{\Pi_B^2}) &= \mathcal{D}(\Pi_B) = \mathcal{D}(\Gamma) \cap \mathcal{D}(\Gamma^* B_2) \\ \|\sqrt{\Pi_B^2}u\| &\simeq \|\Pi_B u\| \simeq \|\Gamma u\| + \|\Gamma^* B_2 u\|. \end{aligned}$$

If the quadratic estimates

$$\int_0^\infty \|t\Pi_B(1+t^2\Pi_B^2)^{-1}u\|^2 \simeq \|u\| \quad (\text{Q})$$

hold for every $u \in \overline{\mathcal{R}(\Pi_B)}$, then, \mathcal{H} decomposes into the spectral subspaces of Π_B as $\mathcal{H} = \mathcal{N}(\Pi_B) \oplus E_+ \oplus E_-$ and

$$\begin{aligned} \mathcal{D}(\sqrt{\Pi_B^2}) &= \mathcal{D}(\Pi_B) = \mathcal{D}(\Gamma) \cap \mathcal{D}(\Gamma^* B_2) \\ \|\sqrt{\Pi_B^2}u\| &\simeq \|\Pi_B u\| \simeq \|\Gamma u\| + \|\Gamma^* B_2 u\|. \end{aligned}$$

The Kato problems are then obtained by letting $\mathcal{H} = L^2(\mathcal{M}) \oplus (L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M}))$ and letting

$$\Gamma = \begin{pmatrix} 0 & 0 \\ S & 0 \end{pmatrix}, \quad \Gamma^* = \begin{pmatrix} 0 & S^* \\ 0 & 0 \end{pmatrix}, \quad B_1 = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}, \quad B_2 = \begin{pmatrix} 0 & 0 \\ 0 & A \end{pmatrix}.$$

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

The idea is to reduce the quadratic estimate (Q) to a *Carleson measure* estimate.

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

The idea is to reduce the quadratic estimate (Q) to a *Carleson measure* estimate. This is achieved via a *local $T(b)$* argument.

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

The idea is to reduce the quadratic estimate (Q) to a *Carleson measure* estimate. This is achieved via a *local $T(b)$* argument.

Geometry enters the picture precisely in the harmonic analysis.

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

The idea is to reduce the quadratic estimate (Q) to a *Carleson measure* estimate. This is achieved via a *local $T(b)$* argument.

Geometry enters the picture precisely in the harmonic analysis. We need to perform harmonic analysis on vector fields, not just functions.

Geometry and harmonic analysis

Harmonic analytic methods are used to prove quadratic estimates (Q).

The idea is to reduce the quadratic estimate (Q) to a *Carleson measure* estimate. This is achieved via a *local $T(b)$* argument.

Geometry enters the picture precisely in the harmonic analysis. We need to perform harmonic analysis on vector fields, not just functions.

One can show that this is *not* artificial - the Kato problem on functions immediately provides a solution to the dual problem on vector fields.

Elements of the proofs

Similar in structure to the proof of [AKMc] which is inspired from the proof in [AHLMcT].

Elements of the proofs

Similar in structure to the proof of [AKMc] which is inspired from the proof in [AHLMcT].

- A dyadic decomposition of the space

Elements of the proofs

Similar in structure to the proof of [AKMc] which is inspired from the proof in [AHLMcT].

- A dyadic decomposition of the space
- A notion of averaging (in an integral sense)

Elements of the proofs

Similar in structure to the proof of [AKMc] which is inspired from the proof in [AHLMcT].

- A dyadic decomposition of the space
- A notion of averaging (in an integral sense)
- Poincaré inequality - on both functions and vector fields

Elements of the proofs

Similar in structure to the proof of [AKMc] which is inspired from the proof in [AHLMcT].

- A dyadic decomposition of the space
- A notion of averaging (in an integral sense)
- Poincaré inequality - on both functions and vector fields
- Control of ∇^2 in terms of Δ .

The case of non-smooth metrics on manifolds

We let \mathcal{M} be a smooth, complete manifold as before but now let g be a C^0 metric. Let μ_g denote the volume measure with respect to g .

The case of non-smooth metrics on manifolds

We let \mathcal{M} be a smooth, complete manifold as before but now let g be a C^0 metric. Let μ_g denote the volume measure with respect to g .

Let $h \in C^0(\mathcal{T}^{(2,0)}\mathcal{M})$. Then, define

$$\|h\|_{\text{op},g} = \sup_{x \in \mathcal{M}} \sup_{|u|_g = |v|_g = 1} |h_x(u, v)|.$$

If \tilde{g} is another C^0 metric satisfying $\|g - \tilde{g}\|_{\text{op},g} \leq \delta < 1$, then $L^2(\mathcal{M}, g) = L^2(\mathcal{M}, \tilde{g})$ and $W^{1,2}(\mathcal{M}, g) = W^{1,2}(\mathcal{M}, \tilde{g})$ with comparable norms.

Π_B under a change of metric

The operator Γ_g does not change under the change of metric.

Π_B under a change of metric

The operator Γ_g does not change under the change of metric. However,

$$\Gamma_g^* = C^{-1} \Gamma_{\tilde{g}}^* C$$

where C is the bounded, invertible, multiplication operator on $L^2(\mathcal{M}) \oplus L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M})$.

Π_B under a change of metric

The operator Γ_g does not change under the change of metric. However,

$$\Gamma_g^* = C^{-1} \Gamma_{\tilde{g}}^* C$$

where C is the bounded, invertible, multiplication operator on $L^2(\mathcal{M}) \oplus L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M})$.

Thus,

$$\Pi_{B,g} = \Gamma_g + B_1 \Gamma_g^* B_2 = \Gamma_{\tilde{g}} + B_1 C^{-1} \Gamma_{\tilde{g}}^* C B_2.$$

Π_B under a change of metric

The operator Γ_g does not change under the change of metric. However,

$$\Gamma_g^* = C^{-1} \Gamma_{\tilde{g}}^* C$$

where C is the bounded, invertible, multiplication operator on $L^2(\mathcal{M}) \oplus L^2(\mathcal{M}) \oplus L^2(T^*\mathcal{M})$.

Thus,

$$\Pi_{B,g} = \Gamma_g + B_1 \Gamma_g^* B_2 = \Gamma_{\tilde{g}} + B_1 C^{-1} \Gamma_{\tilde{g}}^* C B_2.$$

This allows us to reduce the study of $\Pi_{B,g}$ for a C^0 metric g to the study of $\Pi_{\tilde{B},\tilde{g}} = \Gamma_{\tilde{g}} + \tilde{B}_1 \Gamma_{\tilde{g}}^* \tilde{B}_2$ where $\tilde{B}_1 = B_1 C^{-1}$ and $\tilde{B}_2 = C B_2$, but now with a smooth metric \tilde{g} .

Connection to geometric flows

Given a C^0 metric g on a smooth *compact* manifold, we are able to always find C^∞ metric \tilde{g} .

Connection to geometric flows

Given a C^0 metric g on a smooth *compact* manifold, we are able to always find C^∞ metric \tilde{g} .

The metric \tilde{g} has $\text{inj}(\mathcal{M}, \tilde{g}) > \kappa$ and $|\text{Ric}(\tilde{g})|_{\tilde{g}} \leq \eta$ so we obtain a corresponding Kato square root estimate in this setting.

Connection to geometric flows

Given a C^0 metric g on a smooth *compact* manifold, we are able to always find C^∞ metric \tilde{g} .

The metric \tilde{g} has $\text{inj}(\mathcal{M}, \tilde{g}) > \kappa$ and $|\text{Ric}(\tilde{g})|_{\tilde{g}} \leq \eta$ so we obtain a corresponding Kato square root estimate in this setting.

The non-compact situation poses issues.

Connection to geometric flows

Given a C^0 metric g on a smooth *compact* manifold, we are able to always find C^∞ metric \tilde{g} .

The metric \tilde{g} has $\text{inj}(\mathcal{M}, \tilde{g}) > \kappa$ and $|\text{Ric}(\tilde{g})|_{\tilde{g}} \leq \eta$ so we obtain a corresponding Kato square root estimate in this setting.

The non-compact situation poses issues.

Smooth the metric via mean curvature flow for, say, a C^2 imbedding?

Connection to geometric flows

Given a C^0 metric g on a smooth *compact* manifold, we are able to always find C^∞ metric \tilde{g} .

The metric \tilde{g} has $\text{inj}(\mathcal{M}, \tilde{g}) > \kappa$ and $|\text{Ric}(\tilde{g})|_{\tilde{g}} \leq \eta$ so we obtain a corresponding Kato square root estimate in this setting.

The non-compact situation poses issues.

Smooth the metric via mean curvature flow for, say, a C^2 imbedding?

Smooth the metric via Ricci flow in the general case? Regularity of the initial metric?

Application to PDE

In the case we are able to find a suitable C^∞ metric near the C^0 one, then we have Lipschitz estimates.

Application to PDE

In the case we are able to find a suitable C^∞ metric near the C^0 one, then we have Lipschitz estimates.

Possible application to hyperbolic PDE?

Application to PDE

In the case we are able to find a suitable C^∞ metric near the C^0 one, then we have Lipschitz estimates.

Possible application to hyperbolic PDE?

“Stability” of geometries with Ricci bounds and injectivity radius bounds?

References I

- [AHLMcT] Pascal Auscher, Steve Hofmann, Michael Lacey, Alan McIntosh, and Ph. Tchamitchian, *The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n* , Ann. of Math. (2) **156** (2002), no. 2, 633–654.
- [AKMc-2] Andreas Axelsson, Stephen Keith, and Alan McIntosh, *The Kato square root problem for mixed boundary value problems*, J. London Math. Soc. (2) **74** (2006), no. 1, 113–130.
- [AKMc] ———, *Quadratic estimates and functional calculi of perturbed Dirac operators*, Invent. Math. **163** (2006), no. 3, 455–497.
- [Aus] Pascal Auscher, *Lectures on the Kato square root problem*, Surveys in analysis and operator theory (Canberra, 2001), Proc. Centre Math. Appl. Austral. Nat. Univ., vol. 40, Austral. Nat. Univ., Canberra, 2002, pp. 1–18.

References II

- [Christ] Michael Christ, *A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral*, Colloq. Math. **60/61** (1990), no. 2, 601–628.
- [Mc72] Alan McIntosh, *On the comparability of $A^{1/2}$ and $A^{*1/2}$* , Proc. Amer. Math. Soc. **32** (1972), 430–434.
- [Mc82] ———, *On representing closed accretive sesquilinear forms as $(A^{1/2}u, A^{*1/2}v)$* , Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. III (Paris, 1980/1981), Res. Notes in Math., vol. 70, Pitman, Boston, Mass., 1982, pp. 252–267.
- [Morris] Andrew J. Morris, *The Kato square root problem on submanifolds*, J. Lond. Math. Soc. (2) **86** (2012), no. 3, 879–910.