CHAPTER 06

Aldehydes and Ketones I

Content and rendering review packet

Original app content and actual Chromium screenshots. No textbook passages or textbook figures are embedded. Source labels and curriculum headings are not evidence of factual correctness.

Source snapshot SHA-256: 9753d9af98eeba71e4832842ba3fe2740217e0310e4959123b817776406426e2

Review all claims, molecule graphs, lessons, exercises, hints and displayed states. Keys are author claims, not ground truth. Screenshots show static states with reduced motion: they do not prove animation timing or persistence behavior.

For every finding: item ID/page; severity (critical/major/minor); exact claim; derivation or reference evidence; proposed correction; confidence. List unreviewed/uncertain items. Model agreement is not scientific verification.

Coverage checklist

ch06:exercise:addition-site | addition-site:stage-1, addition-site:stage-2, addition-site:stage-3

ch06:exercise:hemiacetal-identify | hemiacetal-identify:stage-1, hemiacetal-identify:stage-2, hemiacetal-identify:stage-3

ch06:exercise:cyanohydrin-count | cyanohydrin-count:stage-1, cyanohydrin-count:stage-2, cyanohydrin-count:stage-3

ch06:exercise:hydride-product | hydride-product:stage-1, hydride-product:stage-2, hydride-product:stage-3

ch06:lesson:carbonyl | carbonyl:rule, carbonyl:cue, carbonyl:trap, carbonyl:app, carbonyl:step-1, carbonyl:step-2

ch06:lesson:oxygen-addition | oxygen-addition:rule, oxygen-addition:cue, oxygen-addition:trap, oxygen-addition:app, oxygen-addition:step-1, oxygen-addition:step-2, oxygen-addition:step-3

ch06:lesson:nitrogen-addition | nitrogen-addition:rule, nitrogen-addition:cue, nitrogen-addition:trap, nitrogen-addition:app, nitrogen-addition:step-1, nitrogen-addition:step-2, nitrogen-addition:step-3

ch06:lesson:carbonyl-redox | carbonyl-redox:rule, carbonyl-redox:cue, carbonyl-redox:trap, carbonyl-redox:app, carbonyl-redox:step-1, carbonyl-redox:step-2, carbonyl-redox:step-3

ch06:molecule:acetaldehyde | acetaldehyde

ch06:molecule:hemiacetal | hemiacetal

ch06:molecule:acetal | acetal

ch06:molecule:geminal | geminal

ch06:molecule:imine | imine

ch06:molecule:enamine | enamine

ch06:molecule:cyanohydrin | cyanohydrin

ch06:molecule:aceticacid | aceticacid

ch06:molecule:ethanol | ethanol

ch06:molecule:acetone | acetone

ch06:curriculum | scope-1, scope-2, scope-3

Curriculum scope

scope-1: 6.1 Description and Properties

scope-2: 6.2 Nucleophilic Addition Reactions

scope-3: 6.3 Oxidation–Reduction Reactions

Objective: Track carbonyl addition electron flow

Objective: Identify hydrates, hemiacetals, acetals, imines, and enamines

Objective: Predict carbonyl redox outcomes

Supplied Kaplan curriculum: Chapter 6; section scope checked against local body and summary | organic_chemistry_kaplan/OEBPS/chapter06b.xhtml

No source passage is included. Request relevant authoritative excerpts when a claim cannot be checked by derivation.

carbonyl | Carbonyl polarity and geometry

Planar electrophile → tetrahedral addition product

carbonyl:rule | Core Rules

The carbonyl is polarized Cδ+–Oδ−. A nucleophile attacks carbon while the pi pair shifts to oxygen; protonation can then give an alcohol. Aldehydes are generally less crowded and less electron-rich than comparable ketones, so they often add nucleophiles more readily.

carbonyl:cue | Memory Cues

Attack C; move pi electrons to O; then handle proton transfers.

carbonyl:trap | Common Mistakes / MCAT Traps

Neutral aldehydes and ketones can accept hydrogen bonds but lack an O–H donor. Do not draw five bonds to carbon when a nucleophile adds. A planar carbonyl can be attacked from either face.

carbonyl:app | MCAT Applications

A chiral environment can favor one face; without such bias, addition that creates a single new stereocenter often yields a racemic mixture.


carbonyl:step-1 | 1. Find the electrophile

The C=O carbon is trigonal planar and electron poor. Oxygen is electron rich.

Displayed molecule ID: acetaldehyde

{"highlights": {"prin": [1], "sub": [2]}, "revealName": false}


Screenshot: carbonyl-step-1



Embedded challenge: Select the electrophilic carbonyl carbon.

{"prompt": "Select the electrophilic carbonyl carbon.", "answer": [1], "explanation": "B is the carbonyl carbon."}

Screenshot: carbonyl-step-1-challenge




carbonyl:step-2 | 2. Move the pi pair

The purple curved arrow shows the carbonyl pi pair moving toward oxygen during nucleophilic attack. This arrow is one part of an addition mechanism; the incoming nucleophile is described in the lesson rather than drawn here.

Displayed molecule ID: acetaldehyde

{"highlights": {"prin": [1, 2], "flow": [{"fromBond": 1, "toAtom": 2}]}, "revealName": false}

Screenshot: carbonyl-step-2



oxygen-addition | Hydrates, hemiacetals, and acetals

Distinguish OH from OR on the same carbon

oxygen-addition:rule | Core Rules

Hydration gives a geminal diol. Alcohol addition gives a hemiacetal (one OH and one OR at the former carbonyl carbon). Further acid-catalyzed reaction gives an acetal (two OR groups). These transformations are reversible and respond to water/alcohol activities.

oxygen-addition:cue | Memory Cues

Two OH = hydrate; OH + OR = hemiacetal; two OR = acetal. Ketone-derived examples are also called hemiketals/ketals.

oxygen-addition:trap | Common Mistakes / MCAT Traps

Acetal formation requires acid catalysis; base alone does not turn OH− into a good leaving group. Cyclic sugar hemiacetals can be stable because intramolecular ring formation is favorable.

oxygen-addition:app | MCAT Applications

Monosaccharides cyclize through hemiacetal or hemiketal formation. The new anomeric center can have two configurations.


oxygen-addition:step-1 | 1. Identify a hemiacetal

This carbon bears both OH and OCH3, along with methyl and H. It is a hemiacetal from ethanal and methanol.

Displayed molecule ID: hemiacetal

{"highlights": {"prin": [1, 2, 3]}, "revealName": true}

Screenshot: oxygen-addition-step-1




oxygen-addition:step-2 | 2. Compare an acetal

This ketone-derived acetal has two OCH3 groups on the central carbon and no OH there. Hydrolysis with aqueous acid returns a carbonyl.

Displayed molecule ID: acetal

{"highlights": {"prin": [1, 3, 4]}, "revealName": true}

Screenshot: oxygen-addition-step-2




oxygen-addition:step-3 | 3. Compare a hydrate

Here both oxygen substituents are OH: a geminal diol. The equilibrium does not equally favor hydrates for every carbonyl.

Displayed molecule ID: geminal

{"highlights": {"prin": [1, 3, 4]}, "revealName": true}

Screenshot: oxygen-addition-step-3



nitrogen-addition | Imines, enamines, and cyanohydrins

Reagent identity changes the product

nitrogen-addition:rule | Core Rules

A primary amine condenses with an aldehyde or ketone to form an imine after addition, proton transfers, and water loss. A secondary amine can form an enamine when an alpha H is available. Related nitrogen reagents form oximes (hydroxylamine), hydrazones (hydrazine derivatives), and semicarbazones (semicarbazide). Cyanide addition followed by protonation forms a cyanohydrin and creates a C–C bond.

nitrogen-addition:cue | Memory Cues

Primary amine → C=N; secondary amine + alpha H → C=C–N. Cyanide adds its carbon.

nitrogen-addition:trap | Common Mistakes / MCAT Traps

Too much acid protonates the amine and suppresses its nucleophilicity. Enamine formation needs an alpha hydrogen. Do not attach cyanide through nitrogen in a standard cyanohydrin product.

nitrogen-addition:app | MCAT Applications

Schiff-base (imine) intermediates help enzymes reorganize electron density; nitrogen chemistry is a bridge to amino-acid metabolism.


nitrogen-addition:step-1 | 1. Imine bond

The C=N bond replaces the starting C=O after condensation with a primary amine. This connectivity drawing does not assign E/Z about C=N.

Displayed molecule ID: imine

{"highlights": {"prin": [1, 2]}, "revealName": true}

Screenshot: nitrogen-addition-step-1




nitrogen-addition:step-2 | 2. Enamine bond

An enamine has N attached to a C=C system. The nitrogen lone pair is conjugated with the double bond, giving nucleophilic carbon character.

Displayed molecule ID: enamine

{"highlights": {"path": [0, 1, 2], "prin": [2]}, "revealName": true}

Screenshot: nitrogen-addition-step-2




nitrogen-addition:step-3 | 3. Cyanohydrin carbon count

Acetone cyanohydrin contains a new C–C bond between the former carbonyl carbon and the nitrile carbon. The nitrile N is triple-bonded to that new carbon.

Displayed molecule ID: cyanohydrin

{"highlights": {"prin": [1, 3], "sub": [4, 5]}, "revealName": true}

Screenshot: nitrogen-addition-step-3



carbonyl-redox | Aldehyde oxidation and hydride reduction

Separate addition from net redox

carbonyl-redox:rule | Core Rules

Aldehydes can be oxidized to carboxylic acids; ordinary ketones resist analogous oxidation without C–C cleavage. NaBH4 and LiAlH4 reduce aldehydes to primary alcohols and ketones to secondary alcohols. Hydride adds first, then oxygen is protonated in workup or by the medium.

carbonyl-redox:cue | Memory Cues

Aldehyde reduction → 1° alcohol; ketone reduction → 2° alcohol.

carbonyl-redox:trap | Common Mistakes / MCAT Traps

Do not predict an alkane from ordinary hydride reduction of C=O. The oxygen remains in the alcohol. A positive aldehyde oxidation test is contextual evidence, not a complete structural identification.

carbonyl-redox:app | MCAT Applications

Biological NADH-dependent carbonyl reductions transfer a hydride equivalent; the enzyme can control stereochemistry.


carbonyl-redox:step-1 | 1. Aldehyde substrate

Ethanal has a hydrogen on its carbonyl carbon, allowing ordinary oxidation to a carboxylic acid.

Displayed molecule ID: acetaldehyde

{"highlights": {"prin": [1, 2]}, "revealName": true}

Screenshot: carbonyl-redox-step-1




carbonyl-redox:step-2 | 2. Oxidized product

Ethanoic acid retains the two-carbon skeleton and adds a carbon–oxygen bond overall.

Displayed molecule ID: aceticacid

{"highlights": {"prin": [1, 2, 3]}, "revealName": true}

Screenshot: carbonyl-redox-step-2




carbonyl-redox:step-3 | 3. Reduced product

The reduction product ethanol retains the same oxygen but has a single C–O bond and more C–H bonding.

Displayed molecule ID: ethanol


{"highlights": {"prin": [1, 2]}, "revealName": true}

Screenshot: carbonyl-redox-step-3



addition-site | Follow nucleophilic addition


Screenshot: addition-site-stage-1-molecule



addition-site:stage-1 | Select the atom attacked by a carbon nucleophile.

Answer type: atoms

Screenshot: addition-site-stage-1-question




{"authorAnswer": [1], "tolerance": null, "highlightsAfterCorrectAnswer": {"prin": [1]}}

Hint 1: Locate the positively polarized end of C=O.

Hint 2: Carbon is less electronegative than oxygen.

Explanation: The nucleophile attacks the carbonyl carbon.

Screenshot: addition-site-stage-1-feedback



addition-site:stage-2 | As the new sigma bond forms, the C=O pi pair moves to...


Answer type: choice

Option 0 (A): Oxygen


Option 1 (B): The incoming nucleophile

Option 2 (C): A methyl hydrogen

Screenshot: addition-site-stage-2-molecule



Screenshot: addition-site-stage-2-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {"flow": [{"fromBond": 1, "toAtom": 2}]}}

Hint 1: Count the carbon bonds during attack.

Hint 2: O can carry the resulting negative charge.

Explanation: Moving the pi pair to O prevents carbon from exceeding an octet.

Screenshot: addition-site-stage-2-feedback



addition-site:stage-3 | The immediate former-carbonyl geometry becomes approximately...


Answer type: choice

Option 0 (A): Tetrahedral


Option 1 (B): Linear

Option 2 (C): Unchanged trigonal planar

Screenshot: addition-site-stage-3-molecule



Screenshot: addition-site-stage-3-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Count electron domains after addition.

Hint 2: Four directions correspond to sp^3-like geometry.


Explanation: Addition replaces the pi bond and gives four sigma directions around carbon.

Screenshot: addition-site-stage-3-feedback




Completion summary: Nucleophilic addition changes planar C=O into a tetrahedral center.

Screenshot: addition-site-complete



hemiacetal-identify | Read oxygen substituents


Screenshot: hemiacetal-identify-stage-1-molecule



hemiacetal-identify:stage-1 | Select the carbon bearing two oxygen substituents.

Answer type: atoms

Screenshot: hemiacetal-identify-stage-1-question




{"authorAnswer": [1], "tolerance": null, "highlightsAfterCorrectAnswer": {"prin": [1, 2, 3]}}

Hint 1: Trace both C–O bonds to their shared carbon.

Hint 2: The O atoms are C and D.

Explanation: B is the former carbonyl carbon.

Screenshot: hemiacetal-identify-stage-1-feedback



hemiacetal-identify:stage-2 | One OH and one OR on that carbon identify a...


Answer type: choice

Option 0 (A): Hemiacetal


Option 1 (B): Acetal

Option 2 (C): Geminal diol

Screenshot: hemiacetal-identify-stage-2-molecule



Screenshot: hemiacetal-identify-stage-2-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Inspect whether each oxygen also has a carbon substituent.

Hint 2: One oxygen is OH; the other links to methyl.

Explanation: A hemiacetal has both OH and OR.

Screenshot: hemiacetal-identify-stage-2-feedback



hemiacetal-identify:stage-3 | Converting it to an acetal with alcohol is favored by...


Answer type: choice

Option 0 (A): Acid catalysis and removal of water


Option 1 (B): Adding abundant water alone

Option 2 (C): Strong base alone

Screenshot: hemiacetal-identify-stage-3-molecule



Screenshot: hemiacetal-identify-stage-3-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Water is a product of acetal formation.

Hint 2: Apply Le Châtelier's principle.


Explanation: Acid enables water departure; removing water shifts the equilibrium toward acetal.

Screenshot: hemiacetal-identify-stage-3-feedback




Completion summary: Hemiacetal: OH + OR on the same carbon.

Screenshot: hemiacetal-identify-complete



cyanohydrin-count | Track the new carbon bond


Screenshot: cyanohydrin-count-stage-1-molecule



cyanohydrin-count:stage-1 | Select the nitrile carbon added by cyanide.

Answer type: atoms

Screenshot: cyanohydrin-count-stage-1-question




{"authorAnswer": [4], "tolerance": null, "highlightsAfterCorrectAnswer": {"prin": [4, 5]}}

Hint 1: Cyanide adds through carbon.

Hint 2: Find the carbon endpoint of C≡N.

Explanation: E is the carbon triple-bonded to N and bonded to the former carbonyl carbon.


Screenshot: cyanohydrin-count-stage-1-feedback




cyanohydrin-count:stage-2 | How many carbons are in the cyanohydrin?

Answer type: number | unit: carbons

Screenshot: cyanohydrin-count-stage-2-molecule



Screenshot: cyanohydrin-count-stage-2-question




{"authorAnswer": 4, "tolerance": 0, "highlightsAfterCorrectAnswer": {}}

Hint 1: Count carbon atoms, not total atoms.

Hint 2: Include the nitrile carbon.

Explanation: Acetone supplies three carbons and cyanide supplies one.

Screenshot: cyanohydrin-count-stage-2-feedback



cyanohydrin-count:stage-3 | A secondary amine plus a ketone with alpha H typically forms...


Answer type: choice

Option 0 (A): An enamine


Option 1 (B): A primary amide directly

Option 2 (C): A cyanohydrin

Screenshot: cyanohydrin-count-stage-3-molecule



Screenshot: cyanohydrin-count-stage-3-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Distinguish primary and secondary amine reagents.

Hint 2: The product contains C=C–N.


Explanation: A secondary amine gives an enamine after condensation and alpha deprotonation.

Screenshot: cyanohydrin-count-stage-3-feedback




Completion summary: Cyanohydrin formation adds one carbon; secondary amines can form enamines.

Screenshot: cyanohydrin-count-complete



hydride-product | Predict carbonyl reduction


Screenshot: hydride-product-stage-1-molecule



hydride-product:stage-1 | Select the carbonyl carbon receiving hydride.

Answer type: atoms

Screenshot: hydride-product-stage-1-question




{"authorAnswer": [1], "tolerance": null, "highlightsAfterCorrectAnswer": {"prin": [1, 3]}}

Hint 1: Hydride is the nucleophile.

Hint 2: Attack the electron-poor carbon.

Explanation: B receives hydride.

Screenshot: hydride-product-stage-1-feedback



hydride-product:stage-2 | After protonation, the product is a...


Answer type: choice

Option 0 (A): Secondary alcohol


Option 1 (B): Primary alcohol

Option 2 (C): Tertiary alcohol

Screenshot: hydride-product-stage-2-molecule



Screenshot: hydride-product-stage-2-question




{"authorAnswer": 0, "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Count carbon neighbors of the new C–OH carbon.

Hint 2: No C–C bonds are added or removed.

Explanation: The alcohol-bearing carbon retains its two methyl substituents.


Screenshot: hydride-product-stage-2-feedback




hydride-product:stage-3 | Name the product.

Answer type: text

Screenshot: hydride-product-stage-3-molecule



Screenshot: hydride-product-stage-3-question




{"authorAnswer": ["propan-2-ol", "2-propanol", "isopropanol"], "tolerance": null, "highlightsAfterCorrectAnswer": {}}

Hint 1: Retain the three-carbon chain.

Hint 2: Place OH at the central carbon.


Explanation: Acetone reduces to propan-2-ol.

Screenshot: hydride-product-stage-3-feedback



Completion summary: Propanone → propan-2-ol.

Screenshot: hydride-product-complete



Molecular repository: author claims to verify

acetaldehyde

Author claims: ethanal | formula C2H4O

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=50.00, y=115.00; charge=0; explicit H override=none

1 (B): C; x=108.00, y=85.00; charge=0; explicit H override=none

2 (C): O; x=108.00, y=25.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 2]]

hemiacetal

Author claims: 1-methoxyethan-1-ol | formula C3H8O2

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=60.00, y=120.00; charge=0; explicit H override=none

1 (B): C; x=120.00, y=85.00; charge=0; explicit H override=none

2 (C): O; x=120.00, y=25.00; charge=0; explicit H override=none

3 (D): O; x=180.00, y=120.00; charge=0; explicit H override=none

4 (E): C; x=240.00, y=85.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [1, 3, 1], [3, 4, 1]]

acetal

Author claims: 2,2-dimethoxypropane | formula C5H12O2

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=75.00, y=130.00; charge=0; explicit H override=none

1 (B): C; x=135.00, y=100.00; charge=0; explicit H override=none

2 (C): C; x=195.00, y=130.00; charge=0; explicit H override=none

3 (D): O; x=105.00, y=45.00; charge=0; explicit H override=none

4 (E): O; x=175.00, y=45.00; charge=0; explicit H override=none

5 (F): C; x=50.00, y=15.00; charge=0; explicit H override=none

6 (G): C; x=230.00, y=15.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [1, 3, 1], [1, 4, 1], [3, 5, 1], [4, 6, 1]]

geminal

Author claims: propane-2,2-diol | formula C3H8O2

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=48.00, y=115.00; charge=0; explicit H override=none

1 (B): C; x=102.00, y=85.00; charge=0; explicit H override=none

2 (C): C; x=156.00, y=115.00; charge=0; explicit H override=none

3 (D): O; x=76.00, y=32.00; charge=0; explicit H override=none

4 (E): O; x=130.00, y=32.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [1, 3, 1], [1, 4, 1]]

imine

Author claims: N-methylethanimine (E/Z unspecified) | formula C3H7N

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=55.00, y=120.00; charge=0; explicit H override=none

1 (B): C; x=115.00, y=85.00; charge=0; explicit H override=none

2 (C): N; x=180.00, y=85.00; charge=0; explicit H override=none

3 (D): C; x=240.00, y=120.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 2], [2, 3, 1]]

enamine

Author claims: N,N-dimethylethenamine | formula C4H9N

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=55.00, y=120.00; charge=0; explicit H override=none

1 (B): C; x=115.00, y=85.00; charge=0; explicit H override=none

2 (C): N; x=180.00, y=120.00; charge=0; explicit H override=none

3 (D): C; x=240.00, y=85.00; charge=0; explicit H override=none

4 (E): C; x=180.00, y=180.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 2], [1, 2, 1], [2, 3, 1], [2, 4, 1]]

cyanohydrin

Author claims: 2-hydroxy-2-methylpropanenitrile (acetone cyanohydrin) | formula C4H7NO

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=60.00, y=120.00; charge=0; explicit H override=none

1 (B): C; x=120.00, y=85.00; charge=0; explicit H override=none

2 (C): C; x=180.00, y=120.00; charge=0; explicit H override=none

3 (D): O; x=120.00, y=25.00; charge=0; explicit H override=none

4 (E): C; x=120.00, y=150.00; charge=0; explicit H override=none

5 (F): N; x=120.00, y=205.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [1, 3, 1], [1, 4, 1], [4, 5, 3]]

aceticacid

Author claims: ethanoic acid | formula C2H4O2

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=50.00, y=115.00; charge=0; explicit H override=none

1 (B): C; x=108.00, y=85.00; charge=0; explicit H override=none

2 (C): O; x=108.00, y=25.00; charge=0; explicit H override=none

3 (D): O; x=166.00, y=115.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 2], [1, 3, 1]]

ethanol

Author claims: ethanol | formula C2H6O

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=50.00, y=115.00; charge=0; explicit H override=none

1 (B): C; x=108.00, y=85.00; charge=0; explicit H override=none

2 (C): O; x=166.00, y=115.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1]]

acetone

Author claims: propanone | formula C3H6O

Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=50.00, y=115.00; charge=0; explicit H override=none

1 (B): C; x=108.00, y=85.00; charge=0; explicit H override=none

2 (C): C; x=166.00, y=115.00; charge=0; explicit H override=none


3 (D): O; x=108.00, y=25.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [1, 3, 2]]


Layout samples

These overview images are scaled for page fit. Detailed structure and stage images appear earlier. Samples cover the first concept on desktop and mobile in both themes, not every responsive state.


Screenshot: layout-desktop-light



Screenshot: layout-mobile-light



Screenshot: layout-mobile-dark



Execution evidence

55 screenshots and 13 recorded interactions. Incorrect answers rejected and encoded keys accepted for captured guided stages. This checks implementation against its data; it does not validate chemistry independently. Static screenshots do not verify animation timing, arbitrary clicks, or screen-reader experience.