CHAPTER 10

Nitrogen- and Phosphorus-Containing Compounds

Independent exercise solving

Original app content and actual Chromium screenshots. No textbook passages or textbook figures are embedded. Source labels and curriculum headings are not evidence of factual correctness.


Source snapshot SHA-256: 4d975bae2603f4ebd33ecb97980538e370fde47bf5493578198d17cd97c2e8af

BLIND PASS: answer keys, author molecule names/formulas, hints, explanations and answer overlays are withheld. Solve every numbered stage independently. Report the stage ID, answer, derivation, and uncertainty. Question wording and displayed diagram captions are retained because they are visible task data.

Do not open the full review packet in this chat before saving your independent attempt. The full packet contains keys and lessons that would contaminate a blind attempt.

glycine-charge | Independent exercise

Screenshot: glycine-charge-stage-1-molecule



Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): N; x=45.00, y=115.00; charge=1; explicit H override=none

1 (B): C; x=105.00, y=85.00; charge=0; explicit H override=none

2 (C): C; x=165.00, y=115.00; charge=0; explicit H override=none

3 (D): O; x=165.00, y=175.00; charge=0; explicit H override=none

4 (E): O; x=225.00, y=85.00; charge=-1; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [2, 3, 2], [2, 4, 1]]

glycine-charge:stage-1 | What is the net formal charge of the displayed zwitterion?

Answer type: choice

Option 0 (A): 0

Option 1 (B): +1

Option 2 (C): −1

glycine-charge:stage-2 | Why is glycine achiral?

Answer type: choice

Option 0 (A): Its alpha carbon has two H substituents

Option 1 (B): All amino acids are achiral


Option 2 (C): Its nitrogen is neutral

glycine-charge:stage-3 | Using pKa values 2.3 and 9.6 for this neutral-side-chain model, calculate pI.

Answer type: number | unit: pH units

peptide-link | Independent exercise

Screenshot: peptide-link-stage-1-molecule



Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): N; x=35.00, y=110.00; charge=1; explicit H override=none

1 (B): C; x=90.00, y=80.00; charge=0; explicit H override=none

2 (C): C; x=145.00, y=110.00; charge=0; explicit H override=none

3 (D): O; x=145.00, y=170.00; charge=0; explicit H override=none

4 (E): N; x=200.00, y=80.00; charge=0; explicit H override=none

5 (F): C; x=255.00, y=110.00; charge=0; explicit H override=none

6 (G): C; x=310.00, y=80.00; charge=0; explicit H override=none

7 (H): O; x=310.00, y=20.00; charge=0; explicit H override=none

8 (I): O; x=365.00, y=110.00; charge=-1; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [2, 3, 2], [2, 4, 1], [4, 5, 1], [5, 6, 1], [6, 7, 2], [6, 8, 1]]

peptide-link:stage-1 | Select the two endpoints of the internal peptide C–N bond.

Answer type: atoms

peptide-link:stage-2 | Why is rotation around this bond restricted?

Answer type: choice

Option 0 (A): Amide resonance gives partial double-bond character

Option 1 (B): Every single bond is completely rigid

Option 2 (C): The bond is an ionic salt bridge

peptide-link:stage-3 | Peptide sequences are read from…


Answer type: choice

Option 0 (A): N-terminus to C-terminus

Option 1 (B): C-terminus to N-terminus by default

strecker-map | Independent exercise

Screenshot: strecker-map-stage-1-molecule



Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): C; x=50.00, y=120.00; charge=0; explicit H override=none

1 (B): C; x=110.00, y=85.00; charge=0; explicit H override=none

2 (C): C; x=170.00, y=120.00; charge=0; explicit H override=none

3 (D): N; x=230.00, y=155.00; charge=0; explicit H override=none

4 (E): N; x=110.00, y=25.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 1], [1, 2, 1], [2, 3, 3], [1, 4, 1]]

strecker-map:stage-1 | Select the carbon that becomes the carboxyl carbon after hydrolysis.

Answer type: atoms

strecker-map:stage-2 | Which group is retained as the alanine side chain?

Answer type: choice

Option 0 (A): Methyl

Option 1 (B): Nitrile nitrogen

Option 2 (C): A new phenyl group

strecker-map:stage-3 | Under achiral conditions, the chiral product is generally…

Answer type: choice


Option 0 (A): Racemic

Option 1 (B): Only L without a chiral influence

Option 2 (C): Only D without a chiral influence

phosphate-links | Independent exercise

Screenshot: phosphate-links-stage-1-molecule



Graph uses zero-based indices; A=0, B=1, etc. Coordinates are drawing positions, not 3D coordinates.

0 (A): P; x=160.00, y=100.00; charge=0; explicit H override=none

1 (B): O; x=160.00, y=30.00; charge=0; explicit H override=none

2 (C): O; x=90.00, y=100.00; charge=0; explicit H override=none

3 (D): O; x=230.00, y=100.00; charge=0; explicit H override=none

4 (E): O; x=160.00, y=170.00; charge=-1; explicit H override=none

5 (F): C; x=35.00, y=140.00; charge=0; explicit H override=none

6 (G): C; x=285.00, y=140.00; charge=0; explicit H override=none

Bonds [atom A, atom B, order]: [[0, 1, 2], [0, 2, 1], [0, 3, 1], [0, 4, 1], [2, 5, 1], [3, 6, 1]]

phosphate-links:stage-1 | Select the phosphorus atom.

Answer type: atoms

phosphate-links:stage-2 | This structure models a…

Answer type: choice

Option 0 (A): Phosphodiester

Option 1 (B): Phosphoanhydride

Option 2 (C): Carboxylic anhydride

phosphate-links:stage-3 | Why can ATP hydrolysis be favorable overall?

Answer type: choice

Option 0 (A): Products have lower free energy under the stated conditions

Option 1 (B): Breaking a bond itself releases energy

Option 2 (C): All phosphate bonds break spontaneously at the same rate