
PCA of Combined Datasets
60
40
20
0
−20
PC2 20.21% Variance
−120
−80
−40
0
PC1 29.33% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
60
40
20
0
−20
PC3 11.34% Variance
−20
0
20
40
60
PC2 20.21% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
20
0
−20
−40
PC4 6.3% Variance
−20
0
20
40
60
PC3 11.34% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
25
0
−25
−50
PC5 5.57% Variance
−40
−20
0
20
PC4 6.3% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
40
20
0
−20
PC6 5.27% Variance
−50
−25
0
25
PC5 5.57% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
20
0
−20
PC7 3.61% Variance
−20
0
20
40
PC6 5.27% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka


PCA of Combined Datasets
40
20
0
−20
PC8 2.74% Variance
−20
0
20
PC7 3.61% Variance
genotype
heterozygous_Vav1−iCre_and_homozygous_Setbp1−floxed_allele
homozygous_Setbp1−floxed_allele
PATH2
PATH3
SETBP1−KO
VUS2
WT
shape
Cardo
Shaw
Tanaka