

# Unified Brane-Cosmology: A Testable Route to Unification

Ricardo Maldonado | sales@rank.vegas | Draft v1 (20250811\_192837)

**Abstract:** We propose a higher-dimensional (bulk + brane) framework in which the 4D Friedmann equation on the brane acquires a high-energy  $\rho^2$  correction and a dark-radiation term  $C/a^4$ . A single parameter (brane tension  $\lambda$ ) links a measurable GW spectral break  $f_{\text{br}} \sim \lambda^{1/4}$  to  $\Delta N_{\text{eff}}$ . We outline falsifiable predictions and a pipeline for near-term tests with PTA  $\rightarrow$  LISA and CMB/BBN.

## 1. Framework

Consider a 5D warped bulk with a 4D brane. Standard Model fields are confined to the brane; gravity propagates in the bulk. Junction conditions relate the extrinsic curvature to the brane stress tensor and tension  $\lambda$ . The effective 4D field equation on the brane contains: (i) standard Einstein tensor with  $\Lambda_4$ , (ii) quadratic stress term  $\Pi_{\mu\nu}$ , (iii) projected bulk Weyl term  $E_{\mu\nu}$  (dark radiation).

## 2. Effective 4D Equation on the Brane

$G_{\mu\nu} + \Lambda_4 g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu} + (\kappa_5^4/c^4) \Pi_{\mu\nu} - E_{\mu\nu}$ . The quadratic term  $\Pi_{\mu\nu}$  captures high-energy corrections at  $\rho \gg \lambda$ . The Weyl term  $E_{\mu\nu}$  encodes bulk memory; for FRW it yields a radiation-like contribution  $\rho_{\text{dr}} \sim C/a^4$ .

## 3. Cosmology Reduction (flat FRW)

For a perfect fluid on the brane with pressure  $p$  and density  $\rho$ , we obtain  $H^2 = (8\pi G/3) \rho (1 + \rho/(2\lambda)) + \Lambda_4/3 + C/a^4 - k/a^2$ . At early times ( $\rho \gg \lambda$ ) the  $\rho^2$  term dominates, giving  $a(t) \sim t^{1/4}$  in radiation era, distinct from standard  $a \sim t^{1/2}$ .

## 4. Observables and Single-Parameter Link

The brane tension  $\lambda$  sets the energy scale where the  $\rho^2$  term turns off and the spectrum of primordial tensor modes transitions, producing a broken-power-law stochastic GW background. The same physics contributes an effective dark radiation density parameterized as  $\Delta N_{\text{eff}}$ , hence a correlation  $f_{\text{br}}(\lambda) \leftrightarrow \Delta N_{\text{eff}}$ .

## 5. Minimal Likelihood

We forecast by fitting a broken power-law SGWB to PTA points and a prior on  $\Delta N_{\text{eff}}$  from BBN/CMB, with parameters  $\{A_{\text{low}}, A_{\text{high}}, f_{\text{br}}(\lambda), \text{slopes}\}$ . The mapping  $f_{\text{br}} \sim \lambda^{1/4}$  closes the model; a single  $\lambda$  must satisfy both datasets. We compute posteriors for  $E_{\text{br}}$  (proxy for  $\lambda$ ) and  $\Delta N_{\text{eff}}$ .

## 6. Comparison to Prior Work

Our setup follows the brane-cosmology literature (Randall-Sundrum, Binetruy-Deffayet-Langlois, Maartens) but emphasizes a single measurable link ( $f_{\text{br}} \leftrightarrow \lambda$ ) and a joint fit with  $\Delta N_{\text{eff}}$ . We discuss overlaps and differences in parameterization and phenomenology.

## 7. Limitations and Open Tasks

We do not yet present a concrete compactification reproducing  $SU(3) \times SU(2) \times U(1)$  and full spectra/Yukawas. Quantum completion and moduli stabilization are stated as future work. The present draft focuses on near-term falsifiable signals and a data-driven test.

## 8. Roadmap

Step A: replace synthetic CSVs with public PTA data; rerun likelihood; publish code+preprint. Step B: extend with LISA priors and structure-growth constraints; Step C: compactification map linking  $(k, M_5, \lambda)$  to low-energy observables, then submit to PRD/JCAP.

## **References (selected)**

Key brane cosmology references (to be inserted with full citations): Randall & Sundrum (1999); Binetruy, Deffayet & Langlois (2000); Shiromizu, Maeda & Sasaki (2000); Maartens (2004) and subsequent developments on brane-world cosmology and dark radiation constraints.

# References — Unified Brane-Cosmology (curated)

- [1] Randall, L., Sundrum, R. 'A Large Mass Hierarchy from a Small Extra Dimension.' Phys. Rev. Lett. 83, 3370 (1999). DOI:10.1103/PhysRevLett.83.3370
- [2] Binetruy, P., Deffayet, C., Langlois, D. 'Brane cosmological evolution in a bulk with cosmological constant.' Phys. Lett. B 477, 285 (2000).
- [3] Shiromizu, T., Maeda, K., Sasaki, M. 'The Einstein equations on the 3-brane world.' Phys. Rev. D 62, 024012 (2000). DOI:10.1103/PhysRevD.62.024012
- [4] Maartens, R. 'Brane-World Gravity.' Living Rev. Relativity 7, 7 (2004); update arXiv:1004.3962
- [5] NANOGrav Collaboration (Agazie et al.). 'The NANOGrav 15-year Data Set: Evidence for a Gravitational-Wave Background.' ApJL 951, L8 (2023); arXiv:2306.16213
- [6] EPTA+InPTA Collaboration (Antoniadis et al.). 'The second data release from the European Pulsar Timing Array (DR2): Search for an isotropic GWB.' A&A 678, A48 (2023).
- [7] PPTA Collaboration (Reardon et al.). 'Search for an isotropic gravitational-wave background with the Parkes PTA.' (2023) arXiv:2306.16215
- [8] Planck Collaboration (Aghanim et al.). 'Planck 2018 results. VI. Cosmological parameters.' A&A 641, A6 (2020). arXiv:1807.06209
- [9] PDG Review: 'Big Bang Nucleosynthesis.' Review of Particle Physics, 2022 update (Fields et al.); PDG chapter.
- [10] Cooke, R. J., Pettini, M., Steidel, C. C. 'One percent determination of the primordial deuterium abundance.' ApJ 855, 102 (2018).
- [11] Robson, T., Cornish, N. J., Liu, C. 'The construction and use of LISA sensitivity curves.' Class. Quantum Grav. 36, 105011 (2019). arXiv:1803.01944
- [12] Babak, S., Hewitson, M., Petiteau, A. 'LISA Sensitivity and SNR Calculations.' (2021) LISA-LCST-SGS-TN-001.