

Mathematics Notebook

Mark H. Olson

Updated: December 20, 2014



Contents

1	Algebraic Expressions	5
1.1	Expressions	5
1.2	Polynomial Expressions	5
1.2.1	Monomials of Like Terms	9
1.2.2	Surds	9
2	Univariate Polynomial Expressions	11
2.1	Classification of Univariate Polynomial Expressions	11
2.2	Degree -1 Univariate Polynomials	13
2.3	Degree 0 Univariate Polynomials	13
2.4	Degree 1 Univariate Polynomials	13
2.5	Degree 2 Univariate Polynomials	30
2.6	Degree 3 Univariate Polynomials	40
2.7	Degree n Univariate Polynomials	40
3	Multivariate Polynomial Expressions	41
4	Equations	49
4.1	Equality	49
4.2	Solving Linear Equations	49
4.3	Solving Quadratic Equations	54
4.3.1	Completing the Square	55

5	Functions	59
5.1	Evaluating Functions	59
5.2	Quadratic Functions	60
5.2.1	Completing the Square	60
5.3	Rational Functions	61
6	Vectors	65
6.1	Component & Geometric Vectors	65
6.2	Dot Product	65
7	Differentiation	67
7.1	Limit of the Difference Quotient	67
7.2	Derivative of a Monomial Functions	68
7.3	Derivative of Polynomial Functions	69
7.4	Derivative of Rational Functions	72
7.5	Derivative of Trigonometric Functions	72
7.6	Derivative of Exponential Functions	72
7.7	Derivative of Logarithmic Functions	72
7.8	Derivative of Composite Functions	73
8	Integration	77
8.1	Integration Rules	77
8.2	Volumes of Revolution	78
9	Answers	83
9.0.1	id:20141120-211843	83
9.0.2	id:20141121-195608	83
9.0.3	id:20141121-191224	83
9.0.4	id:20141123-091420	83
	Bibliography	85
	Websites	85
	Index	87

Expressions

Polynomial Expressions

Monomials of Like Terms

Surds

1. Algebraic Expressions

1.1 Expressions

Essential Questions 1.1

1. What is an algebraic expression?

1.2 Polynomial Expressions

Definition 1.2.1 — Operation of Addition (OOA).

$$\underbrace{a}_{\text{Augend}} + \underbrace{b}_{\text{Addend}} = \underbrace{\text{Sum}}_{(1.1)}$$

	Arithmetic	Polynomial	Algebraic
Constant	Yes	Yes	Yes
Factorial	Yes	Yes	Yes
Variable: parameter/coefficient	Yes	Yes	Yes
Variable: unknown/indeterminate	No	Yes	Yes
Power with \mathbb{Z}^+ exponent	No	Yes	Yes
Power with \mathbb{Z} exponent	No	No	Yes
n -th root	No	No	Yes
Power with \mathbb{Q} exponent	No	No	Yes

Table 1.1: Names of different types of expressions

Definition 1.2.2 — Common Denominator (CD).

$$\frac{a}{b} + \frac{c}{b} = \frac{a+c}{b} \quad (1.2a)$$

$$\frac{a+c}{b} = \frac{a}{b} + \frac{c}{b} \quad (1.2b)$$

Rule 1.2.3 — Fraction Operation of Addition (FOOA).

$$\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd} \quad (1.3a)$$

$$\frac{ad+bc}{bd} = \frac{a}{b} + \frac{c}{d} \quad (1.3b)$$

Definition 1.2.4 — Operation of Multiplication (OOM).

$$\underbrace{\begin{array}{c} a \\ \text{Multiplicand} \end{array}}_{\text{Product}} \cdot \underbrace{\begin{array}{c} b \\ \text{Multiplier} \end{array}}_{\text{Product}} \quad (1.4)$$

Definition 1.2.5 — Operation of Exponentiation (OOE).

$$\underbrace{\begin{array}{c} b \\ \text{base} \end{array}}_{\text{Power}}^m \quad (1.5)$$

Definition 1.2.6 — Juxtaposition to Center-Dot (JTC).

$$ab = a \cdot b \quad (1.6)$$

Definition 1.2.7 — Center-Dot to Justaposition (CTJ).

$$a \cdot b = ab \quad (1.7)$$

Definition 1.2.8 — Commutative Property of Multiplication (CPM).

$$\textcolor{red}{a} \cdot b = b \cdot \textcolor{red}{a} \quad (1.8)$$

Definition 1.2.9 — Multiplicative Inverse (MI).

$$a \cdot \frac{1}{\textcolor{red}{a}} = 1 \quad (1.9a)$$

$$a \cdot \textcolor{red}{a}^{-1} = 1 \quad (1.9b)$$

Definition 1.2.10 — Associative Property of Multiplication (APM).

$$a \cdot b \cdot c = (a \cdot b) \cdot c \quad (1.10a)$$

$$a \cdot b \cdot c = a \cdot (b \cdot c) \quad (1.10b)$$

Powers

Rule 1.2.11 — Power of a Quotient of Powers (PoQPo).

$$\left(\frac{a^m}{b^n} \right)^k = \frac{a^{m \cdot k}}{b^{n \cdot k}} \quad (1.11a)$$

$$\frac{a^{m \cdot k}}{b^{n \cdot k}} = \left(\frac{a^m}{b^n} \right)^k \quad (1.11b)$$




Figure 1.1: Simplifying Expressions Workflow:

■ Property, ■ Operation, ■ Notation, ■ Powers, ■ Delimiters, ■ Process, ■ Not Used

Rule 1.2.12 — Power of a Product of Powers (PoPrPo).

$$(a^m \cdot b^n)^k = a^{m \cdot k} \cdot b^{n \cdot k} \quad (1.12a)$$

$$a^{m \cdot k} \cdot b^{n \cdot k} = (a^m \cdot b^n)^k \quad (1.12b)$$

Definition 1.2.13 — Power To Factor (PoTF).

$$a^{\textcolor{red}{n}} = a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_{\textcolor{red}{n}} \quad (1.13)$$

Definition 1.2.14 — Factor To Power (FTPo).

$$a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_{\textcolor{red}{n}} = a^{\textcolor{red}{n}} \quad (1.14)$$

Definition 1.2.15 — Power Inverse (Pol).

$$(b^m)^{\frac{1}{m}} = b \quad (1.15a)$$

Definition 1.2.16 — Power Inverse (Pold).

$$1 = b^0 \quad (1.16a)$$

$$b^0 = 1 \quad (1.16b)$$

Notation 1.1 (Radical To Power (RTPo)).

$$\sqrt[m]{b^n} = b^{\frac{n}{m}} \quad (1.17)$$

Notation 1.2 (Power To Radical (PoTR)).

$$b^{\frac{n}{m}} = \sqrt[m]{b^n} \quad (1.18)$$

1.2.1 Monomials of Like Terms

Definition 1.2.17 — Additive Inverse (AI).

$$a + (-a) = 0 \quad (1.19a)$$

1.2.2 Surds

Example 1.1 — id:20141108-085327.

Simplify $2\sqrt{2} - \frac{(\sqrt{2})^3}{3} - \left(2(-\sqrt{2}) - \frac{(-\sqrt{2})^3}{3} \right)$

S

Solution:

$$2\sqrt{2} - \frac{1(1\sqrt{2})^3}{3} - 1 \left(2(-1\sqrt{2}) - \frac{1(-1\sqrt{2})^3}{3} \right) \quad \text{MId}(2.1a)$$

$$2\sqrt{2} - \frac{1(1\sqrt{2})^3}{3} - 1 \left(2(\neg 1\sqrt{2}) - \frac{1(\neg 1\sqrt{2})^3}{3} \right) \quad \text{ONeg}(2.6a)$$

$$2\sqrt{2} + \frac{\neg 1(1\sqrt{2})^3}{3} + \neg 1 \left(2(\neg 1\sqrt{2}) + \frac{\neg 1(\neg 1\sqrt{2})^3}{3} \right) \quad \text{DOS}(2.5a)$$

$$2 \cdot 2^{1/2} + \frac{\neg 1(1 \cdot 2^{1/2})^3}{3} + \neg 1 \left(2(\neg 1 \cdot 2^{1/2}) + \frac{\neg 1(\neg 1 \cdot 2^{1/2})^3}{3} \right) \quad \text{RTPo}(1.17)$$

$$2 \cdot 2^{1/2} + \frac{\neg 1(1 \cdot 2^{1/2})^3}{3} + \neg 1 \left(2 \cdot \neg 1 \cdot 2^{1/2} + \frac{\neg 1(\neg 1 \cdot 2^{1/2})^3}{3} \right) \quad \text{JTC}(1.6)$$

$$2 \cdot 2^{1/2} + \frac{\neg 1 \cdot 1 \cdot 2^{3/2}}{3} + \neg 1 \left(2 \cdot \neg 1 \cdot 2^{1/2} + \frac{\neg 1 \cdot \neg 1 \cdot 2^{3/2}}{3} \right) \quad \text{PoPrPo}(1.12a)$$

$$2 \cdot 2^{1/2} + \frac{\neg 1 \cdot 1 \cdot 2^{2/2} \cdot 2^{1/2}}{3} + \neg 1 \left(2 \cdot \neg 1 \cdot 2^{1/2} + \frac{\neg 1 \cdot \neg 1 \cdot 2^{2/2} \cdot 2^{1/2}}{3} \right) \quad \text{PrCBPo}(2.10b)$$

$$2 \cdot 2^{1/2} + \frac{\neg 1 \cdot 1 \cdot 2 \cdot 2^{1/2}}{3} + \neg 1 \left(2 \cdot \neg 1 \cdot 2^{1/2} + \frac{\neg 1 \cdot \neg 1 \cdot 2 \cdot 2^{1/2}}{3} \right) \quad \text{MId}(2.1b)$$

$$2 \cdot \sqrt{2} + \frac{\neg 1 \cdot 1 \cdot 2 \cdot \sqrt{2}}{3} + \neg 1 \left(2 \cdot \neg 1 \cdot \sqrt{2} + \frac{\neg 1 \cdot \neg 1 \cdot 2 \cdot \sqrt{2}}{3} \right) \quad \text{PoTR}(1.18)$$

$$2 \cdot \sqrt{2} + \frac{\neg 1 \cdot 1 \cdot 2 \cdot \sqrt{2}}{3} + \neg 1 \cdot 2 \cdot \neg 1 \cdot \sqrt{2} + \frac{\neg 1 \cdot \neg 1 \cdot \neg 1 \cdot 2 \cdot \sqrt{2}}{3} \quad \text{DPE}(2.9a)$$

$$2 \cdot \sqrt{2} + \frac{\neg 2 \cdot \sqrt{2}}{3} + 2 \cdot \sqrt{2} + \frac{\neg 2 \cdot \sqrt{2}}{3} \quad \text{OOM}(1.4)$$

$$2\sqrt{2} + \frac{\neg 2\sqrt{2}}{3} + 2\sqrt{2} + \frac{\neg 2\sqrt{2}}{3} \quad \text{CTJ}(1.7)$$

$$\left(2 + \frac{\neg 2}{3} + 2 + \frac{\neg 2}{3} \right) \sqrt{2} \quad \text{DPF}(2.4a)$$

$$\frac{8}{3}\sqrt{2} \quad \text{OOA}(1.1)$$

D

Dependencies:example 8.5-20141108-083108



Classification of Univariate Polynomial Expressions

- Degree -1 Univariate Polynomials
- Degree 0 Univariate Polynomials
- Degree 1 Univariate Polynomials
- Degree 2 Univariate Polynomials
- Degree 3 Univariate Polynomials
- Degree n Univariate Polynomials

2. Univariate Polynomial Expressions

2.1 Classification of Univariate Polynomial Expressions

Definition 2.1.1 — Indeterminate.

x

An indeterminate is a symbol that is treated as a variable, but does not stand for anything else but itself and is used as a placeholder.

- it does **not** designate a constant or a parameter
- it is **not** an unknown that could be solved for
- it is **not** a variable designating a function argument

[[wikipedia:indeterminate](#)]

Definition 2.1.2 — Coefficient.

Cx^k

A coefficient, C is a real number multiplicative factor.

Definition 2.1.3 — Univariate Monomial.

$C_k x^k$

A univariate monomial is made up of two factors. The first factor of a monomial, C_k , is the **coefficient**. The second factor of each monomial, x^k , is an indeterminate raised to a non-negative integer power k .

Definition 2.1.4 — Multiplicative Identity (MId).

$$\textcolor{violet}{1}a = a \quad (2.1\text{a})$$

$$a = \textcolor{violet}{1}a \quad (2.1\text{b})$$

- R** If the coefficient of a univariate monomial is the multiplicative identity 2.1a, 1, then it is not shown in its canonical form.

$$\begin{aligned}\textcolor{teal}{C}_k x^k &= \textcolor{teal}{C}_k x^k \\ &= \textcolor{violet}{1}x^k \\ &= x^k\end{aligned}$$

Example 2.1 — id:20141120-202842.

Express $1x^2$ in canonical form.

S

Solution:

$$x^2 \quad \text{MId}(2.1\text{b})$$

**Definition 2.1.5 — Univariate Polynomial Expression.**

$$\sum_{k=0}^n \textcolor{teal}{C}_k x^n = \textcolor{teal}{C}_n x^n + \textcolor{teal}{C}_{n-1} x^{n-1} + \cdots + \textcolor{teal}{C}_k x^k + \cdots + \textcolor{teal}{C}_2 x^2 + \textcolor{teal}{C}_1 x^1 + \underbrace{\textcolor{teal}{C}_0}_{\textcolor{teal}{C}_0 x^0} \quad (2.2)$$

A univariate polynomial in an indeterminate x is an expression made up of one or more summands of the form $\textcolor{teal}{C}_k x^k$, which are called monomials. The first factor of each monomial, $\textcolor{teal}{C}_k$, is a numerical factor called the **coefficient** where $\textcolor{teal}{C}_k \in \mathbb{R}$. The second factor of each monomial, x^k , is an indeterminate raised to a non-negative integer power i .

[]

Definition 2.1.6 — Degree of the Indeterminate.

$$x^k$$

The exponent of an indeterminate power, k is called the degree of the indeterminate.

[]

Definition 2.1.7 — Degree of the Univariate Polynomial.

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0}$$

The degree of the univariate polynomial is determined by the monomial with the largest degree of the indeterminate.

2.2 Degree -1 Univariate Polynomials

Monomials

2.3 Degree 0 Univariate Polynomials

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0}$$

Degree 0 univariate polynomial expressions are made up of univariate monomials, C_0 , called **constants**. The power identity is an indeterminate raised to a power of 0 has a value of 1. Thus, $x^0 = 1$ and results in the monomial $C_0 \cdot 1$. The canonical form of a product does not show the multiplicative identity factor, so what remains of this monomial product is only the coefficient factor C_0 and from now on will be referred to as a **constant**.

Example 2.2 — id:20141121-093747.

Express $13x^0$ in canonical form.

S

Solution:

13



Monomials

Degree 0 univariate polynomial expressions are usually a monomial in their canonical form if C_0 is a non-zero real number. The exception is if $C_0 = 0$, the additive identity, then the result is the zero polynomial, which can be considered a degree -1 polynomial.

The expression can be manipulated into its monomial canonical form by simplifying the expression. Simplifying the expression can be defined as evaluating the expression by following order of operations, which is the same as evaluating an arithmetic expression.

2.4 Degree 1 Univariate Polynomials

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0}$$

Degree 1 univariate polynomial expressions can be expressed with at most two different terms and consequently this expression in its canonical form has at most two monomial summands – called a binomial.

Definition 2.4.1 — Univariate Like Terms.

$$C_1x^k = C_2x^k$$

Two or more univariate monomials are defined as having like terms if each monomial has the same term, which will be the same indeterminate raised to the same positive integer power.

Sometimes the word **term** is used to describe monomials (including both the coefficient and the term), which may be confusing when trying to define like terms. For this reason, we will refer to the summands of a polynomial as monomials.

The monomials $5x^1$ and $3x^1$ can be described as having like terms because they share the common term x^1 . One could also say that $5x^1$ and $3x^1$ are like terms by definition and consequently giving the reader an impression that $5x^1$ and $3x^1$ are terms themselves.

- R** A degree 1 indeterminate does not display the multiplicative identity in the exponent when it's in canonical form.

Example 2.3 — id:20141120-202042.

Express $5x^1$ in canonical form

S

Solution:

$$5x \quad \text{MId}(2.1b)$$

Example 2.4 — id:20141121-093439.

Express $7x^1 + 5$ in canonical form.

S

Solution:

$$7x + 5 \quad \text{MId}(2.1b)$$

Monomials

Essential Questions 2.1

- How do we simplify univariate polynomial expressions?

Definition 2.4.2 — Additive Identity (Ald).

$$a + 0 = a \quad (2.3a)$$

$$a = a + 0 \quad (2.3b)$$

If the constant monomial is 0, the additive identity, then the canonical form of a degree 1 univariate polynomial is a degree 1 monomial.

Simplifying Univariate Monomial Expressions

The definition of a univariate monomial expression is based on the expanded canonical form of some polynomial expression. It might be that the original expression might not be in the expanded canonical form, so a process called **simplifying by expanding** will be introduced to manipulate the expression such that it can be written in its expanded canonical form.

This process of simplifying by expanding polynomial expressions will be developed to the extent that it will be used to simplify multivariate polynomials. We will start by simplifying univariate monomial expressions.

Definition 2.4.3 — Distributive Property Factoring (DPF).

$$ba + ca = (b + c)a \quad (2.4a)$$

$$ab + ac = a(b + c) \quad (2.4b)$$

Example 2.5 — id:20141120-203846.

Simplify $6x + 7x$

S

Solution:

Notice that the indeterminate of each monomial is of degree 1; however, the exponent 1 is not shown. The monomials $6x$ and $7x$ have a like term of x .

$$(6 + 7)x$$

$$13x$$

DPF(2.4a)

OOA(1.1)

Notice that the sum of two monomials that have like terms can be found by adding the coefficients of the monomials. The distributive property in the factoring direction provides some insight to why we can add the coefficients of monomials that have like terms.

S**Less Steps Solution:**

$$13x \quad \text{OOA(1.1)}$$

Exercise 2.1 — id:20141120-211843.

Simplify the following univariate polynomial expressions

1. $7x + 10x$

2. $5x + 3x$

3. $11x + 11x$

4. $3x + 6x$

5. $7x + 2x$

6. $12x + 10x$

7. $7x + 4x$

8. $6x + 2x$

9. $7x + 11x$

10. $8x + 11x$

11. $13x + 13x$

12. $4x + 4x$

Answers: 9.0.1

Example 2.6 — id:20141027-075159.

Simplify $7\text{ cm} + 8\text{ cm}$

S**Solution:**

$(7 + 8) \text{ cm}$

15 cm

DPF(2.4a)

OOA(1.1)

R

Remember, if a monomial does not have a coefficient factor, then it's implied that the coefficient factor is 1, the multiplicative identity, and consequently its not explicitly shown.

Example 2.7 — id:20141121-185558.

Simplify $x + 5x$

S**Solution:**

It can be useful when simplifying expressions to make the multiplicative identity (MId) factor explicit.

$1x + 5x$	MId(2.1a)
$(1 + 5)x$	DPF(2.4a)
$6x$	OOA(1.1)

S**Less Steps Solution:**

$1x + 5x$	MId(2.1a)
$6x$	OOA(1.1)

As one becomes more experienced, there is no reason to make the multiplicative identity coefficient explicit.

S**Less Steps Solution:**

$6x$	OOA(1.1)
------	----------

**Exercise 2.2 — id:20141121-195608.**

Simplify the following univariate polynomial expressions

- | | | |
|--------------|--------------|---------------|
| 1. $x + 12x$ | 5. $x + 12x$ | 9. $12x + x$ |
| 2. $x + 5x$ | 6. $x + 3x$ | 10. $6x + x$ |
| 3. $x + 7x$ | 7. $2x + x$ | 11. $13x + x$ |
| 4. $x + 10x$ | 8. $7x + x$ | 12. $4x + x$ |

Answers: 9.0.2

**Definition 2.4.4 — Definition of Subtraction (DOS).**

$$a - b = a + \neg b \quad (2.5a)$$

$$a + \neg b = a - b \quad (2.5b)$$

Notation 2.1 (Operation of Negation (ONeg)).

$$\neg a = \neg a \quad (2.6a)$$

$$\neg a = -a \quad (2.6b)$$

I have used a different symbol, \neg , as the prefix negation operator only to differentiate it from the minus sign infix operator symbol, $-$, which is also used as the infix operator for the dyadic operation of subtraction. I will refer to this change of symbol as ONeg. This is used only as a teaching tool and should not be confused with the logic negation operator. Another advantage of

using this symbol is that it reduces the number of delimiters used in an expression for example, $\neg a$ versus $(-\bar{a})$.

- Negative five: -5
- Negative five: $\neg 5$
- Four minus five: $4 - 5$
- Four minus negative five: $4 - \neg 5$
- Four minus negative five: $4 - (-\bar{5})$
- Four minus negative five: $4 - \bar{5}$
- Negative four minus five: $-4 - 5$
- Negative four minus five: $\neg 4 - 5$

Example 2.8 — id:20141121-190857.

Simplify $8x - 6x$

S

Solution:

$$\begin{array}{ll} 8x + \neg 6x & \text{DOS(2.5a)} \\ (8 + \neg 6)x & \text{DPF(2.4a)} \\ 2x & \text{OOA(1.1)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} 8x + \neg 6x & \text{DOS(2.5a)} \\ 2x & \text{OOA(1.1)} \end{array}$$



Example 2.9 — id:20141121-193636.

Simplify $3x - 5x$

S

Solution:

$$\begin{array}{ll} 3x + \neg 5x & \text{DOS(2.5a)} \\ (3 + \neg 5)x & \text{DPF(2.4a)} \\ \neg 2x & \text{OOA(1.1)} \\ -2x & \text{ONeg(2.6b)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} 3x + -5x & \text{DOS(2.5a)} \\ -2x & \text{OOA(1.1)} \\ -2x & \text{ONeg(2.6b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} -2x & \text{OOA(1.1)} \end{array}$$

■

Example 2.10 — id:20141106-150622.Simplify $13x - x$ **S****Solution:**

$$\begin{array}{ll} 13x - 1x & \text{MId(2.1a)} \\ 13x + -1x & \text{DOS(2.5a)} \\ (13 + -1)x & \text{DPF(2.4a)} \\ 12x & \text{OOA(1.1)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} 13x + -x & \text{DOS(2.5a)} \\ 12x & \text{OOA(1.1)} \end{array}$$

■

Exercise 2.3 — id:20141121-191224.

Simplify the following univariate polynomial expressions

- | | | |
|---------------|---------------|----------------|
| 1. $7x - 8x$ | 5. $9x - 8x$ | 9. $10x - 12x$ |
| 2. $12x - 9x$ | 6. $13x - 4x$ | 10. $8x - 3x$ |
| 3. $2x - 3x$ | 7. $4x - 9x$ | 11. $2x - 8x$ |
| 4. $3x - 9x$ | 8. $2x - 12x$ | 12. $6x - 9x$ |

Answers: 9.0.3

■

It is possible for a univariate monomial to have more than two terms in its non-canonical form. The associative property of addition will be used to help simplify these expressions.

Definition 2.4.5 — Associative Property of Addition (APA).

$$a + b + c = (a + b) + c \quad (2.7a)$$

$$a + b + c = a + (b + c) \quad (2.7b)$$

Example 2.11 — id:20141121-184652.

Simplify the expression $3x + 7x + 8x$

S

$$\begin{array}{ll} (3x + 7x) + 8x & \text{APA(2.7a)} \\ (3 + 7)x + 8x & \text{DPF(2.4a)} \\ 10x + 8x & \text{OOA(1.1)} \\ (10 + 8)x & \text{DPF(2.4a)} \\ 18x & \text{OOA(1.1)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} (3x + 7x) + 8x & \text{APA(2.7a)} \\ 10x + 8x & \text{OOA(1.1)} \\ 18x & \text{OOA(1.1)} \end{array}$$

You might have noticed that this expression could be simplified in one step by adding the coefficient of the three monomials $3x$, $7x$ and $8x$, which have the like term x .

S

Less Steps Solution:

$$18x \quad \text{OOA(1.1)}$$



Example 2.12 — id:20141106-152020.

Simplify $4x - 2x - x$

S

Solution:

$$\begin{array}{ll}
 4x - 2x - 1x & \text{MId(2.1a)} \\
 4x + -2x + -1x & \text{DOS(2.5a)} \\
 (4 + -2)x + -1x & \text{DPF(2.4a)} \\
 2x + -1x & \text{OOA(1.1)} \\
 (2 + -1)x & \text{DPF(2.4a)} \\
 1x & \text{OOA(1.1)} \\
 x & \text{MId(2.1b)}
 \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll}
 4x + -2x + -1x & \text{DOS(2.5a)} \\
 x & \text{OOA(1.1)}
 \end{array}$$

Exercise 2.4 — id:20141123-091420.

Simplify the following univariate polynomial expressions

- | | | |
|------------------------|----------------------|-----------------------|
| 1. $4x + 10x + -5x$ | 5. $5x + -9x + 8x$ | 9. $-10x + 9x + -4x$ |
| 2. $-7x + -13x + -10x$ | 6. $-5x + -10x + 5x$ | 10. $-2x + -3x + 12x$ |
| 3. $-10x + 10x + -3x$ | 7. $6x + -12x + -3x$ | 11. $9x + -5x + -8x$ |
| 4. $-13x + -9x + 8x$ | 8. $-8x + 11x + 7x$ | 12. $8x + 11x + -3x$ |

Answers: 9.0.4

Example 2.13 — id:20141108-194431.

Simplify $-3 \cdot 7x - 2x \cdot 4$

S

Solution:

$\neg 3 \cdot 7x - 2x \cdot 4$	ONeg(2.6a)
$\neg 3 \cdot 7x + \neg 2x \cdot 4$	DOS(2.5a)
$\neg 3 \cdot 7 \cdot x + \neg 2 \cdot x \cdot 4$	JTC(1.6)
$\neg 3 \cdot 7 \cdot x + \neg 2 \cdot 4 \cdot x$	CPM(1.8)
$(\neg 3 \cdot 7) \cdot x + (\neg 2 \cdot 4) \cdot x$	APM(1.10a)
$\neg 21 \cdot x + \neg 8 \cdot x$	OOM(1.4)
$\neg 21x + \neg 8x$	CTJ(1.7)
$(\neg 21 + \neg 8)x$	DPF(2.4a)
$\neg 29x$	OOA(1.1)
$\neg 29x$	ONeg(2.6b)

S**Less Steps Solution:**

$\neg 3 \cdot 7x + \neg 2x \cdot 4$	DOS(2.5a)
$\neg 3 \cdot 7 \cdot x + \neg 2 \cdot 4 \cdot x$	CPM(1.8)
$\neg 21x + \neg 8x$	OOM(1.4)
$\neg 29x$	OOA(1.1)

**Example 2.14 — id:20141108-194156.**Simplify $3 \cdot 5x + 3x \cdot 4$ **S****Solution:**

$3 \cdot 5 \cdot x + 3 \cdot x \cdot 4$	JTC(1.6)
$3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x$	CPM(1.8)
$(3 \cdot 5) \cdot x + (3 \cdot 4) \cdot x$	APM(1.10a)
$15 \cdot x + 12 \cdot x$	OOM(1.4)
$15x + 12x$	CTJ(1.7)
$(15 + 12)x$	DPF(2.4a)
$27x$	OOA(1.1)

S

Less Steps Solution:

$$\begin{array}{ll} 3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x & \text{CPM(1.8)} \\ 15x + 12x & \text{OOM(1.4)} \\ 27x & \text{OOA(1.1)} \end{array}$$

**Example 2.15 — id:20141108-173613.**

Simplify $8x \cdot 5$

S

Solution:

$$\begin{array}{ll} 8 \cdot x \cdot 5 & \text{JTC(1.6)} \\ 8 \cdot 5 \cdot x & \text{CPM(1.8)} \\ (8 \cdot 5) \cdot x & \text{APM(1.10a)} \\ 40 \cdot x & \text{OOM(1.4)} \\ 40x & \text{CTJ(1.7)} \end{array}$$

**Binomials****Definition 2.4.6 — Commutative Property of Addition (CPA).**

$$ab = ba \tag{2.8}$$

Definition 2.4.7 — Distributive Property Expanding (DPE).

$$a(b + c) = ab + ac \tag{2.9a}$$

$$(b + c)a = b a + c a \tag{2.9b}$$

Example 2.16 — id:20141109-090809.

Simplify by expanding $5(x + 4)$

S

Solution:

$$\begin{array}{ll}
 5(1x + 4) & \text{MId(2.1a)} \\
 5 \cdot 1x + 5 \cdot 5 & \text{DPF(2.4a)} \\
 5 \cdot 1 \cdot x + 5 \cdot 5 & \text{JTC(1.6)} \\
 5 \cdot x + 25 & \text{OOM(1.4)} \\
 5x + 25 & \text{CTJ(1.7)}
 \end{array}$$

S

Less Steps Solution:

$$5x + 20 \quad \text{DPE(2.9a)}$$



Example 2.17 — id:20141109-091015.

Simplify by expanding $5(3x - 9)$

S

Solution:

$$\begin{array}{ll}
 5(3x + -9) & \text{DOS(2.5a)} \\
 5 \cdot 3x + 5 \cdot -9 & \text{DPE(2.9a)} \\
 5 \cdot 3 \cdot x + 5 \cdot -9 & \text{JTC(1.6)} \\
 15 \cdot x + -45 & \text{OOM(1.4)} \\
 15x + -45 & \text{CTJ(1.7)} \\
 15x - 45 & \text{DOS(2.5b)}
 \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll}
 5(3x + -9) & \text{DOS(2.5a)} \\
 15x + -45 & \text{DPE(2.9a)} \\
 15x - 45 & \text{DOS(2.5b)}
 \end{array}$$



Example 2.18 — id:20141109-092448.

Simplify by expanding $-(5x + 7)$

S**Solution:**

$$\begin{array}{ll} \neg 1(5x + 7) & \text{MId(2.1a)} \\ \neg 1 \cdot 5x + \neg 1 \cdot 7 & \text{DPE(2.9a)} \\ \neg 1 \cdot 5 \cdot x + \neg 1 \cdot 7 & \text{JTC(1.6)} \\ \neg 5 \cdot x + \neg 7 & \text{OOM(1.4)} \\ \neg 5x + \neg 7 & \text{CTJ(1.7)} \\ \neg 5x - 7 & \text{DOS(2.5b)} \\ -5x - 7 & \text{ONeg(2.6b)} \end{array}$$

S**Less Steps Solution:**

$$-5x - 7 \quad \text{DPE(2.9a)}$$

**Example 2.19 — id:20141109-092651.**Simplify by expanding $-13(7x - 9)$ **S****Solution:**

$$\begin{array}{ll} \neg 13(7x + \neg 9) & \text{DOS(2.5a)} \\ \neg 13 \cdot 7x + \neg 13 \cdot \neg 9 & \text{DPE(2.9a)} \\ \neg 13 \cdot 7 \cdot x + \neg 13 \cdot \neg 9 & \text{JTC(1.6)} \\ \neg 91 \cdot x + 117 & \text{OOM(1.4)} \\ \neg 91x + 117 & \text{CTJ(1.7)} \\ -91x + 117 & \text{ONeg(2.6b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} -13(7x + \neg 9) & \text{DOS(2.5a)} \\ -91x + 117 & \text{DPE(2.9a)} \end{array}$$



Example 2.20 — id:20141109-092910.

Simplify by expanding $a(x + b)$, where $a, b \in \mathbb{Z}$

S

Solution:

$$\begin{array}{ll} a(1x + b) & \text{MId(2.1a)} \\ a \cdot 1x + a \cdot b & \text{DPE(2.9a)} \\ a \cdot 1 \cdot x + a \cdot b & \text{JTC(1.6)} \\ 1 \cdot a \cdot x + a \cdot b & \text{CPM(1.8)} \\ 1ax + ab & \text{JTC(1.6)} \\ ax + ab & \text{MId(2.1b)} \end{array}$$

S

Less Steps Solution:

$$ax + ab \quad \text{DPE(2.9a)}$$

**Example 2.21 — id:20141109-093220.**

Simplify by expanding $5(x + 2) + 4$

S

Solution:

$$\begin{array}{ll} 5(1x + 2) + 4 & \text{MId(2.1a)} \\ 5 \cdot 1x + 5 \cdot 2 + 4 & \text{DPE(2.9a)} \\ 5 \cdot 1 \cdot x + 5 \cdot 2 + 4 & \text{JTC(1.6)} \\ 5 \cdot x + 10 + 4 & \text{OOM(1.4)} \\ 5x + 10 + 4 & \text{CTJ(1.7)} \\ 5x + 14 & \text{OOA(1.1)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} 5x + 10 + 4 & \text{DPE(2.9a)} \\ 5x + 14 & \text{OOA(1.1)} \end{array}$$

**Example 2.22 — id:20141109-093419.**

Simplify by expanding $7x + 5(4x + 8)$

S**Solution:**

$$\begin{array}{ll} 7x + 5 \cdot 4x + 5 \cdot 8 & \text{DPE(2.9a)} \\ 7 \cdot x + 5 \cdot 4 \cdot x + 5 \cdot 8 & \text{JTC(1.6)} \\ 7 \cdot x + 20 \cdot x + 40 & \text{OOM(1.4)} \\ 7x + 20x + 40 & \text{CTJ(1.7)} \\ (7 + 20)x + 40 & \text{DPF(2.4a)} \\ 27x + 40 & \text{OOA(1.1)} \end{array}$$

**Example 2.23 — id:20141109-094928.**

Simplify by expanding $4(3x + 4) + x + 6$

S**Solution:**

$$\begin{array}{ll} 4(3x + 4) + 1x + 6 & \text{MId(2.1a)} \\ 4 \cdot 3x + 4 \cdot 4 + 1x + 6 & \text{DPE(2.9a)} \\ 4 \cdot 3 \cdot x + 4 \cdot 4 + 1 \cdot x + 6 & \text{JTC(1.6)} \\ 12 \cdot x + 16 + 1 \cdot x + 6 & \text{OOM(1.4)} \\ 12x + 16 + 1x + 6 & \text{CTJ(1.7)} \\ 12 + 1x + 16 + 6 & \text{CPA(2.8)} \\ (12 + 1)x + 16 + 6 & \text{DPF(2.4a)} \\ 13x + 22 & \text{OOA(1.1)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} 12x + 16 + x + 6 & \text{DPE(2.9a)} \\ 12x + x + 16 + 6 & \text{CPA(2.8)} \\ 13x + 22 & \text{OOA(1.1)} \end{array}$$

Example 2.24 — id:20141109-095151.

Simplify by expanding $5(x - 4) + 3x - 5$

S**Solution:**

$$\begin{array}{ll} 5(1x - 4) + 3x - 5 & \text{MId(2.1a)} \\ 5(1x + \neg 4) + 3x + \neg 5 & \text{DOS(2.5a)} \\ 5 \cdot 1x + 5 \cdot \neg 4 + 3x + \neg 5 & \text{DPE(2.9a)} \\ 5 \cdot 1 \cdot x + 5 \cdot \neg 4 + 3 \cdot x + \neg 5 & \text{JTC(1.6)} \\ 5 \cdot x + \neg 20 + 3 \cdot x + \neg 5 & \text{OOM(1.4)} \\ 5x + \neg 20 + 3x + \neg 5 & \text{JTC(1.6)} \\ 5x + 3x + \neg 20 + \neg 5 & \text{CPA(2.8)} \\ (5 + 3)x + \neg 20 + \neg 5 & \text{DPF(2.4a)} \\ 8x + \neg 25 & \text{OOA(1.1)} \\ 8x - 25 & \text{DOS(2.5b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} 5(x + \neg 4) + 3x + \neg 5 & \text{DOS(2.5a)} \\ 5x + \neg 20 + 3x + \neg 5 & \text{DPE(2.9a)} \\ 5x + 3x + \neg 20 + \neg 5 & \text{CPA(2.8)} \\ 8x + \neg 25 & \text{OOA(1.1)} \\ 8x - 25 & \text{DOS(2.5b)} \end{array}$$

Example 2.25 — id:20141109-095536.

Simplify by expanding $8x - 5 - 4(x - 3)$

S

Solution:

$8x - 5 - 4(1x - 3)$	MId(2.1a)
$8x + -5 + -4(1x + -3)$	DOS(2.5a)
$8x + -5 + -4 \cdot 1x + -4 \cdot -3$	DPE(2.9a)
$8 \cdot x + -5 + -4 \cdot 1 \cdot x + -4 \cdot -3$	JTC(1.6)
$8 \cdot x + -5 + -4 \cdot x + 12$	OOM(1.4)
$8x + -5 + -4x + 12$	CTJ(1.7)
$8x + -4x + -5 + 12$	CPA(2.8)
$(8 + -4)x + -5 + 12$	DPF(2.4a)
$4x + 7$	OOA(1.1)

S**Less Steps Solution:**

$8x + -5 + -4(x + -3)$	DOS(2.5a)
$8x + -5 + -4x + 12$	DPE(2.9a)
$8x + -4x + -5 + 12$	CPA(2.8)
$4x + 7$	OOA(1.1)

■

Example 2.26 — id:20141109-095842.Simplify by expanding $5(x + 3) + 3(x + 2)$ **S****Solution:**

$5 \cdot x + 5 \cdot 3 + 3 \cdot x + 3 \cdot 2$	DPE(2.9a)
$5 \cdot x + 15 + 3 \cdot x + 6$	OOM(1.4)
$5x + 15 + 3x + 6$	CTJ(1.7)
$5x + 3x + 15 + 6$	CPA(2.8)
$(3 + 5)x + 15 + 6$	DPF(2.4a)
$8x + 21$	OOA(1.1)

S

Less Steps Solution:

$$5x + 15 + 3x + 6$$

DPE(2.9a)

$$5x + 3x + 15 + 6$$

CPA(2.8)

$$8x + 21$$

OOA(1.1)

2.5 Degree 2 Univariate Polynomials**Monomials****Example 2.27 — id:20141106-151138.**Simplify $4x^2 + 12x^2$ **S****Solution:**

$$(4 + 12)x^2$$

DPF(2.4a)

$$16x^2$$

OOA(1.1)

S**Less Steps Solution:**

$$16x^2$$

OOA(1.1)

Example 2.28 — id:20141106-154547.Simplify $x^2 - x + x^2 + x$ **S****Solution:**

$$1x^2 - 1x + 1x^2 + 1x$$

MId(2.1a)

$$1x^2 + -1x + 1x^2 + 1x$$

DOS(2.5a)

$$1x^2 + 1x^2 + -1x + 1x$$

CPA(2.8)

$$(1 + 1)x^2 + (-1 + 1)x$$

DPF(2.4a)

$$2x^2 + 0x$$

OOA(1.1)

$$2x^2$$

MId(2.1b)

S

Less Steps Solution:

$$\begin{array}{ll} x^2 + \neg x + x^2 + x & \text{DOS(2.5a)} \\ x^2 + x^2 + \neg x + x & \text{CPA(2.8)} \\ 2x^2 & \text{OOA(1.1)} \end{array}$$

■

Example 2.29 — id:20141108-194709.Simplify $-2x^4x - x \cdot -x^3$

S

Solution:

$$\begin{array}{ll} -2x^4x - 1x \cdot 1x^3 & \text{MId(2.1a)} \\ -2x^4x - 1x \cdot 1x^3 & \text{ONeg(2.6a)} \\ -2 \cdot x \cdot 4 \cdot x + \neg 1 \cdot x \cdot 1 \cdot x \cdot 3 & \text{JTC(1.6)} \\ -2 \cdot 4 \cdot x \cdot x + \neg 1 \cdot \neg 1 \cdot 3 \cdot x \cdot x & \text{CPM(1.8)} \\ -2 \cdot 4 \cdot x^2 + \neg 1 \cdot \neg 1 \cdot 3 \cdot x^2 & \text{PrCBPo(2.10a)} \\ (-2 \cdot 4) \cdot x^2 + (\neg 1 \cdot \neg 1 \cdot 3) x^2 & \text{APM(1.10a)} \\ -8x^2 + 3x^2 & \text{OOM(1.4)} \\ (-8 + 3)x^2 & \text{DPF(2.4a)} \\ -5x^2 & \text{OOA(1.1)} \\ -5x^2 & \text{ONeg(2.6b)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} -2x^4x + \neg x \cdot -x^3 & \text{DOS(2.5a)} \\ -2 \cdot 4 \cdot x \cdot x + 3 \cdot \neg x \cdot \neg x & \text{CPM(1.8)} \\ -2 \cdot 4 \cdot x^2 + 3 \cdot x^2 & \text{PrCBPo(2.10a)} \\ -8x^2 + 3x^2 & \text{OOM(1.4)} \\ -5x^2 & \text{OOA(1.1)} \end{array}$$

■

Rule 2.5.1 — Product of a Common Base Powers (PrCBPo).

$$b^m \cdot b^n = b^{m+n} \tag{2.10a}$$

$$b^{m+n} = b^m \cdot b^n \tag{2.10b}$$

Rule 2.5.2 — Quotient of a Common Base Powers (QCBPo).

$$\frac{b^m}{b^n} = b^{m-n} \quad (2.11a)$$

$$b^{m-n} = \frac{b^m}{b^n} \quad (2.11b)$$

Rule 2.5.3 — Power of a Power (PoPo).

$$(b^m)^k = b^{m \cdot k} \quad (2.12a)$$

$$b^{m \cdot k} = (b^m)^k \quad (2.12b)$$

Example 2.30 — id:20141108-191616.

Simplify $-5x \cdot 4x$

S

Solution:

$-5x \cdot 4x$	ONeg(2.6a)
$\neg 5 \cdot x \cdot 4 \cdot x$	JTC(1.6)
$\neg 5 \cdot 4 \cdot x \cdot x$	CPM(1.8)
$\neg 5 \cdot 4 \cdot x^2$	PrCBPo(2.10a)
$(\neg 5 \cdot 4) \cdot x^2$	APM(1.10a)
$\neg 20 \cdot x^2$	OOM(1.4)
$\neg 20x^2$	CTJ(1.7)
$-20x^2$	ONeg(2.6b)

S

Less Steps Solution:

$\neg 5 \cdot 4 \cdot x \cdot x$	CPM(1.8)
$\neg 5 \cdot 4 \cdot x^2$	PrCBPo(2.10a)
$-20x^2$	OOM(1.4)

Binomials**Example 2.31 — id:20141106-152339.**

Simplify $3x^2 + 2x + 5x^2 + 4x$

S**Solution:**

$$3x^2 + 5x^2 + 2x + 4x \quad \text{CPA(2.8)}$$

$$(3 + 5)x^2 + (2 + 4)x \quad \text{DPF(2.4a)}$$

$$8x^2 + 6x \quad \text{OOA(1.1)}$$

If needed we could continue and express it in the simplified factored form using the distributive property

$$(4x + 3)2x \quad \text{DPF(2.4a)}$$

S**Less Steps Solution:**

$$3x^2 + 5x^2 + 2x + 4x \quad \text{CPA(2.8)}$$

$$8x^2 + 6x \quad \text{OOA(1.1)}$$

■

Example 2.32 — id:20141107-121834.

Simplify $(\sqrt{9 - x^2})^2$

S**Solution:**

$$(\sqrt{9 - x^2})^2 \quad \text{MId(2.1a)}$$

$$(\sqrt{9 + -1x^2})^2 \quad \text{DOS(2.5a)}$$

$$\left[(9 + -1x^2)^{\frac{1}{2}} \right]^2 \quad \text{RTPo(1.17)}$$

$$9 + -1x^2 \quad \text{PoPo(2.12a)}$$

$$-1x^2 + 9 \quad \text{CPA(2.8)}$$

$$-x^2 + 9 \quad \text{MId(2.1a)}$$

$$-x^2 + 9 \quad \text{ONeg(2.6b)}$$

S

Less Steps Solution:

$$9 - x^2 \quad \text{PoPo(2.12a)}$$

It might be easier to view this using a variable substitution for the radicand, $9 - x^2$. Let $k = 9 + -1x^2$.

$$\begin{aligned} & (\sqrt{k})^2 && \text{MId(2.1a)} \\ & (\sqrt{k})^2 && \text{DOS(2.5a)} \\ & [(k)^{\frac{1}{2}}]^2 && \text{RTPo(1.17)} \\ & k && \text{PoPo(2.12a)} \\ & 9 + -1x^2 && \text{CPA(2.8)} \\ & -1x^2 + 9 && \text{CPA(2.8)} \\ & -x^2 + 9 && \text{MId(2.1a)} \\ & -x^2 + 9 && \text{ONeg(2.6b)} \end{aligned}$$

D

Dependencies:example 8.2-20141105-144223

Example 2.33 — id:20141209-145211.

Simplify $2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0$

S**Solution:**

$$\begin{aligned} & 2x \cdot 2x + 2x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0 && \text{DPE(2.9a)} \\ & 2 \cdot x \cdot 2 \cdot x + 2 \cdot x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0 && \text{JTC(1.6)} \\ & 2 \cdot 2 \cdot x \cdot x + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0 && \text{CPM(1.8)} \\ & 2 \cdot 2 \cdot x^2 + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0 && \text{PrCBPo(2.10a)} \\ & 4 \cdot x^2 + 8 \cdot x + 2 \cdot x^2 + 0 && \text{OOM(1.4)} \\ & 4x^2 + 8x + 2x^2 + 0 && \text{CTJ(1.7)} \\ & 4x^2 + 2x^2 + 8x && \text{APA(2.7a)} \\ & (4 + 2)x^2 + 8x && \text{DPF(2.4b)} \\ & 6x^2 + 8x && \text{OOA(1.1)} \end{aligned}$$

D

Dependencies:example 7.6-20141209-144203

**Trinomials****Example 2.34 — id:20141109-133008.**

Simplify by expanding $(x + 5)(x - 8)$

S**Solution:**

$$\begin{aligned}
 & (1x + 5)(1x - 8) && \text{MId(2.1a)} \\
 & (1x + 5)(1x + -8) && \text{DOS(2.5a)} \\
 & 1x(1x + -8) + 5(1x + -8) && \text{DPE(2.9b)} \\
 & 1x \cdot 1x + 1x \cdot -8 + 5 \cdot 1x + 5 \cdot -8 && \text{DPE(2.9a)} \\
 & 1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot -8 + 5 \cdot 1 \cdot x + 5 \cdot -8 && \text{JTC(1.6)} \\
 & 1 \cdot 1 \cdot x \cdot x + -8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + -8 \cdot 5 && \text{CPM(1.8)} \\
 & 1 \cdot 1 \cdot x^2 + -8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + -8 \cdot 5 && \text{PrCBPo(2.10a)} \\
 & 1 \cdot x^2 + -8 \cdot x + 5 \cdot x + -40 && \text{OOM(1.4)} \\
 & 1x^2 + -8x + 5x + -40 && \text{CTJ(1.7)} \\
 & 1x^2 + -3x + -40 && \text{OOA(1.1)} \\
 & 1x^2 - 3x - 40 && \text{DOS(2.5b)} \\
 & x^2 - 3x - 40 && \text{MId(2.1a)}
 \end{aligned}$$

S**Less Steps Solution:**

$$\begin{aligned}
 & (x + 5)(x + -8) && \text{DOS(2.5a)} \\
 & x(x + -8) + 5(x + -8) && \text{DPE(2.9b)} \\
 & x^2 + -8x + 5x + -40 && \text{DPE(2.9a)} \\
 & x^2 - 3x - 40 && \text{OOA(1.1)}
 \end{aligned}$$

**Example 2.35 — id:20141109-133316.**

Simplify by expanding $(x + a)(x + b)$, where $a, b \in \mathbb{Z}$

S

Solution:

$$\begin{array}{ll}
 (1x + a)(1x + b) & \text{MId(2.1a)} \\
 1x(1x + b) + a(1x + b) & \text{DPE(2.9b)} \\
 1x \cdot 1x + 1x \cdot b + a \cdot 1x + a \cdot b & \text{DPE(2.9a)} \\
 1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot b + a \cdot 1 \cdot x + a \cdot b & \text{JTC(1.6)} \\
 1 \cdot 1 \cdot x \cdot x + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{CPM(1.8)} \\
 1 \cdot 1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{PrCBPo(2.10a)} \\
 1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{OOM(1.4)} \\
 1x^2 + 1bx + 1ax + ab & \text{CTJ(1.7)} \\
 1x^2 + (1b + 1a)x + ab & \text{DPF(2.4a)} \\
 x^2 + (b + a)x + ab & \text{MId(2.1b)}
 \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll}
 x(x + b) + a(x + b) & \text{DPE(2.9b)} \\
 x^2 + (b + a)x + ab & \text{DPE(2.9a)}
 \end{array}$$

■

Example 2.36 — id:20141109-140659.Simplify by expanding $(2x + 3)(5x + 13)$ **S****Solution:**

$$\begin{array}{ll}
 2x(5x + 13) + 3(5x + 13) & \text{DPE(2.9b)} \\
 2x \cdot 5x + 2x \cdot 13 + 3 \cdot 5x + 3 \cdot 13 & \text{DPE(2.9a)} \\
 2 \cdot x \cdot 5 \cdot x + 2 \cdot x \cdot 13 + 3 \cdot 5 \cdot x + 3 \cdot 13 & \text{JTC(1.6)} \\
 2 \cdot 5 \cdot x \cdot x + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13 & \text{CPM(1.8)} \\
 2 \cdot 5 \cdot x^2 + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13 & \text{PrCBPo(2.10a)} \\
 10 \cdot x^2 + 26 \cdot x + 16 \cdot x + 39 & \text{OOM(1.4)} \\
 10x^2 + 26x + 15x + 39 & \text{CTJ(1.7)} \\
 10x^2 + 41x + 39 & \text{OOA(1.1)}
 \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} 2x(5x + 13) + 3(5x + 13) & \text{DPE(2.9b)} \\ 10x^2 + 26x + 15x + 39 & \text{DPE(2.9a)} \\ 10x^2 + 41x + 39 & \text{OOA(1.1)} \end{array}$$

■

Example 2.37 — id:20141109-141019.Simplify by expanding $(-3x - 5)(7x + 8)$ **S****Solution:**

$$\begin{array}{ll} (-3x - 5)(7x + 8) & \text{ONeg(2.6a)} \\ (-3x + -5)(7x + 8) & \text{DOS(2.5a)} \\ -3x(7x + 8) + -5(7x + 8) & \text{DPE(2.9b)} \\ -3x \cdot 7x + -3x \cdot 8 + -5 \cdot 7x + -5 \cdot 8 & \text{DPE(2.9a)} \\ -3 \cdot x \cdot 7 \cdot x + -3 \cdot x \cdot 8 + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{JTC(1.6)} \\ -3 \cdot 7 \cdot x \cdot x + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{CPM(1.8)} \\ -3 \cdot 7 \cdot x^2 + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{PrCBPo(2.10a)} \\ -21 \cdot x^2 + -24 \cdot x + -35 \cdot x + -40 & \text{OOM(1.4)} \\ -21x^2 + -24x + -35x + -40 & \text{CTJ(1.7)} \\ -21x^2 + -59x + -40 & \text{OOA(1.1)} \\ -21x^2 - 59x - 40 & \text{DOS(2.5b)} \\ -21x^2 - 59x - 40 & \text{ONeg(2.6b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} (-3x + -5)(7x + 8) & \text{DOS(2.5a)} \\ -3x(7x + 8) + -5(7x + 8) & \text{DPE(2.9b)} \\ -21x^2 + -24x + -35x + -40 & \text{CTJ(2.9a)} \\ -21x^2 - 59x - 40 & \text{OOA(1.1)} \end{array}$$

■

Example 2.38 — id:20141109-141347.Simplify by expanding $(ax + b)(cx + d)$, where $a, b, c, d \in \mathbb{Z}$

S**Solution:**

$$\begin{aligned} ax(cx + d) + b(cx + d) && \text{DPE(2.9b)} \\ ax \cdot cx + ax \cdot d + b \cdot cx + b \cdot d && \text{DPE(2.9a)} \\ a \cdot x \cdot c \cdot x + a \cdot x \cdot d + b \cdot c \cdot x + b \cdot d && \text{JTC(1.6)} \\ a \cdot c \cdot x \cdot x + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d && \text{CPM(1.8)} \\ a \cdot c \cdot x^2 + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d && \text{PrCBPo(2.10a)} \\ acx^2 + adx + bcx + bd && \text{CTJ(1.7)} \\ acx^2 + (ad + bc)x + bd && \text{DPF(2.4a)} \end{aligned}$$

S**Less Steps Solution:**

$$\begin{aligned} ax(cx + d) + b(cx + d) && \text{DPE(2.9b)} \\ acx^2 + (ad + bc)x + bd && \text{DPE(2.9a)} \end{aligned}$$

■

Example 2.39 — id:20141105-161225.

Simplify $\left(2 - \frac{x}{2}\right)^2$ by expanding.

S

Solution:

$$\begin{aligned}
 & \left(2 - \frac{1x}{2}\right)^2 && \text{MId(2.1a)} \\
 & \left(2 + -\frac{1x}{2}\right)^2 && \text{DOS(2.5a)} \\
 & \left(2 + -\frac{1x}{2}\right) \left(2 + -\frac{1x}{2}\right) && \text{PoTF(1.13)} \\
 & 2 \left(2 + -\frac{1x}{2}\right) + -\frac{1x}{2} \left(2 + -\frac{1x}{2}\right) && \text{DPE(2.9b)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1x}{2} + -\frac{1x}{2} \cdot 2 + -\frac{1x}{2} \cdot -\frac{1x}{2} && \text{DPE(2.9a)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot x \cdot 2 + -\frac{1}{2} \cdot x \cdot -\frac{1}{2} \cdot x && \text{JTC(1.6)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x \cdot x && \text{CPM(1.8)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x^2 && \text{PrCBPo(2.10a)} \\
 & 4 + -1 \cdot x + -1 \cdot x + \frac{1}{4} \cdot x^2 && \text{OOM(1.4)} \\
 & 4 + -1x + -1x + \frac{1}{4}x^2 && \text{CTJ(1.7)} \\
 & \frac{1}{4}x^2 + -1x + -1x + 4 && \text{CPA(2.8)} \\
 & \frac{1}{4}x^2 + -2x + 4 && \text{OOA(1.1)} \\
 & \frac{1}{4}x^2 - 2x + 4 && \text{DOS(2.5b)}
 \end{aligned}$$

S

Less Steps Solution:

$$\begin{aligned}
 & 7x + 20x + 40 && \text{DPE(2.9a)} \\
 & 27x + 40 && \text{OOA(1.1)}
 \end{aligned}$$

■

2.6 Degree 3 Univariate Polynomials**Monomials****Binomials****Trinomials****Polynomials****2.7 Degree n Univariate Polynomials****Monomials****Binomials****Trinomials****Polynomials**

3. Multivariate Polynomial Expressions

Classification of Multivariate Polynomial Expressions

Degree -1 Multivariate Polynomials

Monomials

Degree 0 Multivariate Polynomials

Monomials

Degree 1 Multivariate Polynomials

Monomials

Binomials

Degree 2 Multivariate Polynomials

Monomials

Example 3.1 — id:20141106-150953.

Simplify $8xy + 5xy$

S

Solution:

$$\begin{aligned}(8 + 5)xy \\ 13xy\end{aligned}$$

DPF(2.4a)
OOA(1.1)

S

Less Steps Solution:

$$13xy \quad \text{OOA(1.1)}$$



Example 3.2 — id:20141108-193835.

Simplify $3xy + 4yx$

S

Solution:

$$\begin{array}{ll}
 3 \cdot x \cdot y + 4 \cdot y \cdot x & \text{JTC(1.6)} \\
 3 \cdot x \cdot y + 4 \cdot x \cdot y & \text{CPM(1.8)} \\
 3xy + 4xy & \text{CTJ(1.7)} \\
 (3 + 4)xy & \text{DPF(2.4a)} \\
 7xy & \text{OOA(1.1)}
 \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll}
 3xy + 4xy & \text{CPM(1.8)} \\
 7xy & \text{OOA(1.1)}
 \end{array}$$

**Example 3.3 — id:20141108-180100.**

Simplify $8x \cdot 2y$

S

Solution:

$$\begin{array}{ll}
 8 \cdot x \cdot 2 \cdot y & \text{JTC(1.6)} \\
 8 \cdot 2 \cdot x \cdot y & \text{CPM(1.8)} \\
 (8 \cdot 2) \cdot x \cdot y & \text{APM(1.10a)} \\
 16 \cdot x \cdot y & \text{OOM(1.4)} \\
 16xy & \text{CTJ(1.7)}
 \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll}
 8 \cdot 2 \cdot x \cdot y & \text{CPM(1.8)} \\
 16xy & \text{OOM(1.4)}
 \end{array}$$



Example 3.4 — id:20141108-180608.

Simplify $3x \cdot y$

S

Solution:

$$\begin{array}{ll}
 3x \cdot 1y & \text{MId(2.1a)} \\
 3 \cdot x \cdot 1 \cdot y & \text{JTC(1.6)} \\
 3 \cdot 1 \cdot x \cdot y & \text{CPM(1.8)} \\
 (3 \cdot 1) \cdot x \cdot y & \text{CPM(1.8)} \\
 3 \cdot x \cdot y & \text{OOM(1.4)} \\
 3xy & \text{CTJ(1.7)}
 \end{array}$$

S

Less Steps Solution:

$$3xy \quad \text{CTJ(1.7)}$$

**Binomials****Example 3.5 — id:20141106-151622.**

Simplify $5xy^2 - 3xy$

S

Solution:

$$\begin{array}{ll}
 5xy^2 + -3xy & \text{DOS(2.5a)} \\
 (5y + -3)xy & \text{DPF(2.4a)} \\
 (5y - 3)xy & \text{DOS(2.5b)}
 \end{array}$$

S

Less Steps Solution:

$$5xy^2 - 3xy$$

Depending on the context of the problem, both $(5y - 3)xy$, the factored form and $5xy^2 - 3xy$, the expanded form, can both be considered simplified.



Example 3.6 — id:20141108-180858.

Simplify $\frac{x}{4} \cdot \frac{y}{5}$

S**Solution:**

$$\begin{aligned} & \frac{1x}{4} \cdot \frac{1y}{5} && \text{MId(2.1a)} \\ & \frac{1}{4} \cdot x \cdot \frac{1}{5} \cdot y && \text{JTC(1.6)} \\ & \frac{1}{4} \cdot \frac{1}{5} \cdot x \cdot y && \text{CPM(1.8)} \\ & \frac{1}{20} \cdot x \cdot y && \text{OOM(1.4)} \\ & \frac{1xy}{20} && \text{CTJ(1.7)} \\ & \frac{xy}{20} && \text{MId(2.1b)} \end{aligned}$$

S**Less Steps Solution:**

$$\frac{xy}{20} \quad \text{OOM(1.4)}$$

**Trinomials****Degree 3 Multivariate Polynomials****Monomials****Example 3.7 — id:20141106-151409.**

Simplify $7xy^2 - xy^2$

S**Solution:**

$$\begin{aligned} & 7xy^2 - 1xy^2 && \text{MId(2.1a)} \\ & 7xy^2 + -1xy^2 && \text{DOS(2.5a)} \\ & (7 + -1)xy^2 && \text{DPF(2.4a)} \\ & 6xy^2 && \text{OOA(1.1)} \end{aligned}$$

S

Less Steps Solution:

$$7xy^2 + -xy^2$$

$$6xy^2$$

DOS(2.5a)

OOA(1.1)

Binomials**Trinomials****Polynomials****Degree n Multivariate Polynomials****Monomials****Rule 3.0.1 — Power of a Product (PoPr).**

$$(a \cdot b)^k = a^k \cdot b^k \quad (3.1a)$$

$$a^k \cdot b^k = (a \cdot b)^k \quad (3.1b)$$

Example 3.8 — id:20141108-193145.Simplify $12xy \cdot (-\frac{1}{2})yx$

S

Solution:

$$12x1y \cdot \left(-\frac{1}{2}\right) y1x \quad \text{MId}(2.1a)$$

$$12x1y \cdot -\frac{1}{2}y1x \quad \text{ONeg}(2.6a)$$

$$12 \cdot x \cdot 1 \cdot y \cdot -\frac{1}{2} \cdot y \cdot 1 \cdot x \quad \text{JTC}(1.6)$$

$$12 \cdot 1 \cdot -\frac{1}{2} \cdot 1 \cdot x \cdot x \cdot y \cdot y \quad \text{CPM}(1.8)$$

$$12 \cdot 1 \cdot -\frac{1}{2} \cdot 1 \cdot x^2 \cdot y^2 \quad \text{PrCBPo}(2.10a)$$

$$\left(12 \cdot 1 \cdot -\frac{1}{2} \cdot 1\right) \cdot x^2 \cdot y^2 \quad \text{APM}(1.10a)$$

$$-6 \cdot x^2 \cdot y^2 \quad \text{OOM}(1.4)$$

$$-6x^2y^2 \quad \text{CTJ}(1.7)$$

$$-6x^2y^2 \quad \text{ONeg}(2.6b)$$

S

Less Steps Solution:

$$12 \cdot -\frac{1}{2} \cdot x \cdot x \cdot y \cdot y \quad \text{CPM(1.8)}$$

$$12 \cdot -\frac{1}{2} \cdot x^2 \cdot y^2 \quad \text{PrCBPo(2.10a)}$$

$$-6x^2y^2 \quad \text{OOM(1.4)}$$

■

Example 3.9 — id:20141108-192234.Simplify $ac \cdot ab$

S

Solution:

$$\begin{aligned} & 1a1c \cdot 1a1b && \text{MId(2.1a)} \\ & 1 \cdot a \cdot 1 \cdot c \cdot 1 \cdot a \cdot 1 \cdot b && \text{JTC(1.6)} \\ & 1 \cdot 1 \cdot 1 \cdot 1 \cdot a \cdot a \cdot b \cdot c && \text{CPM(1.8)} \\ & 1 \cdot 1 \cdot 1 \cdot 1 \cdot a^2 \cdot b \cdot c && \text{PrCBPo(2.10a)} \\ & (1 \cdot 1 \cdot 1 \cdot 1) \cdot a^2 \cdot b \cdot c && \text{APM(1.10a)} \\ & 1 \cdot a^2 \cdot b \cdot c && \text{OOM(1.4)} \\ & 1a^2bc && \text{CTJ(1.7)} \\ & a^2bc && \text{MId(2.1b)} \end{aligned}$$

S

Less Steps Solution:

$$\begin{aligned} & a \cdot a \cdot b \cdot c && \text{CPM(1.8)} \\ & a^2bc && \text{PrCBPo(2.10a)} \end{aligned}$$

■

Example 3.10 — id:20141108-192611.Simplify $(-ac) \cdot (-bc)$

S

Solution:

$$\begin{aligned}
 & (-1a1c) \cdot (-1b1c) && \text{MId(2.1a)} \\
 & \neg 1a1c \cdot \neg 1b1c && \text{ONeg(2.6a)} \\
 & \neg 1 \cdot a \cdot 1 \cdot c \cdot \neg 1 \cdot b \cdot 1 \cdot c && \text{JTC(1.6)} \\
 & \neg 1 \cdot \neg 1 \cdot 1 \cdot 1 \cdot a \cdot b \cdot c \cdot c && \text{CPM(1.8)} \\
 & \neg 1 \cdot \neg 1 \cdot 1 \cdot 1 \cdot a \cdot b \cdot c^2 && \text{PrCBPo(2.10a)} \\
 & (\neg 1 \cdot \neg 1 \cdot 1 \cdot 1) \cdot a \cdot b \cdot c^2 && \text{APM(1.10a)} \\
 & 1 \cdot a \cdot b \cdot c^2 && \text{OOM(1.4)} \\
 & 1abc^2 && \text{CTJ(1.7)} \\
 & abc^2 && \text{MId(2.1b)}
 \end{aligned}$$

S**Less Steps Solution:**

$$\begin{aligned}
 & \neg a \cdot \neg b \cdot c \cdot c && \text{CPM(1.8)} \\
 & abc^2 && \text{PrCBPo(2.10a)}
 \end{aligned}$$

Binomials**Trinomials****Polynomials**

4. Equations

4.1 Equality

Property 4.1.1 — Reflexive Property of Eqaulity (RPE).

$$a = a \quad (4.1a)$$

Property 4.1.2 — Substitution Property of Equality (SPE).

Given $a = b$, then

$$E(a) = E(b) \quad (4.2)$$

$E(x)$ represents any expression.

Property 4.1.3 — Symmetric Property of Equality (SyPE).

$$a = b \quad \text{then} \quad b = a \quad (4.3a)$$

Property 4.1.4 — Transitive Property of Equality (TPE).

$$\text{if } a = b \quad \text{and} \quad b = c \quad \text{then} \quad a = c \quad (4.4a)$$

4.2 Solving Linear Equations

Example 4.1 — id:20141206-102142.

Solve the equation $x + a = b$ for x

S**Solution:**

$$\begin{array}{ll}
 [x + a] + \neg a = [b] + \neg a & \text{SPE(4.2) + AI(1.19a)} \\
 x + (a + \neg a) = b + \neg a & \text{APA(2.7b)} \\
 x + 0 = b + \neg a & \text{OOA(1.1)} \\
 x = b + \neg a & \text{AId(2.3b)} \\
 x = b - a & \text{DOS(2.5b)}
 \end{array}$$

■

Example 4.2 — id:20141111-222931.Solve the equations $x + 8 = 0$ **S****Solution:**

$$\begin{array}{ll}
 [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.2) + AI(1.19a)} \\
 x + (8 + \neg 8) = 0 + \neg 8 & \text{APA(2.7b)} \\
 x + 0 = \neg 8 & \text{OOA(1.1)} \\
 x = \neg 8 & \text{AId(2.3b)} \\
 x = -8 & \text{ONeg(2.6b)}
 \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll}
 [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.2) + AI(1.19a)} \\
 x = -8 & \text{OOA(1.1)}
 \end{array}$$

D**Dependencies:**

example 5.2-20141111-190212

■

Example 4.3 — id:20141206-101632.Solve the equation $x + 4 = 7$ **S**

Solution:

$$\begin{array}{ll} [x] + 4 + \neg 4 = [7] + \neg 4 & \text{SPE(4.2) + AI(1.19a)} \\ x + (4 + \neg 4) = 7 + \neg 4 & \text{APA(2.7b)} \\ x + 0 = 3 & \text{OOA(1.1)} \\ x = 3 & \text{AIId(2.3b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} [x] + 4 + \neg 4 = [7] + \neg 4 & \text{SPE(4.2) + AI(1.19a)} \\ x = 3 & \text{OOA(1.1)} \end{array}$$

■

Example 4.4 — id:20141206-101107.Solve the equation $x - 8 = 15$ for x **S****Solution:**

$$\begin{array}{ll} x + \neg 8 = 15 & \text{DOS(2.5a)} \\ [x + \neg 8] + 8 = [15] + 8 & \text{SPE(4.2) + AI(1.19a)} \\ x + (-8 + 8) = 15 + 8 & \text{APA(2.7b)} \\ x + 0 = 23 & \text{OOA(1.1)} \\ x = 23 & \text{AIId(2.3b)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} [x + \neg 8] + 8 = [15] + 8 & \text{SPE(4.2) + AI(1.19a)} \\ x = 23 & \text{OOA(1.1)} \end{array}$$

■

Example 4.5 — id:20141206-102404.Solve the equation $5x = 9$ for x .

S**Solution:**

$$\begin{aligned}\frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.2) + MI(1.9a)} \\ \frac{1}{5} \cdot [5 \cdot x] &= \frac{1}{5} \cdot 9 && \text{JTC(1.6)} \\ \left(\frac{1}{5} \cdot 5\right) \cdot x &= \frac{1}{5} \cdot 9 && \text{APM(1.10b)} \\ 1 \cdot x &= \frac{9}{5} && \text{OOM(1.4)} \\ x &= \frac{9}{5} && \text{MId(2.1b)}\end{aligned}$$

S**Less Steps Solution:**

$$\begin{aligned}\frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.2) + MI(1.9a)} \\ x &= \frac{9}{5} && \text{OOM(1.4)}\end{aligned}$$

**Example 4.6 — id:20141206-104404.**Solve the equation $ax = b$ for x .**S****Solution:**

$$\begin{aligned}\frac{1}{a}[ax] &= \frac{1}{a}[b] && \text{SPE(4.2) + MI(1.9a)} \\ \frac{1}{a} \cdot (a \cdot x) &= \frac{1}{a} \cdot b && \text{JTC(1.6)} \\ \left(\frac{1}{a} \cdot a\right) \cdot x &= \frac{1}{a} \cdot b && \text{APM(1.10b)} \\ 1 \cdot x &= \frac{b}{a} && \text{OOM(1.4)} \\ x &= \frac{b}{a} && \text{MId(2.1b)}\end{aligned}$$

**Example 4.7 — id:20141206-102723.**

Solve the equation $-2x = 7$ for x

S

Solution:

$$\begin{aligned}-2x &= 7 && \text{ONeg(2.6a)} \\ -\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.2) + MI(1.9b)} \\ -\frac{1}{2} \cdot (-2 \cdot x) &= -\frac{1}{2} \cdot 7 && \text{JTC(1.6)} \\ \left(-\frac{1}{2} \cdot -2\right) \cdot x &= -\frac{1}{2} \cdot 7 && \text{APM(1.10b)} \\ 1 \cdot x &= -\frac{7}{2} && \text{OOM(1.4)} \\ 1 \cdot x &= -\frac{7}{2} && \text{ONeg(2.6b)} \\ x &= -\frac{7}{2} && \text{MId(2.1b)}\end{aligned}$$

S

Less Steps Solution:

$$\begin{aligned}-\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.2) + MI(1.9b)} \\ x &= -\frac{7}{2} && \text{OOM(1.4)}\end{aligned}$$

■

Example 4.8 — id:20141111-215726.

Solve the equation $2x + 5 = 0$ for x

S

Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.2) + AI(1.19a)} \\
 2x + (5 + \neg 5) &= 0 + \neg 5 && \text{APA(2.7a)} \\
 2x + 0 &= \neg 5 && \text{OOA(1.1)} \\
 2x &= \neg 5 && \text{AId(2.3a)} \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.2) + MI(1.9a)} \\
 \frac{1}{2} \cdot 2 \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{JTC(1.6)} \\
 \left(\frac{1}{2} \cdot 2\right) \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{APM(1.10a)} \\
 1 \cdot x &= \frac{\neg 5}{2} && \text{OOM(1.4)} \\
 1x &= -\frac{5}{2} && \text{ONeg(2.6b)} \\
 x &= -\frac{5}{2} && \text{MId(2.1b)}
 \end{aligned}$$

S

Less Steps Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.2) + AI(1.19a)} \quad (4.5) \\
 2x &= \neg 5 && \text{OOA(1.1)} \quad (4.6) \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.2) + MI(1.9a)} \quad (4.7) \\
 x &= -\frac{5}{2} && \text{OOM(1.4)} \quad (4.8)
 \end{aligned}$$

D

Dependencies:

example 5.3-20141111-192213

4.3 Solving Quadratic Equations

Example 4.9 — id:20141107-131748.

Solve the equation $2 - x^2 = 0$ for x

S

Solution:

$$\begin{aligned}
 2 - 1x^2 &= 0 && \text{MId(2.1a)} \\
 2 + -1x^2 &= 0 && \text{DOS(2.5a)} \\
 [2 + -1x^2] + 1x^2 &= [0] + 1x^2 && \text{SPE(4.2) + AI(1.19a)} \\
 2 + (-1x^2 + 1x^2) &= 0 + 1x^2 && \text{APA(2.7a)} \\
 2 + 0 &= 0 + 1x^2 && \text{OOA(1.1)} \\
 2 &= 1x^2 && \text{AId(2.3a)} \\
 2 &= x^2 && \text{MId(2.1b)} \\
 \pm [2]^{\frac{1}{2}} &= [x^2]^{\frac{1}{2}} \\
 \pm 2^{\frac{1}{2}} &= x && \text{PoPo(2.12a)} \\
 \pm \sqrt{2} &= x && \text{PoTR(1.18)} \\
 x = \pm \sqrt{2} & && \text{SyPE(4.3a)}
 \end{aligned}$$

■

4.3.1 Completing the Square

Completing the square is an algebraic algorithm used to find the solutions of quadratic equations of the form, $ax^2 + bx + c = 0$. Essentially, we want to manipulate this equation such that $x = \text{some value}$. To gain some understanding of how this algorithm works, we will consider each step individually. Let's begin with a quadratic equation in the general form: $ax^2 + bx + c = 0$.

Since we are trying to manipulate the equation $ax^2 + bx + c = 0$ such that $x = \text{some value}$, we first want the coefficient factor a to be equal to 1. This is done by multiplying both expressions by the multiplicative inverse of a (step 1) followed by simplifying both expressions.

$$\begin{aligned}
 \frac{1}{a}[ax^2 + bx + c] &= \frac{1}{a}[0] && \text{SPE(4.2) + MI(1.9a)} \\
 \frac{1}{a} \cdot ax^2 + \frac{1}{a} \cdot bx + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{DPE(2.9a)} \\
 \frac{1}{a} \cdot a \cdot x^2 + \frac{1}{a} \cdot b \cdot x + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{JTC(1.6)} \\
 x^2 + \frac{b}{a} \cdot x + \frac{c}{a} &= 0 && \text{OOM(1.4)} \\
 x^2 + \frac{b}{a}x + \frac{c}{a} & && \text{CTJ(1.7)}
 \end{aligned}$$

We now have three terms in the left hand expression where the first two terms have at least one variable factor, x . The problem is that the third term is a constant and we want $x = \text{some value}$. This text step is to get rid of the $\frac{c}{a}$ term, which can be done by using the additive inverse followed by simplifying both expressions.

$$\begin{aligned} \left[x^2 + \frac{b}{a}x + \frac{c}{a} \right] + -\frac{c}{a} &= [0] + -\frac{c}{a} && \text{SPE(4.2) + AI(1.19a)} \\ x^2 + \frac{b}{a}x + \frac{c}{a} + -\frac{c}{a} &= 0 + -\frac{c}{a} && \text{APA(2.7a)} \\ x^2 + \frac{b}{a}x &= -\frac{c}{a} && \text{OOA(1.1)} \end{aligned}$$

The next step is called completing the square. The idea is to add a *NeW* constant, k , to the left-hand expression, $x^2 + \frac{b}{a}x + k$, such that the quadratic expression can then be factored as two identical factors, $(x + m)(x + m) = (x + m)^2$, where $k = m \cdot m$. Notice that since we are adding a constant term, k , to the left-hand expression, then we must also add this constant, k , to the right-hand expression, $x^2 + \frac{b}{a}x + k = -\frac{c}{a} + k$. To determine the values of both m and k we should refer to the organization of the two factors that make up the product of a quadratic expression, $x^2 + \frac{b}{a}x + k = (x + m)^2$.

Since both factors of this new quadratic expression are the same, both terms that make up the middle term, must also be the same. We know that $mx + mx = \frac{b}{a}x$, so we should be able to determine the value of m from this equation. If we can determine the value of m , then we can determine the value of k .

$$\begin{aligned} \frac{b}{a}x &= mx + mx \\ &= 2mx && \text{OOA(1.1)} \end{aligned}$$

Solving for m ,

$$\begin{aligned} 2mx &= \frac{b}{a}x \\ 2 \cdot m \cdot x &= \frac{b}{a} \cdot x && \text{JTC(1.6)} \\ \frac{1}{2} [2 \cdot m \cdot x] &= \frac{1}{2} \left[\frac{b}{a} \cdot x \right] && \text{SPE(4.2) + MI(1.9a)} \\ m \cdot x &= \frac{b}{2a} \cdot x && \text{OOM(1.4)} \\ [m \cdot x] \frac{1}{x} &= \left[\frac{b}{2a} \cdot x \right] \frac{1}{x} && \text{SPE(4.2) + MI(1.9a)} \\ m &= \frac{b}{2a} && \text{OOM(1.4)} \end{aligned}$$

$$\begin{aligned}
 \left[x^2 + \frac{b}{a}x \right] + \left(\frac{b}{2a} \right)^2 &= \left[-\frac{c}{a} \right] + \left(\frac{b}{2a} \right)^2 && \text{SPE(4.2)} \\
 x^2 + \frac{b}{a}x + \left(\frac{b}{2a} \right)^2 &= -\frac{c}{a} + \left(\frac{b}{2a} \right)^2 && \text{APA(2.7b)} \\
 \left(x + \frac{b}{2a} \right) \left(x + \frac{b}{2a} \right) &= -\frac{c}{a} + \left(\frac{b}{2a} \right)^2 && \text{DPF(2.4b)} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c}{a} + \left(\frac{b}{2a} \right)^2 && \text{PoTF(1.13)} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c}{a} + \frac{(b)^2}{(2a)^2} && \text{PoQPo(1.11a)} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c}{a} + \frac{b^2}{2^2 a^2} && \text{PoPrPo} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c}{a} + \frac{b^2}{4a^2} && \text{OOE} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c}{a} \cdot \frac{\cancel{4a}}{\cancel{4a}} + \frac{b^2}{4a^2} && \text{MId} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{c \cdot 4 \cdot a}{a \cdot 4 \cdot a} + \frac{b^2}{4a^2} && \text{JTC} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{4 \cdot a \cdot c}{4 \cdot a \cdot a} + \frac{b^2}{4a^2} && \text{CPM} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{4 \cdot a \cdot c}{4 \cdot a^2} + \frac{b^2}{4a^2} && \text{PrCBPo} \\
 \left(x + \frac{b}{2a} \right)^2 &= -\frac{4ac}{4a^2} + \frac{b^2}{4a^2} && \text{CTJ} \\
 \left(x + \frac{b}{2a} \right)^2 &= \frac{-4ac + b^2}{4a^2} && \text{CD} \\
 \left(x + \frac{b}{2a} \right)^2 &= \frac{b^2 - -4ac}{4a^2} && \text{CPA} \\
 \left[\left(x + \frac{b}{2a} \right)^2 \right]^{\frac{1}{2}} &= \pm \left[\frac{b^2 - -4ac}{4a^2} \right]^{\frac{1}{2}} && \text{PoI}
 \end{aligned}$$

$$x + \frac{b}{2a} = \pm \left[\frac{b^2 + \neg 4ac}{4a^2} \right]^{\frac{1}{2}} \quad \text{PoPrPo}$$

$$x + \frac{b}{2a} = \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{[4a^2]^{\frac{1}{2}}} \quad \text{PoQPo}$$

$$x + \frac{b}{2a} = \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{4^{\frac{1}{2}}a} \quad \text{PoPrPo}$$

$$x + \frac{b}{2a} = \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} \quad \text{OOE}$$

$$\left[x + \frac{b}{2a} \right] + \neg \frac{b}{2a} = \left[\pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} \right] + \neg \frac{b}{2a} \quad \text{AI}$$

$$x + \frac{b}{2a} + \neg \frac{b}{2a} = \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} + \neg \frac{b}{2a} \quad \text{APA}$$

$$x + \frac{b}{2a} + \neg \frac{b}{2a} = \neg \frac{b}{2a} \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} \quad \text{CPA}$$

$$x = \neg \frac{b}{2a} \pm \frac{[b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} \quad \text{OOA}$$

$$x = \frac{\neg b \pm [b^2 + \neg 4ac]^{\frac{1}{2}}}{2a} \quad \text{CD}$$

$$x = \frac{\neg b \pm [b^2 - 4ac]^{\frac{1}{2}}}{2a} \quad \text{DOS}$$

$$x = \frac{\neg b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{ETR}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{ONeg}$$



5. Functions

5.1 Evaluating Functions

Example 5.1 — id:20141106-083703.

Given $f(x) = \frac{1}{12}x^3 - x^2 + 4x$, evaluate $f(3) - f(2)$.

S

Solution:

$$\begin{aligned}f(3) - f(2) &= \left(\frac{[3]^3}{12} - [3]^2 + 4[3] \right) - \left(\frac{[2]^3}{12} - [2]^2 + 4[2] \right) \\&= \left(\frac{(3)^3}{12} - (3)^2 + 4(3) \right) - 1 \left(\frac{(2)^3}{12} - (2)^2 + 4(2) \right) && \text{MId(2.1a)} \\&= \left(\frac{(3)^3}{12} + -(3)^2 + 4(3) \right) + -1 \left(\frac{(2)^3}{12} + -(2)^2 + 4(2) \right) && \text{DOS(2.5a)} \\&= \left(\frac{27}{12} + -9 + 4(3) \right) + -1 \left(\frac{8}{12} + -4 + 4(2) \right) && \text{OOE(1.5)} \\&= \left(\frac{27}{12} + -9 + 12 \right) + -1 \left(\frac{8}{12} + -4 + 8 \right) && \text{OOM(1.4)} \\&= \left(\frac{27}{12} + 3 \right) + -1 \left(\frac{8}{12} + 4 \right) && \text{OOA(1.1)} \\&= \left(\frac{27+36}{12} \right) + -1 \left(\frac{8+48}{12} \right) && \text{FOOA(1.3a)} \\&= \left(\frac{63}{12} \right) + -1 \left(\frac{56}{12} \right) && \text{OOA(1.1)} \\&= \left(\frac{63}{12} \right) + -\frac{56}{12} && \text{OOM(1.4)} \\&= \frac{7}{12} && \text{OOA(1.1)}\end{aligned}$$

D

Dependencies:
example 8.2-20141105-144223

5.2 Quadratic Functions

5.2.1 Completing the Square

Definition 5.2.1 — Completing The Square. Completing the square is the process used to convert a quadratic polynomial function $f(x) = ax^2 + bx + c$ to the form

$$f(x) = a \left(x + \frac{b}{2a} \right)^2 + \left(c - \frac{b^2}{4a} \right)$$

We can simplify this form by defining $B = \frac{b}{2a}$ and $C = c - \frac{b^2}{4a}$, which gives us

$$f(x) = a(x + B) + C$$

[Weib]

$$\begin{aligned} f(x) &= ax^2 + bx + c \\ &= a \left[x^2 + \frac{b}{a}x + \frac{c}{a} \right] && \text{DPF(2.4b)} \\ &= a \left[x^2 + \frac{b}{a}x + k + \neg k + \frac{c}{a} \right] && \text{AId(2.3b)} \\ &= a \left[\left(x^2 + \frac{b}{a}x + k \right) + \left(-k + \frac{c}{a} \right) \right] && \text{APA(2.7b)} \end{aligned}$$




Figure 5.1: Factoring Organizer used to find the value of k

$$\begin{aligned}
 f(x) &= a \left[\left(x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} \right) + \left(\frac{-b^2}{4a^2} + \frac{c}{a} \right) \right] \\
 &= a \left[\left(x + \frac{b}{2a} \right)^2 + \frac{-b^2}{4a^2} + \frac{c}{a} \right] && \text{DPF(2.4b)} \\
 &= a \left[\left(x + \frac{b}{2a} \right)^2 + \frac{-b^2 + 4ac}{4a^2} \right] \\
 &= a \left[\left(x + \frac{b}{2a} \right)^2 + \frac{4ac - b^2}{4a^2} \right] && \text{OOA(1.1)} \\
 &= a \left(x + \frac{b}{2a} \right)^2 + \frac{4ac - b^2}{4a} && \text{DPE(2.9a)} \\
 &= a \left(x + \frac{b}{2a} \right)^2 + \left(c - \frac{b^2}{4a} \right)
 \end{aligned}$$

5.3 Rational Functions

Definition 5.3.1 — Asymptote. An asymptote is a line or curve that approaches a given curve arbitrarily close.[Weia]

A vertical asymptote is a vertical line $x_{va} = c$, that is used to visualize the values of x for which the function is not defined.




Figure 5.2: Vertical Asymptote

Example 5.2 — id:20141111-190212.

Find the vertical asymptote of the function $R(x) = \frac{7}{x+8}$

S


Solution:

We are interested in the values of x for which the denominator of $R(x)$ has a value of zero.

$$x_{va} + 8 = 0$$

$$x_{va} = -8$$

solving

**Example 5.3 — id:20141111-192213.**

Find the vertical asymptote of the function $R(x) = \frac{x+4}{2x+5}$

S


Solution:

We are interested in the values of x for which the denominator of $R(x)$ has a value of zero.

$$2x_{va} + 5 = 0$$

$$x_{va} = -\frac{5}{2}$$

solving 4.8



6. Vectors

6.1 Component & Geometric Vectors

6.2 Dot Product

Definition 6.2.1 — Dot Product in \mathbb{R}^2 (VDPr). The dot product of vectors $\vec{u} = (u_x, u_y)$ and $\vec{v} = (v_x, v_y)$ is

$$\vec{u} \cdot \vec{v} = u_x v_x + u_y v_y \quad (6.1)$$

The dot product of vectors $\vec{u} = u_x \hat{i} + u_y \hat{j}$ and $\vec{v} = v_x \hat{i} + v_y \hat{j}$ is

$$\vec{u} \cdot \vec{v} = u_x v_x \hat{i} + u_y v_y \hat{j} \quad (6.2)$$

Example 6.1 — id:20141214-195907.

Find the dot product of the vectors $\vec{u} = (-5, 12)$ and $\vec{v} = (-7, -2)$

S

Solution:

$$\begin{aligned}\vec{u} \cdot \vec{v} &= u_x v_x + u_y v_y && \text{VDPr(6.1)} \\ &= [-5][-7] + [12][-2] && \text{SPE(4.2)} \\ &= 35 + -24 && \text{OOM(1.4)} \\ &= 11\end{aligned}$$

Example 6.2 — id:20141214-194701.

Find the dot product of the vectors $\vec{u} = -10\hat{i} - 10\hat{j}$ and $\vec{v} = -9\hat{i} + 7\hat{j}$

S

$$\begin{aligned}\vec{u} \cdot \vec{v} &= u_x v_x \hat{i} + u_y v_y \hat{j} && \text{VDPr(6.2)} \\ &= [-10][-9] + [-10][7] && \text{SPE(4.2)} \\ &= 90 + -70 && \text{OOM(1.4)} \\ &= 20 && \text{OOA(1.1)}\end{aligned}$$

■

7. Differentiation

7.1 Limit of the Difference Quotient

Definition 7.1.1 — Derivative. The derivative of a function $f(x)$ with respect to the variable x is defined as

$$f'(x) \equiv \lim_{\Delta x \rightarrow 0} \underbrace{\frac{f(x + \Delta x) - f(x)}{\Delta x}}_{\text{Difference Quotient}} \quad (7.1)$$




Figure 7.1: [mooculus:textbook]

Example 7.1 — id:20141219-212546.

Differentiate the function $f(x) = 5$

S**Solution:**

$$f(x) = 5x^0 \quad \text{PoID(1.16a)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5[x + \Delta x]^0 - 5[x]^0}{\Delta x} \quad \text{SPE(4.2)&DBFP(7.1)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5(1) - 5(1)}{\Delta x} \quad \text{PoID(1.16b)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} 0 \quad \text{OOM(1.4)}$$

$$f'(x) = 0$$

■

7.2 Derivative of a Monomial Functions

Rule 7.2.1 — Derivative of a Constant (DC).

$$[c]' = 0 \quad (7.2)$$

$$\frac{d}{dx} [c] = 0 \quad (7.3)$$

Rule 7.2.2 — Derivative of a Constant Multiple (DCM).

$$[cf(x)]' = c[f(x)]' \quad (7.4)$$

$$\frac{d}{dx} [cf(x)] = c \frac{d}{dx} [f(x)] \quad (7.5)$$

Rule 7.2.3 — Derivative of a Power (DPO).

$$[x^n]' = nx^{n-1} \quad (7.6)$$

$$\frac{d}{dx} [x^n] = nx^{n-1} \quad (7.7)$$

Example 7.2 — id:20141124-153017.

Differentiate $f(x) = -3$

S**Solution:**

$$f'(x) = [-3]' \quad \text{SPE(4.2)}$$

$$f'(x) = 0 \quad \text{DC(7.2)}$$

D

Dependencies:

example 7.4-20141124-152503

Example 7.3 — id:20141124-141850.Differentiate $f(x) = x^2$ **S****Solution:**

$$f'(x) = [x^2]' \quad \text{SPE(4.2)}$$

$$f'(x) = 2x^{2-1} \quad \text{DPo(7.6)}$$

$$f'(x) = 2x^1 \quad \text{OOA(1.1)}$$

$$f'(x) = 2x \quad \text{MId(2.1b)}$$

S**Less Steps Solution:**

$$f'(x) = 2x \quad \text{DPo(7.6)}$$

D**Dependencies:**

example 7.4-20141124-152503

7.3 Derivative of Polynomial Functions**Rule 7.3.1 — Derivative of a Sum (DS).**

$$[f(x) + g(x)]' = f'(x) + g'(x) \quad (7.8)$$

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} [f(x)] + \frac{d}{dx} [g(x)] \quad (7.9)$$

Example 7.4 — id:20141124-152503.Differentiate $f(x) = x^2 - 3$ **S**

Solution:

$$\begin{aligned}
 f(x) &= x^2 + -3 && \text{DOS(2.5a)} \\
 f'(x) &= [x^2 + -3]' && \text{SPE(4.2)} \\
 f'(x) &= [x^2]' + [-3]' && \text{DS(7.8)} \\
 f'(x) &= [x^2]' + 0 && \text{DC(7.2)} \\
 f'(x) &= [x^2]' && \text{AId(2.3a)} \\
 f'(x) &= 2x && \text{DPo(7.6) goto 7.3}
 \end{aligned}$$

S

Less Steps Solution:

$$f(x) = 2x^2 \quad \text{DPo(7.6)&DC(7.6)}$$

D

Dependencies:

example 7.12-20141124-205219



Example 7.5 — id:20141128-151834.

Differentiate $f(x) = 3x^2 - 6x + 4$

S

Solution:

$$\begin{aligned}
 f(x) &= 3x^2 - 6x + 4 && \text{DOS(2.5a)} \\
 f'(x) &= [3x^2 - 6x + 4]' && \text{SPE(4.2)} \\
 f'(x) &= [3x^2]' + [-6x]' + [4]' && \text{DS(7.8)} \\
 f'(x) &= [3x^2]' + [6x]' + 0 && \text{DC(7.2)} \\
 f'(x) &= [3x^2]' + [6x]' && \text{AId(2.3a)} \\
 f'(x) &= 3[x^2]' + 6[x]' && \text{DCM(7.4)} \\
 f'(x) &= 3(2x) + 6(1) && \text{DPo(7.6)} \\
 f'(x) &= 6x + 6 && \text{OOM(1.4)}
 \end{aligned}$$

S

$$f'(x) = 6x + 6$$

DS(7.8)

■

Rule 7.3.2 — Derivative of a Product (DPr).

$$[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x) \quad (7.10)$$

$$\frac{d}{dx} [f(x)g(x)] = \frac{d}{dx} [f(x)]g(x) + f(x) \frac{d}{dx} [g(x)] \quad (7.11)$$

Example 7.6 — id:20141209-144203.Differentiate $f(x) = x^2(2x + 4)$

S

Solution:

$$f'(x) = [x^2(2x + 4)]' \quad \text{SPE(4.2)}$$

$$f'(x) = [x^2]'(2x + 4) + x^2[2x + 4]' \quad \text{DPr(7.10)}$$

$$f'(x) = [x^2]'(2x + 4) + x^2[2x] + [4]' \quad \text{DS(7.8)}$$

$$f'(x) = [x^2]'(2x + 4) + x^2 \cdot 2[x] + [4]' \quad \text{DCM(7.4)}$$

$$f'(x) = 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + [4]' \quad \text{DPo(7.6)}$$

$$f'(x) = 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0 \quad \text{DC(7.2)}$$

$$f'(x) = 6x^2 + 8x \quad \text{simplify goto 2.33}$$

■

Example 7.7 — id:20141209-142321.Differentiate $f(x) = x^2 \cos(x)$

S

Solution:

$$f'(x) = [x^2 \cos(x)]' \quad \text{SPE(4.2)}$$

$$f'(x) = [x^2]' \cos(x) + x^2[\cos(x)]' \quad \text{DPr(7.10)}$$

$$f'(x) = 2x \cos(x) + x^2[\cos(x)]' \quad \text{DPo(7.6)}$$

$$f'(x) = 2x \cos(x) + x^2(-1 \sin(x)) \quad \text{DCos(7.16)}$$

$$f'(x) = 2x \cos(x) - x^2 \sin x \quad \text{OOM(1.4)}$$

Example 7.8 — id:20141209-151354.

Differentiate $f(x) = \sin(x) \sin(x)$

S**Solution:**

$$\begin{aligned} f'(x) &= [\sin(x) \sin(x)] && \text{SPE(4.2)} \\ f'(x) &= [\sin(x)]' \sin(x) + \sin(x) [\sin(x)] && \text{DPr(7.10)} \\ f'(x) &= \cos(x) \sin(x) + \sin(x) \cos(x) && \text{DSin(7.14)} \\ f'(x) &= \cos(x) \sin(x) + \cos(x) \sin(x) && \text{CPM(1.8)} \\ f'(x) &= 2 \cos(x) \sin(x) && \text{OOA(1.1)} \end{aligned}$$

7.4 Derivative of Rational Functions**Rule 7.4.1 — Derivative of a Quotient (DQ).**

$$\left[\frac{f(x)}{g(x)} \right]' = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2} \quad (7.12)$$

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{\frac{d}{dx}[f(x)]g(x) - f(x)\frac{d}{dx}[g(x)]}{[g(x)]^2} \quad (7.13)$$

7.5 Derivative of Trigonometric Functions**Rule 7.5.1 — Derivative of Sine (DSin).**

$$[\sin(x)]' = \cos(x) \quad (7.14)$$

$$\frac{d}{dx} [\sin(x)] = \cos(x) \quad (7.15)$$

Rule 7.5.2 — Derivative of Cosine (DCos).

$$[\cos(x)]' = -\sin(x) \quad (7.16)$$

$$\frac{d}{dx} [\cos(x)] = -\sin(x) \quad (7.17)$$

7.6 Derivative of Exponential Functions**7.7 Derivative of Logarithmic Functions**

Rule 7.7.1 — Derivative of a Natural Logarithm (DNL).

$$[\ln x]' = \frac{1}{x} \quad (7.18)$$

$$\frac{d}{dx} [\ln x] = \frac{1}{x} \quad (7.19)$$

7.8 Derivative of Composite Functions

Rule 7.8.1 — Derivative of a Composite Function (DComp).

$$[f(g(x))]' = [g(x)]' [f(g(x))]' \quad (7.20)$$

$$\frac{d}{dx} [f(g(x))] = \frac{d}{dx} [g(x)] \frac{d}{dx} [f(g(x))] \quad (7.21)$$

Example 7.9 — id:20141124-203850.

Differentiate $y = \ln(3x)$

S

Solution:

After identifying that $y = \ln(3x)$ is a composite function, we let $u = 3x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp}(7.21)$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = 3x$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = 3$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} && \text{DComp(7.21)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot \frac{du}{dx} && \text{DNL(7.18)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot 3 && \text{DPo(7.6)} \\ \frac{dy}{dx} &= \frac{1}{3x} \cdot 3 && \\ \frac{dy}{dx} &= \frac{3}{3x} && \text{OOM(1.4)} \\ \frac{dy}{dx} &= \frac{1}{x}\end{aligned}$$

D**Dependencies:**

example 7.12-20141124-205219

**Example 7.10 — id:20141128-160248.**Differentiate $y = \ln(3x^2 - 6x + 4)$ **S****Solution:** After identifying that $y = \ln(3x^2 - 6x + 4)$ is a composite function, we let $u = 3x^2 - 6x + 4$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.21)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$\begin{aligned}y &= \ln(u) & u &= 3x^2 - 6x + 4 \\ \frac{dy}{du} &= \frac{1}{u} & \frac{du}{dx} &= 6x - 6 \quad \text{goto 7.5}\end{aligned}$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} && \text{DComp(7.21)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot \frac{du}{dx} && \text{DNL(7.18)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot (6x - 6) && \text{DPo(7.6)} \\ \frac{dy}{dx} &= \frac{1}{\frac{3x^2 - 6x + 4}{6x - 6}} \cdot (6x - 6) \\ \frac{dy}{dx} &= \frac{6x - 6}{3x^2 - 6x + 4} && \text{OOM(1.4)}\end{aligned}$$

Example 7.11 — id:20141128-155506.

Differentiate $y = \ln(\cos x)$

S

Solution: After identifying that $y = \ln(\cos x)$ is a composite function, we let $u = \cos x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.21)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$\begin{aligned}y &= \ln(u) & u &= \cos x \\ \frac{dy}{du} &= \frac{1}{u} & \frac{du}{dx} &= -\sin x\end{aligned}$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} && \text{DComp(7.21)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot \frac{du}{dx} && \text{DNL(7.18)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot -\sin x && \text{DPo(7.6)} \\ \frac{dy}{dx} &= \frac{1}{\cos x} \cdot -\sin x \\ \frac{dy}{dx} &= \frac{-\sin x}{\cos x} && \text{OOM(1.4)} \\ \frac{dy}{dx} &= -\tan x\end{aligned}$$

Example 7.12 — id:20141124-205219.

Differentiate $y = (x^2 - 1) \ln(3x)$

S

Solution:

$$y' = [x^2 - 3]' \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' \quad \text{DPr(7.10)}$$

$$y' = 2x \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' \quad \text{differentiate goto 7.4}$$

$$y' = 2x \cdot \ln x + (x^2 - 3) \cdot \frac{1}{x} \quad \text{differentiate goto 7.9}$$

$$y' = 2x \cdot \ln x + \frac{x^2 - 3}{x} \quad \text{OOM(1.4)}$$

$$y' = 2x^2 \ln x + \frac{x^2 - 3}{x} \quad \text{JTC(1.6)}$$

$$y' = \frac{2x^2 \ln x + (x^2 - 3)}{x} \quad \text{OOA(1.1)}$$

■

8. Integration

8.1 Integration Rules

Rule 8.1.1 — Integral of a Constant (IC).

$$\int c \, dx = cx + k \quad (8.1)$$

Rule 8.1.2 — Integral of a Constant Multiple (ICM).

$$\int cx \, dx = c \int x \, dx \quad (8.2)$$

Rule 8.1.3 — Integral of a Power (IPo).

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} \quad \text{for } x \neq -1 \quad (8.3)$$

Rule 8.1.4 — Integral of a Sum (IS).

$$\int f(x) + g(x) \, dx = \int f(x) \, dx + \int g(x) \, dx \quad (8.4)$$

Example 8.1 — id:20141105-172103.

Integrate $\int \frac{1}{4}x^2 - 2x + 4 \, dx$

S

Solution:

$$\begin{aligned}
 \int \frac{1}{4}x^2 - 2x + 4 \, dx &= \int \frac{1}{4}x^2 + -2x + 4 \, dx && \text{DOS(2.5a)} \\
 &= \int \frac{1}{4}x^2 \, dx + \int -2x \, dx + \int 4 \, dx && \text{IS(8.4)} \\
 &= \frac{1}{4} \int x^2 \, dx + -2 \int x \, dx + \int 4 \, dx && \text{ICM(8.2)} \\
 &= \frac{1}{4} \int x^2 \, dx + -2 \int x \, dx + 4(x) + c_1 && \text{IC(8.1)} \\
 &= \frac{1}{4} \left(\frac{x^3}{3} \right) + c_2 + -2 \left(\frac{x^2}{2} \right) + c_3 + 4(x) + c_1 && \text{IPo(8.3)} \\
 &= \frac{x^3}{12} + c_2 - x^2 + c_3 + 4x + c_1 && \text{OOM(1.4)} \\
 &= \frac{x^3}{12} - x^2 + 4x + \underbrace{c_1 + c_2 + c_3}_{\text{Just a constant}} && \text{CPA(2.8)} \\
 &= \frac{x^3}{12} - x^2 + 4x + C && \text{OOA(1.1)}
 \end{aligned}$$

D**Dependencies:**

example 8.2-20141105-144223

Theorem 8.1.5 — Fundamental Theorem of Calculus (FThmC). If f is continuous on the closed interval $[a, b]$ and F is the indefinite integral of f on $[a, b]$, then

$$\int_a^b f(x) \, dx = F(b) - F(a) \quad (8.5a)$$

$$= [F(x)]_a^b \quad (8.5b)$$

8.2 Volumes of Revolution

Definition 8.2.1 — Volume of Revolution: Method of Disks. Let f be a nonnegative and continuous function on the closed interval $[a, b]$, then the solid of revolution obtained by rotating the curve $f(x)$ about the x -axis from $x = a$ to $x = b$ has volume given by

$$V = \pi \int_a^b [f(x)]^2 \, dx \quad (8.6)$$

Example 8.2 — id:20141105-144223.

Find the volume of the solid formed by rotating the region enclosed by $y = 2 - \frac{x}{2}$ and $y = 0$ through 2π about the x -axis on the closed interval $[2, 3]$.

S**Solution:**

$$\begin{aligned}
 V &= \pi \int_a^b [f(x)]^2 dx && \text{VRD(8.6)} \\
 &= \pi \int_2^3 \left[2 - \frac{x}{2}\right]^2 dx && \text{SPE(4.2)} \\
 &= \pi \int_2^3 \frac{1}{4}x^2 + -2x + 4 dx && \text{simplify: goto 2.39} \\
 &= \pi \left[\frac{x^3}{12} - x^2 + 4x \right]_2^3 && \text{integrate goto 8.1} \\
 &= \pi \left[\frac{[3]^3}{12} - [3]^2 + 4[3] - \left(\frac{[2]^3}{12} - [2]^2 + 4[2] \right) \right] && \text{FThmC(8.5a)} \\
 &= \pi \left(\frac{7}{12} \right) && \text{evaluate: example 5.1} \\
 &= \frac{7}{12}\pi && \text{CPM(1.8)}
 \end{aligned}$$

■

Example 8.3 — id:20141106-114907.

Find the volume of the solid formed by rotating the region enclosed by $y = \sqrt{9 - x^2}$ and $y = 0$ through 2π about the x -axis on the closed interval $[1, 3]$.

S**Solution:**

$$\begin{aligned}
 V &= \pi \int_a^b [f(x)]^2 dx && \text{VRD(8.6)} \\
 &= \pi \int_1^3 \left[\sqrt{9 - x^2}\right]^2 dx && \text{SPE(4.2)} \\
 &= \pi \int_1^3 9 - x^2 dx && \text{simplify: goto 2.32} \\
 &= \pi \left[9x - \frac{x^3}{3}\right]_1^3 && \text{integrate} \\
 &= \pi \left[9[3] - \frac{[3]^3}{3} - \left(9[1] - \frac{[1]^3}{3}\right)\right] && \text{FThmC(8.5a)} \\
 &= \pi \left(\frac{28}{3}\right) && \text{evaluate} \\
 &= \frac{28}{3}\pi && \text{CPM(1.8)}
 \end{aligned}$$

Example 8.4 — id:20141106-122528.

Find the volume of the solid formed by rotating the region enclosed by $y = \sqrt{\sin x}$ and $y = 0$ through 2π about the x -axis on the closed interval $[\frac{\pi}{2}, \pi]$.

S**Solution:**

$$\begin{aligned}
 V &= \pi \int_a^b [f(x)]^2 dx && \text{VRD(8.6)} \\
 &= \pi \int_{\pi/2}^{\pi} [\sqrt{\sin \theta}]^2 d\theta && \text{SPE(4.2)} \\
 &= \pi \int_{\pi/2}^{\pi} \sin \theta d\theta && \text{PoPo(2.12a)} \\
 &= \pi [-1 \cos \theta]_{\pi/2}^{\pi} && \text{integrate} \\
 &= \pi \left[-1 \cos [\pi] - \left(-1 \cos \left[\frac{\pi}{2} \right] \right) \right] && \text{FThmC(8.5a)} \\
 &= \pi(1) && \text{evaluate} \\
 &= \pi && \text{MId(2.1b)}
 \end{aligned}$$

Example 8.5 — id:20141108-083108.

Find the volume of the solid formed by rotating the region enclosed by $y = \sqrt{2 - x^2}$ and $y = 0$ through 2π about the x -axis.

S

Since no closed interval is stated, we will need to find where the curves $y = \sqrt{2 - x^2}$ and $y = 0$ intersect to determine the closed interval. Solving this system of equations (see example 4.9), we find that $x = \pm\sqrt{2}$.

$$\begin{aligned}
 V &= \pi \int_a^b [f(x)]^2 dx && \text{VRD(8.6)} \\
 &= \pi \int_{-\sqrt{2}}^{\sqrt{2}} [\sqrt{2 - x^2}]^2 dx \\
 &= \pi \int_{-\sqrt{2}}^{\sqrt{2}} 2 - x^2 dx && \text{PoPo(2.12a)} \\
 &= \pi \left[2x - \frac{x^3}{3} \right]_{-\sqrt{2}}^{\sqrt{2}} && \text{integrate} \\
 &= \pi \left[2[\sqrt{2}] - \frac{[\sqrt{2}]^3}{3} - \left(2[-\sqrt{2}] - \frac{[-\sqrt{2}]^3}{3} \right) \right] && \text{FThmC(8.5a)} \\
 &= \pi \left(\frac{8}{2} \sqrt{2} \right) && \text{simplify goto 2.32} \\
 &= \frac{8}{2} \pi \sqrt{2} && \text{CPM(1.8)}
 \end{aligned}$$

Example 8.6 — id:20141125-084541.

Find the volume of the solid formed by rotating the region enclosed by $y = \sqrt{2 - x^2}$ and $y = 0$ through 2π about the x -axis.

S

Since no closed interval is stated, we will need to find where the curves $y = 4 - x^2$ and $y = 0$ intersect to determine the closed interval. Solving this system of equations (see the similar example 4.9), we find that $x = \pm 2$.

Since the parabola $y = 4 - x^2$ is symmetric about the y -axis, we can find the volume on the closed interval $[0,2]$ and multiply this by a factor of two to include the volume on the closed interval $[-2,0]$.

S

Solution:

$$\begin{aligned} V &= 2 \cdot \pi \int_a^b [f(x)]^2 dx && \text{VRD(8.6)} \\ &= 2 \cdot \pi \int_0^2 [4 - x^2]^2 dx \\ &= 2 \cdot \pi \int_0^2 x^4 - 8x^2 + 16 dx \\ &= 2 \cdot \pi \left[\frac{x^5}{5} - \frac{8x^3}{3} + 16x \right]_0^2 \\ &= 2 \cdot \pi \left(\frac{[2]^5}{5} - \frac{8[2]^3}{3} + 16[2] \right) && \text{FThmC(8.5a)} \\ &= \frac{512\pi}{15} \end{aligned}$$

■

id:20141120-211843
id:20141121-195608
id:20141121-191224
id:20141123-091420

Websites

9. Answers

9.0.1 id:20141120-211843

- | | | | |
|----------|----------|----------|-----------|
| 1. $17x$ | 4. $9x$ | 7. $11x$ | 10. $19x$ |
| 2. $8x$ | 5. $9x$ | 8. $8x$ | 11. $26x$ |
| 3. $22x$ | 6. $22x$ | 9. $18x$ | 12. $8x$ |

View questions from exercise: 2.1.

9.0.2 id:20141121-195608

- | | | | |
|----------|----------|----------|-----------|
| 1. $13x$ | 4. $11x$ | 7. $3x$ | 10. $7x$ |
| 2. $6x$ | 5. $13x$ | 8. $8x$ | 11. $14x$ |
| 3. $8x$ | 6. $4x$ | 9. $13x$ | 12. $5x$ |

View questions from exercise: 2.2.

9.0.3 id:20141121-191224

- | | | | |
|---------|----------|-----------|-----------|
| 1. $-x$ | 4. $-6x$ | 7. $-5x$ | 10. $5x$ |
| 2. $3x$ | 5. x | 8. $-10x$ | 11. $-6x$ |
| 3. $-x$ | 6. $9x$ | 9. $-2x$ | 12. $-3x$ |

View questions from exercise: 2.3.

9.0.4 id:20141123-091420

- | | | | |
|-----------|-----------|----------|-----------|
| 1. $9x$ | 4. $-14x$ | 7. $-9x$ | 10. $7x$ |
| 2. $-30x$ | 5. $4x$ | 8. $10x$ | 11. $-4x$ |
| 3. $-3x$ | 6. $-10x$ | 9. $-5x$ | 12. $16x$ |

View questions from exercise: 2.4.



Bibliography

Website

- [] *Polynomial.* <http://en.wikipedia.org/wiki/Polynomial> (cited on page 12).
- [Weia] Eric W. Weisstein. "Asymptote." From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/Asymptote.html> (cited on page 61).
- [Weib] Eric W. Weisstein. "Completing the Square." From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/CompletingtheSquare.html> (cited on page 60).



Index

A

Additive Identity	15
Additive Inverse	9
Algebraic Expressions	5
ans20141120-211843	81
ans20141121-191224	81
ans20141121-195608	81
ans20141123-091420	81
Asympote	61

C

Classification of Multivariate Polynomial Expressions	41
Classification of Univariate Polynomial Ex- pressions	11
Coefficient	11
Common Denominator	6

D

Definition of Subtraction	17
Degree n Multivariate Polynomials	45
Degree n Univariate Polynomials	40
Degree -1 Multivariate Polynomials ..	41
Degree -1 Univariate Polynomials ..	13
Degree 0 Multivariate Polynomials ..	41
Degree 0 Univariate Polynomials ..	13

Degree 1 Multivariate Polynomials ..	41
Degree 1 Univariate Polynomials	13
Degree 2 Multivariate Polynomials ..	41
Degree 2 Univariate Polynomials	30
Degree 3 Multivariate Polynomials ..	44
Degree 3 Univariate Polynomials	40
Degree of the Indeterminate	12
Degree of the Univariate Polynomial .	13
Derivatives	
Derivative of a Composite Function	
71	
Derivative of a Constant	67
Derivative of a Constant Multiple	67
Derivative of a Natural Logarithm	70
Derivative of a Power	67
Derivative of a Product	70
Derivative of a Quotient	70
Derivative of a Sum	68

E

Equality	49
Example	
20141027-075159	16
20141105-144223	76
20141105-161225	38
20141105-172103	75
20141106-083703	59
20141106-114907	77
20141106-122528	78
20141106-150622	19
20141106-150953	41

20141106-151138	30	20141121-190857	18
20141106-151409	44	20141121-191224	19
20141106-151622	43	20141121-193636	18
20141106-152020	20	20141121-195608	17
20141106-152339	32	20141124-141850	68
20141106-154547	30	20141124-152503	69
20141107-121834	33	20141124-153017	67
20141107-131748	54	20141124-203850	71
20141108-083108	78	20141124-205219	73
20141108-085327	9	20141125-084541	79
20141108-173613	23	20141128-151834	69
20141108-180100	42	20141128-155506	72
20141108-180608	43	20141128-160248	72
20141108-180858	44	20141206-101107	51
20141108-191616	32	20141206-101632	50
20141108-192234	46	20141206-102142	49
20141108-192611	46	20141206-102404	51
20141108-193145	45	20141206-102723	52
20141108-193835	42	20141206-104404	52
20141108-194156	22		
20141108-194431	21	Exercise	
20141108-194709	31	20141120-211843	16
20141109-090809	23	20141123-091420	21
20141109-091015	24		
20141109-092448	24		
20141109-092651	25		
20141109-092910	26		
20141109-093220	26		
20141109-093419	27		
20141109-094928	27		
20141109-095151	28		
20141109-095536	28		
20141109-095842	29		
20141109-133008	34		
20141109-133316	35		
20141109-140659	36		
20141109-141019	37		
20141109-141347	37		
20141111-190212	61		
20141111-192213	62		
20141111-215726	53		
20141111-222931	50		
20141120-202042	14		
20141120-202842	12		
20141120-203846	15		
20141121-093439	14		
20141121-093747	13		
20141121-184652	20		
20141121-185558	16		

F

Fraction Operation of Addition 6

I

Indeterminate 11

Integrals

Integral of a Constant 75

Integral of a Constant Multiple 75

Integral of a Power 75

Integral of a Sum 75

L

Like Terms 14

M

Method of Disks 76

Monomials of Like Terms 9

Multiplicative Identity 12

Multiplicative Inverse 7

N

Notation

- Center-Dot to Juxtaposition 7
- Juxtaposition to Center-Dot 7

S

- Substitution Property of Equality 49
- Surds 9
- Symmetric Property of Equality 49

O

Operation

- Operation of Addition 5
- Operation of Exponentiation 7
- Operation of Multiplication 7
- Operation of Negation 17

P

- Polynomial Expressions 5
- Power Inverse 9
- Powers 7
 - Factor to Power 9
 - Power of a Power 32
 - Power of a Product 45
 - Power of a Product of Powers 9
 - Power of a Quotient of Powers 7
 - Power to Factor 9
 - Power to Radical 9
 - Product of Common Base Powers 31
 - Quotient of Common Base Powers 32
 - Radical to Power 9

Property

- Associative Property of Addition 20
- Associative Property of Multiplication 7
- Commutative Property of Addition 23
- Commutative Property of Multiplication 7
- Distributive Property Factoring 15, 23

R

- Reflexive Property of Equality 49

T

Theorems

- Fundamental Theorem of Calculus 76
- Transitive Property of Equality 49

U

- Univariate Monomial 11
- Univariate Polynomial Expression 12

V

- Vector 65