

Mathematics Notebook

by Mark Olson : v42.org

Updated: October 15, 2015

Contents

I	Reference
1	Numbers 9
1.1	Number Systems 9
1.2	Prime Numbers 9
1.2.1	Listing of Prime Numbers 2-997 10
2	Operations 11
2.1	Dyadic Operations 11
3	Notation 13
3.1	Negation Notation 13
3.2	Multiplication Notation 13
3.3	Power Notation 14
3.4	Logarithm Notation 14
3.5	Derivative Notation 14
4	Properties 17
4.1	Summary of Field Properties 17
4.2	Properties of Addition 17
4.3	Properties of Multiplication 18
4.4	Properties of Subtraction 19
4.5	Properties of Powers 19
4.6	Properties of Equality 19
4.7	Properties of Inequality 20
5	Identities 21
5.1	Power Identities 21
5.2	Logarithm Identities 22
5.3	Trigonometric Identities 22
6	Calculus Rules 27
6.1	Monomial Derivative Rules 27
6.2	Derivative Structural Rules 27

6.3	Trigonometric Derivative Rules	28
6.4	Logarithm Derivative Rules	29
6.5	Exponential Derivative Rules	29

II Algebra

7	Simplifying Univariate Polynomials	33
7.1	Simplifying Degree 0 Univariate Monomials	33
7.2	Simplifying Degree 1 Univariate Monomials	35
7.3	Simplifying Degree 2 Univariate Monomials	41
7.4	Simplifying Degree 3 Univariate Monomials	44
7.5	Simplifying Degree 1 Univariate Binomials	44
7.6	Simplifying Degree 2 Univariate Binomials	50
7.7	Simplifying Degree 2 Univariate Trinomials	53
7.8	Simplifying Degree 1 Univariate Polynomials	57
7.9	Simplifying Degree 2 Univariate Polynomials	57
7.10	Simplifying Degree n Univariate Polynomials	57
8	Simplifying Univariate Monomials	59
8.1		59
9	Solving Linear Equations	61
9.1	Power Inverse	61
10	Solving Quadratic Equations	67
10.1	Power Inverse	67
10.2	Completing The Square	68

III Functions

11	Functions	75
11.1	Inverse Functions	76
11.2	Inverses	76

IV Differential Calculus

12	Derivative by First Principles	79
12.1	Limit of the Difference Quotient	79
13	Derivative Rules	81
13.1	Derivative of a Monomial Functions	81

13.2	Derivative of Polynomial Functions	82
13.3	Derivative of a Quotient	88
13.4	Derivative of a Rational Function	89
14	Equations of Tangent & Secant Lines	91
14.1	Essential Questions	91
14.2	Finding the Equation of the Tangent Line	91
15	First Derivative Test	95
16	Curve Sketching	97
16.0.1	Finding the vertex of a quadratic function using differentiation.	97
17	Second Derivative Test	99
18	Curve Sketching	101
	Bibliography	103
	Bibliography	103
	Websites	103
	Articles	103
	Index	105

Reference

1	Numbers	9
1.1	Number Systems	
1.2	Prime Numbers	
2	Operations	11
2.1	Dyadic Operations	
3	Notation	13
3.1	Negation Notation	
3.2	Multiplication Notation	
3.3	Power Notation	
3.4	Logarithm Notation	
3.5	Derivative Notation	
4	Properties	17
4.1	Summary of Field Properties	
4.2	Properties of Addition	
4.3	Properties of Multiplication	
4.4	Properties of Subtraction	
4.5	Properties of Powers	
4.6	Properties of Equality	
4.7	Properties of Inequality	
5	Identities	21
5.1	Power Identities	
5.2	Logarithm Identities	
5.3	Trigonometric Identities	
6	Calculus Rules	27
6.1	Monomial Derivative Rules	
6.2	Derivative Structural Rules	
6.3	Trigonometric Derivative Rules	
6.4	Logarithm Derivative Rules	
6.5	Exponential Derivative Rules	

1. Numbers

1.1 Number Systems

Definition 1.1.1 – Natural Numbers.

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

-  It is not uncommon for zero to be excluded from the natural numbers. In fact, some exclude zero from the natural numbers and then describe the set of natural numbers that include zero the whole numbers.

$$\mathbb{W} = \{0, 1, 2, 3, \dots\}$$

For the purposes of these notes, zero will be included within the set of natural numbers.

Definition 1.1.2 – Integers.

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Definition 1.1.3 – Positive Integers.

$$\mathbb{Z}^+ = \{1, 2, 3, \dots\}$$

Definition 1.1.4 – Rational Numbers.

$$\mathbb{Q} = \{m/n \mid m, n \in \mathbb{Z}, n \neq 0\}$$

Definition 1.1.5 – Proper Fraction. Given $m < n$, then the fraction m/n is called **proper**.

Definition 1.1.6 – Improper Fraction. Given $m > n$, then the fraction m/n is called **improper**.

1.2 Prime Numbers

Definition 1.2.1 – Greatest Common Divisor. Suppose that m and n are positive integers. The greatest common divisor is the largest divisor (factor) common to both m and n .

Definition 1.2.2 – Relatively Prime. Two integers m and n are relatively prime to each other, $m \perp n$, if they share no common positive integer divisors (factors) except 1.

$$m \perp n \text{ if } \gcd(m, n) = 1.$$

1.2.1 Listing of Prime Numbers 2-997

2	3	5	7	11	13	17	19	23	29	31	37
41	43	47	53	59	61	67	71	73	79	83	89
97	101	103	107	109	113	127	131	137	139	149	151
157	163	167	173	179	181	191	193	197	199	211	223
227	229	233	239	241	251	257	263	269	271	277	281
283	293	307	311	313	317	331	337	347	349	353	359
367	373	379	383	389	397	401	409	419	421	431	433
439	443	449	457	461	463	467	479	487	491	499	503
509	521	523	541	547	557	563	569	571	577	587	593
599	601	607	613	617	619	631	641	643	647	653	659
661	673	677	683	691	701	709	719	727	733	739	743
751	757	761	769	773	787	797	809	811	821	823	827
829	839	853	857	859	863	877	881	883	887	907	911
919	929	937	941	947	953	967	971	977	983	991	997

2. Operations

2.1 Dyadic Operations

Definition 2.1.1 – Operation of Addition (OOA).

$$\underbrace{\begin{array}{c} a \\ \text{Augend} \end{array}}_{\text{Sum}} + \underbrace{\begin{array}{c} b \\ \text{Addend} \end{array}}_{\text{Sum}} \quad (2.1)$$

More generally,

$$\underbrace{\begin{array}{c} a \\ \text{Summand} \end{array}}_{\text{Sum}} + \underbrace{\begin{array}{c} b \\ \text{Summand} \end{array}}_{\text{Sum}} \quad (2.2)$$

Definition 2.1.2 – Operation of Multiplication (OOM).

$$\underbrace{\begin{array}{c} a \\ \text{Multiplicand} \end{array}}_{\text{Product}} \times \underbrace{\begin{array}{c} b \\ \text{Multiplier} \end{array}}_{\text{Product}} \quad (2.3)$$

More generally,

$$\underbrace{\begin{array}{c} a \\ \text{Factor} \end{array}}_{\text{Product}} \times \underbrace{\begin{array}{c} b \\ \text{Factor} \end{array}}_{\text{Product}} \quad (2.4)$$

Definition 2.1.3 – Operation of Exponentiation (OOE).

$$\underbrace{\begin{array}{c} b \\ \text{base} \end{array}}_{\text{Power}}^m \quad (2.5)$$

Definition 2.1.4 – Common Denominator (CD).

$$\frac{a}{b} + \frac{c}{b} = \frac{a+c}{b} \quad (2.6a)$$

$$\frac{a+c}{b} = \frac{a}{b} + \frac{c}{b} \quad (2.6b)$$

Rule 2.1.1 – Fraction Operation of Addition (FOOA).

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd} \quad (2.7a)$$

$$\frac{ad + bc}{bd} = \frac{a}{b} + \frac{c}{d} \quad (2.7b)$$

3. Notation

3.1 Negation Notation

Notation 3.1.1 – Operation of Negation (ONeg).

$$\neg a = \neg a \quad (3.1a)$$

$$\neg a = -a \quad (3.1b)$$

I have used a different symbol, \neg , as the prefix negation operator only to differentiate it from the minus sign infix operator symbol, $-$, which is also used as the infix operator for the dyadic operation of subtraction. I will refer to this change of symbol as ONeg. This is used only as a teaching tool and should not be confused with the logic negation operator. Another advantage of using this symbol is that it reduces the number of delimiters used in an expression for example, $\neg a$ versus $(-a)$.

- Negative five: -5
- Negative five: $\neg 5$
- Four minus five: $4 - 5$
- Four minus negative five: $4 - \neg 5$
- Four minus negative five: $4 - (-5)$
- Four minus negative five: $4 - \neg 5$
- Negative four minus five: $-4 - 5$
- Negative four minus five: $\neg 4 - 5$

3.2 Multiplication Notation

Notation 3.2.1 – Multiplication Center-Dot (MC).

$$a \cdot b \quad (3.2)$$

Notation 3.2.2 – Multiplication Juxtaposition (MJ).

$$ab, a(b), (a)b, (a)(b), a[b], [a]b, [a][b] \quad (3.3)$$

Notation 3.2.3 – Multiplication Times (MT).

$$a \times b \quad (3.4)$$

Notation 3.2.4 – Juxtaposition to Center-Dot (JTC).

$$ab = a \cdot b \quad (3.5)$$

Notation 3.2.5 – Center-Dot to Justaposition (CTJ).

$$a \cdot b = ab \quad (3.6)$$

3.3 Power Notation

Notation 3.3.1 – Power Exponent Negative Exponent (PoNegE).

$$b^{-k} = \frac{1}{b^k} \quad (3.7)$$

$$\frac{1}{b^k} = b^{-k} \quad (3.8)$$

Notation 3.3.2 – Power To Factor (PoTF).

$$a^n = a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_n \quad (3.9)$$

Notation 3.3.3 – Power To Logarithm (PoTL).

$$y = b^x \Rightarrow x = \log_b y \quad (3.10)$$

Notation 3.3.4 – Factor To Power (FTPo).

$$a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_n = a^n \quad (3.11)$$

Notation 3.3.5 – Radical To Power (RTPo).

$$\sqrt[m]{b^n} = b^{\frac{n}{m}} \quad (3.12)$$

3.4 Logarithm Notation

Notation 3.4.1 – Logarithm Exponent Visible (LEV).

$$\log_b y \Rightarrow \log_b y = x \quad (3.13)$$

Notation 3.4.2 – Logarithm to Power (LTPo).

$$x = \log_b y \Rightarrow y = b^x \quad (3.14)$$

3.5 Derivative Notation

Notation 3.5.1 – Leibniz's first derivative.

$$\frac{dy}{dx} = \frac{d[f(x)]}{dx} = \frac{d}{dx}[f(x)] \quad (3.15)$$

Notation 3.5.2 – Leibniz's second derivative.

$$\frac{d^2y}{dx^2} \quad (3.16)$$

Notation 3.5.3 – Leibniz's nth derivative.

$$\frac{d^n y}{dx^n} \quad (3.17)$$

Notation 3.5.4 – Leibniz's Evaluate derivative.

$$\left. \frac{dy}{dx} \right|_{x=a} = \frac{dy}{dx}(a) \quad (3.18)$$

Notation 3.5.5 – LaGrange's first derivative.

$$f'(x) \quad (3.19)$$

Notation 3.5.6 – LaGrange's second derivative.

$$f''(x) \quad (3.20)$$

Notation 3.5.7 – LaGrange's nth derivative.

$$f^{(n)}(x) \quad (3.21)$$

Notation 3.5.8 – LaGrange's Evaluate derivative.

$$f'(a) \quad (3.22)$$

Notation 3.5.9 – Euler's first derivative.

$$Df = D_x f \quad (3.23)$$

Notation 3.5.10 – Euler's second derivative.

$$D^2 f = D_x^2 f \quad (3.24)$$

Notation 3.5.11 – Euler's nth derivative.

$$D^n f = D_x^n \quad (3.25)$$

4. Properties

4.1 Summary of Field Properties

Name	Addition	Multiplication
Commutative	$a + b = b + a$	$a \cdot b = b \cdot a$
Associative	$(a + b) + c = a + (b + c)$	$(a \cdot b) \cdot c = a \cdot (b \cdot c)$
Distributive	$a(b + c) = ab + ac$	$(a + b)c = ac + bc$
Identity	$a + \mathbf{0} = a = \mathbf{0} + a$	$a \cdot \mathbf{1} = a = \mathbf{1} \cdot a$
Inverse	$a + (-a) = 0 = (-a) + a$	$a \cdot a^{-1} = 1 = a^{-1} \cdot a$

Table 4.1: Summary of the Field Properties

4.2 Properties of Addition

Property 4.2.1 – Commutative Property of Addition (CPA).

$$ab = ba \quad (4.1)$$

Property 4.2.2 – Associative Property of Addition (APA).

$$a + b + c = (a + b) + c \quad (4.2a)$$

$$a + b + c = a + (b + c) \quad (4.2b)$$

Property 4.2.3 – Distributive Property Factoring (DPF).

$$ba + ca = (b + c)a \quad (4.3a)$$

$$ab + ac = a(b + c) \quad (4.3b)$$

Property 4.2.4 – Additive Identity (AId).

$$a + \mathbf{0} = a \quad (4.4a)$$

$$a = a + \mathbf{0} \quad (4.4b)$$

Property 4.2.5 – Additive Inverse (AI).

$$a + (-a) = 0 \quad (4.5a)$$

4.3 Properties of Multiplication

Property 4.3.1 – Commutative Property of Multiplication (CPM).

$$a \cdot b = b \cdot a \quad (4.6)$$

Property 4.3.2 – Associative Property of Multiplication (APM).

$$a \cdot b \cdot c = (a \cdot b) \cdot c \quad (4.7a)$$

$$a \cdot b \cdot c = a \cdot (b \cdot c) \quad (4.7b)$$

Property 4.3.3 – Distributive Property Expanding (DPE).

$$a(b + c) = ab + ac \quad (4.8a)$$

$$(b + c)a = ba + ca \quad (4.8b)$$

Property 4.3.4 – Multiplicative Identity (Mid).

$$1a = a \quad (4.9a)$$

$$a = 1a \quad (4.9b)$$

- R** If the coefficient of a univariate monomial is the multiplicative identity 4.9a, 1, then it is not shown in its canonical form.

$$\begin{aligned} C_k x^k &= C_k x^k \\ &= 1x^k \\ &= x^k \end{aligned}$$

Property 4.3.5 – Multiplicative Inverse (MI).

$$a \cdot \frac{1}{a} = 1 \quad (4.10a)$$

$$a \cdot a^{-1} = 1 \quad (4.10b)$$

Property 4.3.6 – Zero Product (ZPr).

$$\text{if } a \cdot b = 0, \text{ then } a = 0 \text{ or } b = 0 \quad (4.11a)$$

4.4 Properties of Subtraction

Definition 4.4.1 – Definition of Subtraction (DOS).

$$a - b = a + \neg b \quad (4.12a)$$

$$a + \neg b = a - b \quad (4.12b)$$

4.5 Properties of Powers

Property 4.5.1 – Power Inverse (Pold).

$$1 = b^0 \quad (4.13a)$$

$$b^0 = 1 \quad (4.13b)$$

4.6 Properties of Equality

Property 4.6.1 – Reflexive Property of Equality (RPE).

$$a = a \quad (4.14a)$$

Property 4.6.2 – Substitution Property of Equality (SPE).

Given $a = b$, then

$$E(a) = E(b) \quad (4.15)$$

$E(x)$ represents any expression.

Property 4.6.3 – Symmetric Property of Equality (SyPE).

$$a = b \quad \text{then} \quad b = a \quad (4.16a)$$

Property 4.6.4 – Transitive Property of Equality (TPE).

$$\text{if } a = b \quad \text{and} \quad b = c \quad \text{then} \quad a = c \quad (4.17a)$$

Property 4.6.5 – Zero Factor Property (ZFP).

$$\text{if } a \cdot b = 0 \text{ then } a = 0 \text{ or } b = 0 \quad (4.18a)$$

4.7 Properties of Inequality

Property 4.7.1 – Substitution Property of Inequality (SPIn).

$$a < b \text{ then } a + c < b + c \quad (4.19a)$$

$$a < b \text{ and } c > 0, \text{ then } ca < cb \quad (4.19b)$$

$$a < b \text{ and } c < 0, \text{ then } ca > cb \quad (4.19c)$$

Property 4.7.2 – Transitive Property of Inequality (TPIn).

$$\text{if } a < b \text{ and } b < c \text{ then } a > c \quad (4.20a)$$

5. Identities

5.1 Power Identities

Identity 5.1.1 – Power of Power (PoPo).

$$(b^m)^k = b^{m \cdot k} \quad (5.1a)$$

$$b^{m \cdot k} = (b^m)^k \quad (5.1b)$$

Identity 5.1.2 – Power of a Product (PoPr).

$$(a \cdot b)^k = a^k \cdot b^k \quad (5.2a)$$

$$a^k \cdot b^k = (a \cdot b)^k \quad (5.2b)$$

Identity 5.1.3 – Product Common Base Powers (PrCBPo).

$$b^m \cdot b^n = b^{m+n} \quad (5.3a)$$

$$b^{m+n} = b^m \cdot b^n \quad (5.3b)$$

Identity 5.1.4 – Quotient Common Base Powers (QCBPo).

$$\frac{b^m}{b^n} = b^{m-n} \quad (5.4a)$$

$$b^{m-n} = \frac{b^m}{b^n} \quad (5.4b)$$

Identity 5.1.5 – Power of a Quotient of Powers (PoQPo).

$$\left(\frac{a^m}{b^n}\right)^k = \frac{a^{m \cdot k}}{b^{n \cdot k}} \quad (5.5a)$$

$$\frac{a^{m \cdot k}}{b^{n \cdot k}} = \left(\frac{a^m}{b^n}\right)^k \quad (5.5b)$$

Identity 5.1.6 – Power of a Product of Powers (PoPrPo).

$$(a^m \cdot b^n)^k = a^{m \cdot k} \cdot b^{n \cdot k} \quad (5.6a)$$

$$a^{m \cdot k} \cdot b^{n \cdot k} = (a^m \cdot b^n)^k \quad (5.6b)$$

5.2 Logarithm Identities

Identity 5.2.1 – Logarithm Power of a Power (LPoPo).

$$\log_b x^n = n \log_b x \quad (5.7a)$$

$$n \log_b x = \log_b x^n \quad (5.7b)$$

Identity 5.2.2 – Logarithm Product of Common Base Powers (LPrCBPo).

$$\log_b(mn) = \log_b m + \log_b n \quad (5.8a)$$

$$\log_b m + \log_b n = \log_b(mn) \quad (5.8b)$$

Identity 5.2.3 – Logarithm Quotient of Common Base Powers (LQCBPo).

$$\log_b \left(\frac{m}{n} \right) = \log_b m - \log_b n \quad (5.9a)$$

$$\log_b m - \log_b n = \log_b \left(\frac{m}{n} \right) \quad (5.9b)$$

5.3 Trigonometric Identities

Identity 5.3.1 – Trigonometric Reciprocal Identities (TRId).

$$\sin \theta = \frac{1}{\csc \theta} \quad (5.10a)$$

$$\cos \theta = \frac{1}{\sec \theta} \quad (5.10b)$$

$$\cot \theta = \frac{1}{\tan \theta} \quad (5.10c)$$

$$\csc \theta = \frac{1}{\sin \theta} \quad (5.10d)$$

$$\sec \theta = \frac{1}{\cos \theta} \quad (5.10e)$$

$$\tan \theta = \frac{1}{\cot \theta} \quad (5.10f)$$

Identity 5.3.2 – Trigonometric Pythagorean Identities (TPythagId).

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (5.11a)$$

$$\sec^2 \theta = \tan^2 \theta + 1 \quad (5.11b)$$

$$\csc^2 \theta = 1 + \cot^2 \theta \quad (5.11c)$$

Identity 5.3.3 – Trigonometric Tangent Identity (TanId).

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad (5.12a)$$

Identity 5.3.4 – Trigonometric Cotangent Identity (CotId).

$$\cot \theta = \frac{\cos \theta}{\sin \theta} \quad (5.13a)$$

Identity 5.3.5 – Sine Double Angle Identity (SinDAId).

$$\sin 2\theta = 2 \sin \theta \cos \theta \quad (5.14a)$$

Identity 5.3.6 – Cosine Double Angle Identity (CosDAId).

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta \quad (5.15a)$$

$$= 1 - 2 \sin^2 \theta \quad (5.15b)$$

$$= 2 \cos^2 \theta - 1 \quad (5.15c)$$

Identity 5.3.7 – Tangent Double Angle Identity (TanDAId).

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} \quad (5.16a)$$

Identity 5.3.8 – Sine Sum of Angles Identity (SinSAId).

$$\sin(\theta + \phi) = \sin \theta \cos \phi + \cos \theta \sin \phi \quad (5.17a)$$

Identity 5.3.9 – Sine Difference of Angles Identity (SinDiffAId).

$$\sin(\theta - \phi) = \sin \theta \cos \phi - \cos \theta \sin \phi \quad (5.18a)$$

Identity 5.3.10 – Cosine Sum of Angles Identity (CosSAId).

$$\cos(\theta + \phi) = \cos \theta \cos \phi - \sin \theta \sin \phi \quad (5.19a)$$

Identity 5.3.11 – Cosine Difference of Angles Identity (CosDiffAId).

$$\cos(\theta - \phi) = \cos \theta \cos \phi + \sin \theta \sin \phi \quad (5.20a)$$

Identity 5.3.12 – Tangent Sum of Angles Identity (TanSAId).

$$\tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} \quad (5.21a)$$

Identity 5.3.13 – Tangent Difference of Angles Identity (TanDiffAId).

$$\tan(\theta - \phi) = \frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi} \quad (5.22a)$$




Figure 5.1: Simplifying Expressions Workflow:

■ Property, ■ Operation, ■ Notation, ■ Powers, ■ Delimiters, ■ Process, ■ Not Used

6. Calculus Rules

6.1 Monomial Derivative Rules

Rule 6.1.1 – Derivative of a Constant (DC).

$$[c]' = 0 \quad (6.1)$$

$$\frac{d}{dx} [c] = 0 \quad (6.2)$$

Rule 6.1.2 – Derivative of a Constant Multiple (DCM).

$$[cf(x)]' = c[f(x)]' \quad (6.3)$$

$$\frac{d}{dx} [cf(x)] = c \frac{d}{dx} [f(x)] \quad (6.4)$$

Rule 6.1.3 – Derivative of a Power (DPo).

$$[x^n]' = nx^{n-1} \quad (6.5)$$

$$\frac{d}{dx} [x^n] = nx^{n-1} \quad (6.6)$$

6.2 Derivative Structural Rules

Rule 6.2.1 – Derivative of a Sum (DS).

$$[f(x) + g(x)]' = f'(x) + g'(x) \quad (6.7)$$

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} [f(x)] + \frac{d}{dx} [g(x)] \quad (6.8)$$

Rule 6.2.2 – Derivative of a Difference (DD).

$$[f(x) - g(x)]' = f'(x) - g'(x) \quad (6.9)$$

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} [f(x)] - \frac{d}{dx} [g(x)] \quad (6.10)$$

Rule 6.2.3 – Derivative of a Product (DPr).

$$[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x) \quad (6.11)$$

$$\frac{d}{dx} [f(x)g(x)] = \frac{d}{dx} [f(x)]g(x) + f(x)\frac{d}{dx} [g(x)] \quad (6.12)$$

Rule 6.2.4 – Derivative of a Quotient (DQ).

$$\left[\frac{f(x)}{g(x)} \right]' = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2} \quad (6.13)$$

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{\frac{d}{dx} [f(x)] g(x) - f(x) \frac{d}{dx} [g(x)]}{[g(x)]^2} \quad (6.14)$$

Rule 6.2.5 – Derivative of a Composite Function (DComp).

$$[f(g(x))]' = [g(x)]' [f(g(x))]' \quad (6.15)$$

$$\frac{d}{dx} [f(g(x))] = \frac{d}{dx} [g(x)] \frac{d}{dx} [f(g(x))] \quad (6.16)$$

6.3 Trigonometric Derivative Rules

Rule 6.3.1 – Derivative of Sine (DSin).

$$[\sin(x)]' = \cos(x) \quad (6.17)$$

$$\frac{d}{dx} [\sin(x)] = \cos(x) \quad (6.18)$$

Rule 6.3.2 – Derivative of Cosine (DCos).

$$[\cos(x)]' = -\sin(x) \quad (6.19)$$

$$\frac{d}{dx} [\cos(x)] = -\sin(x) \quad (6.20)$$

Rule 6.3.3 – Derivative of Tangent (DTan).

$$[\tan x]' = \sec^2 \quad (6.21)$$

$$\frac{d}{dx} [\tan x] = \sec^2 \quad (6.22)$$

Rule 6.3.4 – Derivative of Cosecant (DCsc).

$$[\csc x]' = -\csc x \cot x \quad (6.23)$$

$$\frac{d}{dx} [\csc x] = -\csc x \cot x \quad (6.24)$$

Rule 6.3.5 – Derivative of Secant (DSec).

$$[\sec x]' = \sec x \tan x \quad (6.25)$$

$$\frac{d}{dx} [\sec x] = \sec x \tan x \quad (6.26)$$

Rule 6.3.6 – Derivative of Cotangent (DCot).

$$[\cot x]' = -\csc^2 x \quad (6.27)$$

$$\frac{d}{dx} [\cot x] = -\csc^2 x \quad (6.28)$$

6.4 Logarithm Derivative Rules

Rule 6.4.1 – Derivative of a Logarithm (DL).

$$[\log_a x]' = \frac{1}{x \ln a} \quad (6.29)$$

$$\frac{d}{dx} [\log_a x] = \frac{1}{x \ln a} \quad (6.30)$$

Rule 6.4.2 – Derivative of a Natural Logarithm (DNL).

$$[\ln x]' = \frac{1}{x} \quad (6.31)$$

$$\frac{d}{dx} [\ln x] = \frac{1}{x} \quad (6.32)$$

6.5 Exponential Derivative Rules

Rule 6.5.1 – Derivative of an Exponential(DExp).


$$[a^x]' = a^x \ln a \quad (6.33)$$

$$\frac{d}{dx} [a^x] = a^x \ln a \quad (6.34)$$

Rule 6.5.2 – Derivative of a Natural Exponential(DNExp).

$$[e^x]' = e^x \quad (6.35)$$

$$\frac{d}{dx} [e^x] = e^x \quad (6.36)$$



Algebra

7	Simplifying Univariate Polynomials	33
7.1	Simplifying Degree 0 Univariate Monomials		
7.2	Simplifying Degree 1 Univariate Monomials		
7.3	Simplifying Degree 2 Univariate Monomials		
7.4	Simplifying Degree 3 Univariate Monomials		
7.5	Simplifying Degree 1 Univariate Binomials		
7.6	Simplifying Degree 2 Univariate Binomials		
7.7	Simplifying Degree 2 Univariate Trinomials		
7.8	Simplifying Degree 1 Univariate Polynomials		
7.9	Simplifying Degree 2 Univariate Polynomials		
7.10	Simplifying Degree n Univariate Polynomials		
8	Simplifying Univariate Monomials	59
8.1			
9	Solving Linear Equations	61
9.1	Power Inverse		
10	Solving Quadratic Equations	67
10.1	Power Inverse		
10.2	Completing The Square		

7. Simplifying Univariate Polynomials

The definition of a univariate polynomial expression is based on the expanded canonical form of some polynomial expression. It might be that the original expression might not be in the expanded canonical form, so a process called **simplifying by expanding** will be introduced to manipulate the expression such that it can be written in its expanded canonical form.

This process of simplifying by expanding polynomial expressions will be developed to the extent that it will be used to simplify multivariate polynomials. We will start by simplifying univariate monomial expressions.

7.1 Simplifying Degree 0 Univariate Monomials

Definition 7.1.1 – Indeterminate.

x

An indeterminate is a symbol that is treated as a variable, but does not stand for anything else but itself and is used as a placeholder.

- it does **not** designate a constant or a parameter
- it is **not** an unknown that could be solved for
- it is **not** a variable designating a function argument

[1]

Definition 7.1.2 – Degree of the Indeterminate.

x^k

The exponent of an indeterminate power, k is called the degree of the indeterminate.

Polynomial [3]

Definition 7.1.3 – Coefficient.

Cx^k

A coefficient, C is a real number multiplicative factor.

Definition 7.1.4 – Univariate Monomial.

$C_k x^k$

A univariate monomial is made up of two factors. The first factor of a monomial, C_k , is the **coefficient**. The second factor of each monomial, x^k , is an indeterminate raised to a non-negative integer power k .

Degree 0 univariate polynomial expressions are made up of univariate monomials, C_0 , called **constants**. The power identity is an indeterminate raised to a power of 0 has a value of 1. Thus, $x^0 = 1$ and results in the monomial $C_0 \cdot 1$. The canonical form of a product does not show the multiplicative identity factor, so what remains of this monomial product is only the coefficient factor C_0 and from now on will be referred to as a **constant**.

Degree 0 univariate polynomial expressions are usually a monomial in their canonical form if C_0 is a non-zero real number. The exception is if $C_0 = 0$, the additive identity, then the result is the zero polynomial, which can be considered a degree -1 polynomial.

The expression can be manipulated into its monomial canonical form by simplifying the expression. Simplifying the expression can be defined as evaluating the expression by following order of operations, which is the same as evaluating an arithmetic expression.

Definition 7.1.5 – Univariate Like Terms.

$$C_1x^k = C_2x^k$$

Two or more univariate monomials are defined as having like terms if each monomial has the same term, which will be the same indeterminate raised to the same positive integer power.

Sometimes the word **term** is used to describe monomials (including both the coefficient and the term), which may be confusing when trying to define like terms. For this reason, we will refer to the summands of a polynomial as monomials.

The monomials $5x^1$ and $3x^1$ can be described as having like terms because they share the common term x^1 . One could also say that $5x^1$ and $3x^1$ are like terms by definition and consequently giving the reader an impression that $5x^1$ and $3x^1$ are terms themselves.



A degree 1 indeterminate does not display the multiplicative identity in the exponent when its in canonical form.

Example 7.1 – id:20141121-093747.

Express $13x^0$ in canonical form.



Solution:

7.2 Simplifying Degree 1 Univariate Monomials

Example 7.2 – id:20141120-202042.

Express $5x^1$ in canonical form

(S) _____

Solution:

$$5x \quad \text{MId}(4.9b)$$

■

Example 7.3 – id:20141121-093439.

Express $7x^1 + 5$ in canonical form.

(S) _____

Solution:

$$7x + 5 \quad \text{MId}(4.9b)$$

■

If the constant monomial is 0, the additive identity, then the canonical form of a degree 1 univariate polynomial is a degree 1 monomial.

Example 7.4 – id:20141120-203846.

Simplify $6x + 7x$

(S) _____

Solution:

Notice that the indeterminate of each monomial is of degree 1; however, the exponent 1 is not shown. The monomials $6x$ and $7x$ have a like term of x .

$$\begin{array}{ll} (6 + 7)x & \text{DPF}(4.3a) \\ 13x & \text{OOA}(2.1) \end{array}$$

Notice that the sum of two monomials that have like terms can be found by adding the coefficients of the monomials. The distributive property in the factoring direction provides some insight to why we can add the coefficients of monomials that have like terms.

S**Less Steps Solution:**

$$13x \quad \text{OOA(2.1)}$$

**Example 7.5 – id:20141027-075159.**

Simplify $7 \text{ cm} + 8 \text{ cm}$

S**Solution:**

$$\begin{array}{ll} (7 + 8) \text{ cm} & \text{DPF(4.3a)} \\ 15 \text{ cm} & \text{OOA(2.1)} \end{array}$$

**R**

Remember, if a monomial does not have a coefficient factor, then it's implied that the coefficient factor is 1, the multiplicative identity, and consequently its not explicitly shown.

Example 7.6 – id:20141121-185558.

Simplify $x + 5x$

S**Solution:**

It can be useful when simplifying expressions to make the multiplicative identity (MId) factor explicit.

$$\begin{array}{ll} 1x + 5x & \text{MId(4.9a)} \\ (1 + 5)x & \text{DPF(4.3a)} \\ 6x & \text{OOA(2.1)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} 1x + 5x & \text{MId(4.9a)} \\ 6x & \text{OOA(2.1)} \end{array}$$

As one becomes more experienced, there is no reason to make the multiplicative identity coefficient explicit.

(S) _____

Less Steps Solution:

$$6x \quad \text{OOA(2.1)}$$

■

Example 7.7 – id:20141121-190857.

Simplify $8x - 6x$

(S) _____

Solution:

$$\begin{array}{ll} 8x + -6x & \text{DOS(4.12a)} \\ (8 + -6)x & \text{DPF(4.3a)} \\ 2x & \text{OOA(2.1)} \end{array}$$

■

(S) _____

Less Steps Solution:

$$\begin{array}{ll} 8x + -6x & \text{DOS(4.12a)} \\ 2x & \text{OOA(2.1)} \end{array}$$

■

Example 7.8 – id:20141121-193636.

Simplify $3x - 5x$

(S) _____

Solution:

$$\begin{array}{ll} 3x + -5x & \text{DOS(4.12a)} \\ (3 + -5)x & \text{DPF(4.3a)} \\ -2x & \text{OOA(2.1)} \\ -2x & \text{ONeg(3.1b)} \end{array}$$

■

(S) _____

Less Steps Solution:

$$\begin{array}{ll} 3x + -5x & \text{DOS(4.12a)} \\ -2x & \text{OOA(2.1)} \\ -2x & \text{ONeg(3.1b)} \end{array}$$

(S)

Less Steps Solution:

$$-2x \quad \text{OOA(2.1)}$$

■

Example 7.9 – id:20141106-150622.Simplify $13x - x$

(S)

Solution:

$$\begin{array}{ll} 13x - 1x & \text{MId(4.9a)} \\ 13x + -1x & \text{DOS(4.12a)} \\ (13 + -1)x & \text{DPF(4.3a)} \\ 12x & \text{OOA(2.1)} \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll} 13x + -x & \text{DOS(4.12a)} \\ 12x & \text{OOA(2.1)} \end{array}$$

■

It is possible for a univariate monomial to have more than two terms in its non-canonical form. The associative property of addition will be used to help simplify these expressions.

Example 7.10 – id:20141121-184652.Simplify the expression $3x + 7x + 8x$

(S)

$(3x + 7x) + 8x$	APA(4.2a)
$(3 + 7)x + 8x$	DPF(4.3a)
$10x + 8x$	OOA(2.1)
$(10 + 8)x$	DPF(4.3a)
$18x$	OOA(2.1)

S**Less Steps Solution:**

$(3x + 7x) + 8x$	APA(4.2a)
$10x + 8x$	OOA(2.1)
$18x$	OOA(2.1)

You might have noticed that this expression could be simplified in one step by adding the coefficient of the three monomials $3x$, $7x$ and $8x$, which have the like term x .

S**Less Steps Solution:**

$18x$	OOA(2.1)
-------	----------

Example 7.11 – id:20141106-152020.

Simplify $4x - 2x - x$

S**Solution:**

$4x - 2x - 1x$	MId(4.9a)
$4x + -2x + -1x$	DOS(4.12a)
$(4 + -2)x + -1x$	DPF(4.3a)
$2x + -1x$	OOA(2.1)
$(2 + -1)x$	DPF(4.3a)
$1x$	OOA(2.1)
x	MId(4.9b)

S

Less Steps Solution:

$$\begin{array}{rcl} 4x + -2x + -x & & \text{DOS(4.12a)} \\ x & & \text{OOA(2.1)} \end{array}$$

■

Example 7.12 – id:20141108-194431.Simplify $-3 \cdot 7x - 2x \cdot 4$

(S) _____

Solution:

$$\begin{array}{rcl} -3 \cdot 7x - 2x \cdot 4 & & \text{ONeg(3.1a)} \\ -3 \cdot 7x + -2x \cdot 4 & & \text{DOS(4.12a)} \\ -3 \cdot 7 \cdot x + -2 \cdot x \cdot 4 & & \text{JTC(3.5)} \\ -3 \cdot 7 \cdot x + -2 \cdot 4 \cdot x & & \text{CPM(4.6)} \\ (-3 \cdot 7) \cdot x + (-2 \cdot 4) \cdot x & & \text{APM(4.7a)} \\ -21 \cdot x + -8 \cdot x & & \text{OOM(2.3)} \\ -21x + -8x & & \text{CTJ(3.6)} \\ (-21 + -8)x & & \text{DPF(4.3a)} \\ -29x & & \text{OOA(2.1)} \\ -29x & & \text{ONeg(3.1b)} \end{array}$$

(S) _____

Less Steps Solution:

$$\begin{array}{rcl} -3 \cdot 7x + -2x \cdot 4 & & \text{DOS(4.12a)} \\ -3 \cdot 7 \cdot x + -2 \cdot 4 \cdot x & & \text{CPM(4.6)} \\ -21x + -8x & & \text{OOM(2.3)} \\ -29x & & \text{OOA(2.1)} \end{array}$$

■

Example 7.13 – id:20141108-194156.Simplify $3 \cdot 5x + 3x \cdot 4$

(S) _____

Solution:

$3 \cdot 5 \cdot x + 3 \cdot x \cdot 4$	JTC(3.5)
$3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x$	CPM(4.6)
$(3 \cdot 5) \cdot x + (3 \cdot 4) \cdot x$	APM(4.7a)
$15 \cdot x + 12 \cdot x$	OOM(2.3)
$15x + 12x$	CTJ(3.6)
$(15 + 12)x$	DPF(4.3a)
$27x$	OOA(2.1)

(S)

Less Steps Solution:

$3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x$	CPM(4.6)
$15x + 12x$	OOM(2.3)
$27x$	OOA(2.1)

Example 7.14 – id:20141108-173613.Simplify $8x \cdot 5$

(S)

Solution:

$8 \cdot x \cdot 5$	JTC(3.5)
$8 \cdot 5 \cdot x$	CPM(4.6)
$(8 \cdot 5) \cdot x$	APM(4.7a)
$40 \cdot x$	OOM(2.3)
$40x$	CTJ(3.6)

7.3 Simplifying Degree 2 Univariate Monomials**Example 7.15 – id:20141120-202842.**Express $1x^2$ in canonical form.

(S)

Solution:

$$x^2 \quad \text{MId(4.9b)}$$

■

Example 7.16 – id:20141106-151138.Simplify $4x^2 + 12x^2$

(S) _____

Solution:

$$\begin{aligned} (4 + 12)x^2 & \quad \text{DPF(4.3a)} \\ 16x^2 & \quad \text{OOA(2.1)} \end{aligned}$$

(S) _____

Less Steps Solution:

$$16x^2 \quad \text{OOA(2.1)}$$

■

Example 7.17 – id:20141106-154547.Simplify $x^2 - x + x^2 + x$

(S) _____

Solution:

$$\begin{aligned} 1x^2 - 1x + 1x^2 + 1x & \quad \text{MId(4.9a)} \\ 1x^2 + -1x + 1x^2 + 1x & \quad \text{DOS(4.12a)} \\ 1x^2 + 1x^2 + -1x + 1x & \quad \text{CPA(4.1)} \\ (1 + 1)x^2 + (-1 + 1)x & \quad \text{DPF(4.3a)} \\ 2x^2 + 0x & \quad \text{OOA(2.1)} \\ 2x^2 & \quad \text{MId(4.9b)} \end{aligned}$$

(S) _____

■

Less Steps Solution:

$$x^2 + \neg x + x^2 + x$$

DOS(4.12a)

$$x^2 + x^2 + \neg x + x$$

CPA(4.1)

$$2x^2$$

OOA(2.1)

Example 7.18 – id:20141108-194709.Simplify $-2x^4x - x \cdot -x^3$

(S)

Solution:

$$-2x^4x - 1x \cdot 1x^3$$

MId(4.9a)

$$\neg 2x^4x - 1x \cdot 1x^3$$

ONeg(3.1a)

$$\neg 2 \cdot x \cdot 4 \cdot x + \neg 1 \cdot x \cdot 1 \cdot x \cdot 3$$

JTC(3.5)

$$\neg 2 \cdot 4 \cdot x \cdot x + \neg 1 \cdot \neg 1 \cdot 3 \cdot x \cdot x$$

CPM(4.6)

$$\neg 2 \cdot 4 \cdot x^2 + \neg 1 \cdot \neg 1 \cdot 3 \cdot x^2$$

PrCBPo(5.3a)

$$(\neg 2 \cdot 4) \cdot x^2 + (\neg 1 \cdot \neg 1 \cdot 3)x^2$$

APM(4.7a)

$$\neg 8x^2 + 3x^2$$

OOM(2.3)

$$(\neg 8 + 3)x^2$$

DPF(4.3a)

$$\neg 5x^2$$

OOA(2.1)

$$-5x^2$$

ONeg(3.1b)

(S)

Less Steps Solution:

$$-2x^4x + \neg x \cdot -x^3$$

DOS(4.12a)

$$\neg 2 \cdot 4 \cdot x \cdot x + 3 \cdot \neg x \cdot \neg x$$

CPM(4.6)

$$\neg 2 \cdot 4 \cdot x^2 + 3 \cdot x^2$$

PrCBPo(5.3a)

$$\neg 8x^2 + 3x^2$$

OOM(2.3)

$$-5x^2$$

OOA(2.1)

Example 7.19 – id:20141108-191616.Simplify $-5x \cdot 4x$

(S)

Solution:

$-5x \cdot 4x$	ONeg(3.1a)
$-5 \cdot x \cdot 4 \cdot x$	JTC(3.5)
$-5 \cdot 4 \cdot x \cdot x$	CPM(4.6)
$-5 \cdot 4 \cdot x^2$	PrCBPo(5.3a)
$(-5 \cdot 4) \cdot x^2$	APM(4.7a)
$-20 \cdot x^2$	OOM(2.3)
$-20x^2$	CTJ(3.6)
$-20x^2$	ONeg(3.1b)

(S)

Less Steps Solution:

$-5 \cdot 4 \cdot x \cdot x$	CPM(4.6)
$-5 \cdot 4 \cdot x^2$	PrCBPo(5.3a)
$-20x^2$	OOM(2.3)

7.4 Simplifying Degree 3 Univariate Monomials

7.5 Simplifying Degree 1 Univariate Binomials

Example 7.20 – id:20141109-090809.

Simplify by expanding $5(x + 4)$

(S)

Solution:

$5(1x + 4)$	MId(4.9a)
$5 \cdot 1x + 5 \cdot 5$	DPF(4.3a)
$5 \cdot 1 \cdot x + 5 \cdot 5$	JTC(3.5)
$5 \cdot x + 25$	OOM(2.3)
$5x + 25$	CTJ(3.6)

(S)

Less Steps Solution:

$5x + 20$	DPE(4.8a)
-----------	-----------

Example 7.21 – id:20141109-091015.

Simplify by expanding $5(3x - 9)$

(S)**Solution:**

$$\begin{array}{ll} 5(3x + \neg 9) & \text{DOS(4.12a)} \\ 5 \cdot 3x + 5 \cdot \neg 9 & \text{DPE(4.8a)} \\ 5 \cdot 3 \cdot x + 5 \cdot \neg 9 & \text{JTC(3.5)} \\ 15 \cdot x + \neg 45 & \text{OOM(2.3)} \\ 15x + \neg 45 & \text{CTJ(3.6)} \\ 15x - 45 & \text{DOS(4.12b)} \end{array}$$

(S)**Less Steps Solution:**

$$\begin{array}{ll} 5(3x + \neg 9) & \text{DOS(4.12a)} \\ 15x + \neg 40 & \text{DPE(4.8a)} \\ 15x - 40 & \text{DOS(4.12b)} \end{array}$$

■

Example 7.22 – id:20141109-092448.

Simplify by expanding $-(5x + 7)$

(S)**Solution:**

$$\begin{array}{ll} \neg 1(5x + 7) & \text{MId(4.9a)} \\ \neg 1 \cdot 5x + \neg 1 \cdot 7 & \text{DPE(4.8a)} \\ \neg 1 \cdot 5 \cdot x + \neg 1 \cdot 7 & \text{JTC(3.5)} \\ \neg 5 \cdot x + \neg 7 & \text{OOM(2.3)} \\ \neg 5x + \neg 7 & \text{CTJ(3.6)} \\ \neg 5x - 7 & \text{DOS(4.12b)} \\ \neg 5x - 7 & \text{ONeg(3.1b)} \end{array}$$

(S)

Less Steps Solution:

$$-5x - 7$$

DPE(4.8a)

■

Example 7.23 – id:20141109-092651.Simplify by expanding $-13(7x - 9)$

(S) _____

Solution:

$$-13(7x + -9) \quad \text{DOS(4.12a)}$$

$$-13 \cdot 7x + -13 \cdot -9 \quad \text{DPE(4.8a)}$$

$$-13 \cdot 7 \cdot x + -13 \cdot -9 \quad \text{JTC(3.5)}$$

$$-91 \cdot x + 117 \quad \text{OOM(2.3)}$$

$$-91x + 117 \quad \text{CTJ(3.6)}$$

$$-91x + 117 \quad \text{ONeg(3.1b)}$$

(S) _____

Less Steps Solution:

$$-13(7x + -9) \quad \text{DOS(4.12a)}$$

$$-91x + 117 \quad \text{DPE(4.8a)}$$

■

Example 7.24 – id:20141109-092910.Simplify by expanding $a(x + b)$, where $a, b \in \mathbb{Z}$

(S) _____

Solution:

$$a(1x + b) \quad \text{MId(4.9a)}$$

$$a \cdot 1x + a \cdot b \quad \text{DPE(4.8a)}$$

$$a \cdot 1 \cdot x + a \cdot b \quad \text{JTC(3.5)}$$

$$1 \cdot a \cdot x + a \cdot b \quad \text{CPM(4.6)}$$

$$1ax + ab \quad \text{JTC(3.5)}$$

$$ax + ab \quad \text{MId(4.9b)}$$

S**Less Steps Solution:**

$$ax + ab$$

DPE(4.8a)

■

Example 7.25 – id:20141109-093220.Simplify by expanding $5(x + 2) + 4$ **S****Solution:**

$$5(1x + 2) + 4$$

MId(4.9a)

$$5 \cdot 1x + 5 \cdot 2 + 4$$

DPE(4.8a)

$$5 \cdot 1 \cdot x + 5 \cdot 2 + 4$$

JTC(3.5)

$$5 \cdot x + 10 + 4$$

OOM(2.3)

$$5x + 10 + 4$$

CTJ(3.6)

$$5x + 14$$

OOA(2.1)

S**Less Steps Solution:**

$$5x + 10 + 4$$

DPE(4.8a)

$$5x + 14$$

OOA(2.1)

■

Example 7.26 – id:20141109-093419.Simplify by expanding $7x + 5(4x + 8)$ **S**

Solution:

$$\begin{array}{ll}
 7x + 5 \cdot 4x + 5 \cdot 8 & \text{DPE(4.8a)} \\
 7 \cdot x + 5 \cdot 4 \cdot x + 5 \cdot 8 & \text{JTC(3.5)} \\
 7 \cdot x + 20 \cdot x + 40 & \text{OOM(2.3)} \\
 7x + 20x + 40 & \text{CTJ(3.6)} \\
 (7 + 20)x + 40 & \text{DPF(4.3a)} \\
 27x + 40 & \text{OOA(2.1)}
 \end{array}$$

■

Example 7.27 – id:20141109-094928.Simplify by expanding $4(3x + 4) + x + 6$

(S)

Solution:

$$\begin{array}{ll}
 4(3x + 4) + 1x + 6 & \text{MId(4.9a)} \\
 4 \cdot 3x + 4 \cdot 4 + 1x + 6 & \text{DPE(4.8a)} \\
 4 \cdot 3 \cdot x + 4 \cdot 4 + 1 \cdot x + 6 & \text{JTC(3.5)} \\
 12 \cdot x + 16 + 1 \cdot x + 6 & \text{OOM(2.3)} \\
 12x + 16 + 1x + 6 & \text{CTJ(3.6)} \\
 12 + 1x + 16 + 6 & \text{CPA(4.1)} \\
 (12 + 1)x + 16 + 6 & \text{DPF(4.3a)} \\
 13x + 22 & \text{OOA(2.1)}
 \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll}
 12x + 16 + x + 6 & \text{DPE(4.8a)} \\
 12x + x + 16 + 6 & \text{CPA(4.1)} \\
 13x + 22 & \text{OOA(2.1)}
 \end{array}$$

■

Example 7.28 – id:20141109-095151.Simplify by expanding $5(x - 4) + 3x - 5$

(S)

Solution:

$5(1x - 4) + 3x - 5$	MId(4.9a)
$5(1x + \neg 4) + 3x + \neg 5$	DOS(4.12a)
$5 \cdot 1x + 5 \cdot \neg 4 + 3x + \neg 5$	DPE(4.8a)
$5 \cdot 1 \cdot x + 5 \cdot \neg 4 + 3 \cdot x + \neg 5$	JTC(3.5)
$5 \cdot x + \neg 20 + 3 \cdot x + \neg 5$	OOM(2.3)
$5x + \neg 20 + 3x + \neg 5$	JTC(3.5)
$5x + 3x + \neg 20 + \neg 5$	CPA(4.1)
$(5 + 3)x + \neg 20 + \neg 5$	DPF(4.3a)
$8x + \neg 25$	OOA(2.1)
$8x - 25$	DOS(4.12b)

S**Less Steps Solution:**

$5(x + \neg 4) + 3x + \neg 5$	DOS(4.12a)
$5x + \neg 20 + 3x + \neg 5$	DPE(4.8a)
$5x + 3x + \neg 20 + \neg 5$	CPA(4.1)
$8x + \neg 25$	OOA(2.1)
$8x - 25$	DOS(4.12b)

Example 7.29 – id:20141109-095536.Simplify by expanding $8x - 5 - 4(x - 3)$ **S****Solution:**

$8x - 5 - 4(1x - 3)$	MId(4.9a)
$8x + \neg 5 + \neg 4(1x + \neg 3)$	DOS(4.12a)
$8x + \neg 5 + \neg 4 \cdot 1x + \neg 4 \cdot \neg 3$	DPE(4.8a)
$8 \cdot x + \neg 5 + \neg 4 \cdot 1 \cdot x + \neg 4 \cdot \neg 3$	JTC(3.5)
$8 \cdot x + \neg 5 + \neg 4 \cdot x + 12$	OOM(2.3)
$8x + \neg 5 + \neg 4x + 12$	CTJ(3.6)
$8x + \neg 4x + \neg 5 + 12$	CPA(4.1)
$(8 + \neg 4)x + \neg 5 + 12$	DPF(4.3a)
$4x + 7$	OOA(2.1)

S**Less Steps Solution:**

$$\begin{array}{ll} 8x + -5 + -4(x + -3) & \text{DOS(4.12a)} \\ 8x + -5 + -4x + 12 & \text{DPE(4.8a)} \\ 8x + -4x + -5 + 12 & \text{CPA(4.1)} \\ 4x + 7 & \text{OOA(2.1)} \end{array}$$

■

Example 7.30 – id:20141109-095842.Simplify by expanding $5(x + 3) + 3(x + 2)$ **S****Solution:**

$$\begin{array}{ll} 5 \cdot x + 5 \cdot 3 + 3 \cdot x + 3 \cdot 2 & \text{DPE(4.8a)} \\ 5 \cdot x + 15 + 3 \cdot x + 6 & \text{OOM(2.3)} \\ 5x + 15 + 3x + 6 & \text{CTJ(3.6)} \\ 5x + 3x + 15 + 6 & \text{CPA(4.1)} \\ (3 + 5)x + 15 + 6 & \text{DPF(4.3a)} \\ 8x + 21 & \text{OOA(2.1)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} 5x + 15 + 3x + 6 & \text{DPE(4.8a)} \\ 5x + 3x + 15 + 6 & \text{CPA(4.1)} \\ 8x + 21 & \text{OOA(2.1)} \end{array}$$

■

7.6 Simplifying Degree 2 Univariate Binomials**Example 7.31 – id:20141106-152339.**Simplify $3x^2 + 2x + 5x^2 + 4x$ **S**

Solution:

$$\begin{array}{ll} 3x^2 + 5x^2 + 2x + 4x & \text{CPA(4.1)} \\ (3+5)x^2 + (2+4)x & \text{DPF(4.3a)} \\ 8x^2 + 6x & \text{OOA(2.1)} \end{array}$$

If needed we could continue and express it in the simplified factored form using the distributive property

$$(4x + 3)2x \quad \text{DPF(4.3a)}$$

(S) _____**Less Steps Solution:**

$$\begin{array}{ll} 3x^2 + 5x^2 + 2x + 4x & \text{CPA(4.1)} \\ 8x^2 + 6x & \text{OOA(2.1)} \end{array}$$

■

Example 7.32 – id:20141107-121834.

$$\text{Simplify } (\sqrt{9 - x^2})^2$$

(S) _____**Solution:**

$$\begin{array}{ll} (\sqrt{9 - 1x^2})^2 & \text{MId(4.9a)} \\ (\sqrt{9 + -1x^2})^2 & \text{DOS(4.12a)} \\ [(9 + -x^2)^{\frac{1}{2}}]^2 & \text{RTPo(3.12)} \\ 9 + -1x^2 & \text{PoPo(5.1a)} \\ -1x^2 + 9 & \text{CPA(4.1)} \\ -x^2 + 9 & \text{MId(4.9a)} \\ -x^2 + 9 & \text{ONeg(3.1b)} \end{array}$$

(S) _____**Less Steps Solution:**

$$9 - x^2 \quad \text{PoPo(5.1a)}$$

It might be easier to view this using a variable substitution for the radicand, $9 - x^2$. Let $k = 9 + \neg 1x^2$.

$(\sqrt{k})^2$	MId(4.9a)
$(\sqrt{k})^2$	DOS(4.12a)
$\left[(k)^{\frac{1}{2}}\right]^2$	RTPo(3.12)
k	PoPo(5.1a)
$9 + \neg 1x^2$	CPA(4.1)
$\neg 1x^2 + 9$	CPA(4.1)
$\neg x^2 + 9$	MId(4.9a)
$-x^2 + 9$	ONeg(3.1b)

(D) _____

Dependencies:example ??-20141105-144223

**Example 7.33 – id:20141209-145211.**Simplify $2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0$

(S) _____

Solution:

$2x \cdot 2x + 2x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0$	DPE(4.8a)
$2 \cdot x \cdot 2 \cdot x + 2 \cdot x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0$	JTC(3.5)
$2 \cdot 2 \cdot x \cdot x + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0$	CPM(4.6)
$2 \cdot 2 \cdot x^2 + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0$	PrCBPo(5.3a)
$4 \cdot x^2 + 8 \cdot x + 2 \cdot x^2 + 0$	OOM(2.3)
$4x^2 + 8x + 2x^2 + 0$	CTJ(3.6)
$4x^2 + 2x^2 + 8x$	APA(4.2a)
$(4 + 2)x^2 + 8x$	DPF(4.3b)
$6x^2 + 8x$	OOA(2.1)

(D) _____

Dependencies:example 13.7-20141209-144203



7.7 Simplifying Degree 2 Univariate Trinomials

Example 7.34 – id:20141109-133008.

Simplify by expanding $(x + 5)(x - 8)$

(S)

Solution:

$$\begin{aligned}
 & (1x + 5)(1x - 8) && \text{MId(4.9a)} \\
 & (1x + 5)(1x + \neg 8) && \text{DOS(4.12a)} \\
 & 1x(1x + \neg 8) + 5(1x + \neg 8) && \text{DPE(4.8b)} \\
 & 1x \cdot 1x + 1x \cdot \neg 8 + 5 \cdot 1x + 5 \cdot \neg 8 && \text{DPE(4.8a)} \\
 & 1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot \neg 8 + 5 \cdot 1 \cdot x + 5 \cdot \neg 8 && \text{JTC(3.5)} \\
 & 1 \cdot 1 \cdot x \cdot x + \neg 8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + \neg 8 \cdot 5 && \text{CPM(4.6)} \\
 & 1 \cdot 1 \cdot x^2 + \neg 8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + \neg 8 \cdot 5 && \text{PrCBPo(5.3a)} \\
 & 1 \cdot x^2 + \neg 8 \cdot x + 5 \cdot x + \neg 40 && \text{OOM(2.3)} \\
 & 1x^2 + \neg 8x + 5x + \neg 40 && \text{CTJ(3.6)} \\
 & 1x^2 + \neg 3x + \neg 40 && \text{OOA(2.1)} \\
 & 1x^2 - 3x - 40 && \text{DOS(4.12b)} \\
 & x^2 - 3x - 40 && \text{MId(4.9a)}
 \end{aligned}$$

(S)

Less Steps Solution:

$$\begin{aligned}
 & (x + 5)(x + \neg 8) && \text{DOS(4.12a)} \\
 & x(x + \neg 8) + 5(x + \neg 8) && \text{DPE(4.8b)} \\
 & x^2 + \neg 8x + 5x + \neg 40 && \text{DPE(4.8a)} \\
 & x^2 - 3x - 40 && \text{OOA(2.1)}
 \end{aligned}$$

Example 7.35 – id:20141109-133316.

Simplify by expanding $(x + a)(x + b)$, where $a, b \in \mathbb{Z}$

(S)

Solution:

$(1x + a)(1x + b)$	MId(4.9a)
$1x(1x + b) + a(1x + b)$	DPE(4.8b)
$1x \cdot 1x + 1x \cdot b + a \cdot 1x + a \cdot b$	DPE(4.8a)
$1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot b + a \cdot 1 \cdot x + a \cdot b$	JTC(3.5)
$1 \cdot 1 \cdot x \cdot x + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b$	CPM(4.6)
$1 \cdot 1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b$	PrCBPo(5.3a)
$1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b$	OOM(2.3)
$1x^2 + 1bx + 1ax + ab$	CTJ(3.6)
$1x^2 + (1b + 1a)x + ab$	DPF(4.3a)
$x^2 + (b + a)x + ab$	MId(4.9b)

S

Less Steps Solution:

$x(x + b) + a(x + b)$	DPE(4.8b)
$x^2 + (b + a)x + ab$	DPE(4.8a)

Example 7.36 – id:20141109-140659.

Simplify by expanding $(2x + 3)(5x + 13)$

S

Solution:

$2x(5x + 13) + 3(5x + 13)$	DPE(4.8b)
$2x \cdot 5x + 2x \cdot 13 + 3 \cdot 5x + 3 \cdot 13$	DPE(4.8a)
$2 \cdot x \cdot 5 \cdot x + 2 \cdot x \cdot 13 + 3 \cdot 5 \cdot x + 3 \cdot 13$	JTC(3.5)
$2 \cdot 5 \cdot x \cdot x + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13$	CPM(4.6)
$2 \cdot 5 \cdot x^2 + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13$	PrCBPo(5.3a)
$10 \cdot x^2 + 26 \cdot x + 16 \cdot x + 39$	OOM(2.3)
$10x^2 + 26x + 15x + 39$	CTJ(3.6)
$10x^2 + 41x + 39$	OOA(2.1)

S

Less Steps Solution:

$$\begin{array}{ll} 2x(5x + 13) + 3(5x + 13) & \text{DPE(4.8b)} \\ 10x^2 + 26x + 15x + 39 & \text{DPE(4.8a)} \\ 10x^2 + 41x + 39 & \text{OOA(2.1)} \end{array}$$

Example 7.37 – id:20141109-141019.

Simplify by expanding $(-3x - 5)(7x + 8)$

(S)**Solution:**

$$\begin{array}{ll} (-3x - 5)(7x + 8) & \text{ONeg(3.1a)} \\ (-3x + -5)(7x + 8) & \text{DOS(4.12a)} \\ -3x(7x + 8) + -5(7x + 8) & \text{DPE(4.8b)} \\ -3x \cdot 7x + -3x \cdot 8 + -5 \cdot 7x + -5 \cdot 8 & \text{DPE(4.8a)} \\ -3 \cdot x \cdot 7 \cdot x + -3 \cdot x \cdot 8 + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{JTC(3.5)} \\ -3 \cdot 7 \cdot x \cdot x + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{CPM(4.6)} \\ -3 \cdot 7 \cdot x^2 + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 & \text{PrCBPo(5.3a)} \\ -21 \cdot x^2 + -24 \cdot x + -35 \cdot x + -40 & \text{OOM(2.3)} \\ -21x^2 + -24x + -35x + -40 & \text{CTJ(3.6)} \\ -21x^2 + -59x + -40 & \text{OOA(2.1)} \\ -21x^2 - 59x - 40 & \text{DOS(4.12b)} \\ -21x^2 - 59x - 40 & \text{ONeg(3.1b)} \end{array}$$

(S)**Less Steps Solution:**

$$\begin{array}{ll} (-3x + -5)(7x + 8) & \text{DOS(4.12a)} \\ -3x(7x + 8) + -5(7x + 8) & \text{DPE(4.8b)} \\ -21x^2 + -24x + -35x + -40 & \text{CTJ(4.8a)} \\ -21x^2 - 59x - 40 & \text{OOA(2.1)} \end{array}$$

Example 7.38 – id:20141109-141347.

Simplify by expanding $(ax + b)(cx + d)$, where $a, b, c, d \in \mathbb{Z}$

S**Solution:**

$$\begin{aligned} ax(cx + d) + b(cx + d) & \quad \text{DPE(4.8b)} \\ ax \cdot cx + ax \cdot d + b \cdot cx + b \cdot d & \quad \text{DPE(4.8a)} \\ a \cdot x \cdot c \cdot x + a \cdot x \cdot d + b \cdot c \cdot x + b \cdot d & \quad \text{JTC(3.5)} \\ a \cdot c \cdot x \cdot x + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d & \quad \text{CPM(4.6)} \\ a \cdot c \cdot x^2 + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d & \quad \text{PrCBPo(5.3a)} \\ acx^2 + adx + bcx + bd & \quad \text{CTJ(3.6)} \\ acx^2 + (ad + bc)x + bd & \quad \text{DPF(4.3a)} \end{aligned}$$

S**Less Steps Solution:**

$$\begin{aligned} ax(cx + d) + b(cx + d) & \quad \text{DPE(4.8b)} \\ acx^2 + (ad + bc)x + bd & \quad \text{DPE(4.8a)} \end{aligned}$$

■

Example 7.39 – id:20141105-161225.

Simplify $\left(2 - \frac{x}{2}\right)^2$ by expanding.

S

Solution:

$$\begin{aligned}
 & \left(2 - \frac{1x}{2}\right)^2 && \text{MId(4.9a)} \\
 & \left(2 + -\frac{1x}{2}\right)^2 && \text{DOS(4.12a)} \\
 & \left(2 + -\frac{1x}{2}\right) \left(2 + -\frac{1x}{2}\right) && \text{PoTF(3.9)} \\
 & 2 \left(2 + -\frac{1x}{2}\right) + -\frac{1x}{2} \left(2 + -\frac{1x}{2}\right) && \text{DPE(4.8b)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1x}{2} + -\frac{1x}{2} \cdot 2 + -\frac{1x}{2} \cdot -\frac{1x}{2} && \text{DPE(4.8a)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot x \cdot 2 + -\frac{1}{2} \cdot x \cdot -\frac{1}{2} \cdot x && \text{JTC(3.5)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x \cdot x && \text{CPM(4.6)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x^2 && \text{PrCBPo(5.3a)} \\
 & 4 + -1 \cdot x + -1 \cdot x + \frac{1}{4} \cdot x^2 && \text{OOM(2.3)} \\
 & 4 + -1x + -1x + \frac{1}{4}x^2 && \text{CTJ(3.6)} \\
 & \frac{1}{4}x^2 + -1x + -1x + 4 && \text{CPA(4.1)} \\
 & \frac{1}{4}x^2 + -2x + 4 && \text{OOA(2.1)} \\
 & \frac{1}{4}x^2 - 2x + 4 && \text{DOS(4.12b)}
 \end{aligned}$$

(S)**Less Steps Solution:**

$$\begin{aligned}
 & 7x + 20x + 40 && \text{DPE(4.8a)} \\
 & 27x + 40 && \text{OOA(2.1)}
 \end{aligned}$$

7.8 Simplifying Degree 1 Univariate Polynomials

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + \underbrace{C_1 x^1}_{C_0 x^0} + \underbrace{C_0}_{C_0 x^0}$$

Degree 1 univariate polynomial expressions can be expressed with at most two different terms and consequently this expression in its canonical form has at most two monomial summands – called a binomial.

7.9 Simplifying Degree 2 Univariate Polynomials

7.10 Simplifying Degree n Univariate Polynomials

Definition 7.10.1 – Univariate Polynomial Expression.

$$\sum_{k=0}^n C_k x^k = C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0} \quad (7.1)$$

A univariate polynomial in an indeterminate x is an expression made up of one or more summands of the form $C_k x^k$, which are called monomials. The first factor of each monomial, C_k , is a numerical factor called the **coefficient** where $C_k \in$. The second factor of each monomial, x^k , is an indeterminate raised to a non-negative integer power i .

Polynomial [3]

Definition 7.10.2 – Degree of the Univariate Polynomial.

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0}$$

The degree of the univariate polynomial is determined by the monomial with the largest degree of the indeterminate.

8. Simplifying Univariate Monomials

8.1

9. Solving Linear Equations

9.1 Power Inverse

Example 9.1 – id:20141206-102142.

Solve the equation $x + a = b$ for x

(S) _____

Solution:

$$\begin{array}{ll} [x + a] + \neg a = [b] + \neg a & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x + (a + \neg a) = b + \neg a & \text{APA(4.2b)} \\ x + 0 = b + \neg a & \text{OOA(2.1)} \\ x = b + \neg a & \text{AId(4.4b)} \\ x = b - a & \text{DOS(4.12b)} \end{array}$$

■

Example 9.2 – id:20141111-222931.

Solve the equations $x + 8 = 0$

(S) _____

Solution:

$$\begin{array}{ll} [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x + (8 + \neg 8) = 0 + \neg 8 & \text{APA(4.2b)} \\ x + 0 = \neg 8 & \text{OOA(2.1)} \\ x = \neg 8 & \text{AId(4.4b)} \\ x = -8 & \text{ONeg(3.1b)} \end{array}$$

(S) _____

Less Steps Solution:

$$\begin{array}{ll} [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x = -8 & \text{OOA(2.1)} \end{array}$$

D

Dependencies:
example ??-20141111-190212

Example 9.3 – id:20141206-101632.

Solve the equation $x + 4 = 7$

S

Solution:

$$\begin{array}{ll} [x + 4] + \neg 4 = [7] + \neg 4 & \text{SPE(4.15) + AI(4.5a)} \\ x + (4 + \neg 4) = 7 + \neg 4 & \text{APA(4.2b)} \\ x + 0 = 3 & \text{OOA(2.1)} \\ x = 3 & \text{AId(4.4b)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} [x] + 4 + \neg 4 = [7] + \neg 4 & \text{SPE(4.15) + AI(4.5a)} \\ x = 3 & \text{OOA(2.1)} \end{array}$$

Example 9.4 – id:20141206-101107.

Solve the equation $x - 8 = 15$ for x

S

Solution:

$$\begin{array}{ll} x + \neg 8 = 15 & \text{DOS(4.12a)} \\ [x + \neg 8] + 8 = [15] + 8 & \text{SPE(4.15) + AI(4.5a)} \\ x + (\neg 8 + 8) = 15 + 8 & \text{APA(4.2b)} \\ x + 0 = 23 & \text{OOA(2.1)} \\ x = 23 & \text{AId(4.4b)} \end{array}$$

S

Less Steps Solution:

$$\begin{aligned} [x + \cancel{-8}] + 8 &= [15] + 8 && \text{SPE(4.15) + AI(4.5a)} \\ x &= 23 && \text{OOA(2.1)} \end{aligned}$$

■

Example 9.5 – id:20141206-102404.

Solve the equation $5x = 9$ for x .

(S)

Solution:

$$\begin{aligned} \frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{5} \cdot [5 \cdot x] &= \frac{1}{5} \cdot 9 && \text{JTC(3.5)} \\ \left(\frac{1}{5} \cdot 5\right) \cdot x &= \frac{1}{5} \cdot 9 && \text{APM(4.7b)} \\ 1 \cdot x &= \frac{9}{5} && \text{OOM(2.3)} \\ x &= \frac{9}{5} && \text{MId(4.9b)} \end{aligned}$$

(S)

Less Steps Solution:

$$\begin{aligned} \frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.15) + MI(4.10a)} \\ x &= \frac{9}{5} && \text{OOM(2.3)} \end{aligned}$$

■

Example 9.6 – id:20141206-104404.

Solve the equation $ax = b$ for x .

(S)

Solution:

$$\begin{aligned}\frac{1}{a}[\mathbf{ax}] &= \frac{1}{a}[\mathbf{b}] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{a} \cdot (a \cdot x) &= \frac{1}{a} \cdot b && \text{JTC(3.5)} \\ \left(\frac{1}{a} \cdot a\right) \cdot x &= \frac{1}{a} \cdot b && \text{APM(4.7b)} \\ 1 \cdot x &= \frac{b}{a} && \text{OOM(2.3)} \\ x &= \frac{b}{a} && \text{MId(4.9b)}\end{aligned}$$

■

Example 9.7 – id:20141206-102723.Solve the equation $-2x = 7$ for x

(S)

Solution:

$$\begin{aligned}-2x &= 7 && \text{ONeg(3.1a)} \\ -\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.15) + MI(4.10b)} \\ -\frac{1}{2} \cdot (-2 \cdot x) &= -\frac{1}{2} \cdot 7 && \text{JTC(3.5)} \\ \left(-\frac{1}{2} \cdot -2\right) \cdot x &= -\frac{1}{2} \cdot 7 && \text{APM(4.7b)} \\ 1 \cdot x &= -\frac{7}{2} && \text{OOM(2.3)} \\ 1 \cdot x &= -\frac{7}{2} && \text{ONeg(3.1b)} \\ x &= -\frac{7}{2} && \text{MId(4.9b)}\end{aligned}$$

(S)

Less Steps Solution:

$$\begin{aligned}-\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.15) + MI(4.10b)} \\ x &= -\frac{7}{2} && \text{OOM(2.3)}\end{aligned}$$

■

Example 9.8 – id:20141111-215726.

Solve the equation $2x + 5 = 0$ for x

(S) _____

Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.15) + AI(4.5a)} \\
 2x + (5 + \neg 5) &= 0 + \neg 5 && \text{APA(4.2a)} \\
 2x + 0 &= \neg 5 && \text{OOA(2.1)} \\
 2x &= \neg 5 && \text{AId(4.4a)} \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.15) + MI(4.10a)} \\
 \frac{1}{2} \cdot 2 \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{JTC(3.5)} \\
 \left(\frac{1}{2} \cdot 2\right) \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{APM(4.7a)} \\
 1 \cdot x &= \frac{\neg 5}{2} && \text{OOM(2.3)} \\
 1x &= -\frac{5}{2} && \text{ONeg(3.1b)} \\
 x &= -\frac{5}{2} && \text{MId(4.9b)}
 \end{aligned}$$

(S) _____

Less Steps Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.15) + AI(4.5a)} \quad (9.1) \\
 2x &= \neg 5 && \text{OOA(2.1)} \quad (9.2) \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.15) + MI(4.10a)} \quad (9.3) \\
 x &= -\frac{5}{2} && \text{OOM(2.3)} \quad (9.4)
 \end{aligned}$$

(D) _____

Dependencies:

example ??-20141111-192213

■

Example 9.9 – id:20151015-104754.

Solve the equation $2ax + b = 0$ for x .

(S) _____

Solution:

$$\begin{aligned}
 [2ax + b] + \neg b &= [0] + \neg b && \text{AI(4.5a), SPE(4.15)} \\
 2ax + (b + \neg b) &= 0 + \neg b && \text{APA(4.2b)} \\
 2ax + 0 &= 0 + \neg b && \text{OOA(2.1)} \\
 2ax = \neg b &&& \text{AId(4.4a)} \\
 \frac{1}{2a} [2ax] &= \frac{1}{2a} [\neg b] && \text{MI(4.5a), SPE(4.15)} \\
 \frac{1}{2a} \cdot (2ax) &= \frac{1}{2a} \cdot (\neg b) && \text{JTC(3.5)} \\
 \left(\frac{1}{2a} \cdot 2a\right) \cdot x &= \frac{1}{2a} \cdot \neg b && \text{APA(4.2a)} \\
 \left(\frac{1}{2a} \cdot \frac{2a}{1}\right) x &= \frac{1}{2a} \cdot \frac{\text{neg } b}{1} && \text{MId(4.9a)} \\
 \frac{2a}{2a} \cdot x &= \frac{\neg b}{2a} && \text{OOM(2.3)} \\
 1 \cdot x &= \frac{\neg b}{2a} x && = \frac{\neg b}{2a} && \text{MId(4.9b)} \\
 x &= -\frac{b}{2a} && \text{ONeg(3.1b)}
 \end{aligned}$$

(D)

Dependencies:example 16.1-20151008-110208

10. Solving Quadratic Equations

10.1 Power Inverse

Example 10.1 – id:20141107-131748.

Solve the equation $2 - x^2 = 0$ for x

(S) _____

Solution:

$$\begin{array}{ll} 2 - 1x^2 = 0 & \text{MId(4.9a)} \\ 2 + -1x^2 = 0 & \text{DOS(4.12a)} \\ [2 + -1x^2] + 1x^2 = [0] + 1x^2 & \text{SPE(4.15) + AI(4.5a)} \\ 2 + (-1x^2 + 1x^2) = 0 + 1x^2 & \text{APA(4.2a)} \\ 2 + 0 = 0 + 1x^2 & \text{OOA(2.1)} \\ 2 = 1x^2 & \text{AIId(4.4a)} \\ 2 = x^2 & \text{MId(4.9b)} \\ \pm [2]^{\frac{1}{2}} = [x^2]^{\frac{1}{2}} & \\ \pm 2^{\frac{1}{2}} = x & \text{PoPo(5.1a)} \\ \pm \sqrt{2} = x & \text{PoTR(??)} \\ x = \pm \sqrt{2} & \text{SyPE(4.16a)} \end{array}$$

■

Example 10.2 – id:20151012-192313.

Solve the equation $6x^2 - 6x - 12 = 0$


(S) _____

Solution:

$$\begin{array}{ll} 6x^2 + -6x + -12 = 0 & \text{DOS(4.12a)} \\ 6(1x^2 - 1x + -2) = 0 & \text{DPF(4.3b)} \end{array}$$

■

Using the factor organizer,



Solving the linear equations using ZPr (4.11a)

$$x_1 + -2 = 0$$

$$[x_1 + -2] + 2 = [0] + -2$$

$$x_1 + (-2 + 2) = 0 + -2$$

$$x_1 + 0 = -2$$

$$x_1 = -2$$

$$x_1 = -2$$

Case I

AI(4.5a), SPE(4.15)

APA(4.2a)

OOA(2.1)

AId(4.4b)

ONeg(3.1b)

$$x_2 + 1 = 0$$

$$[x_2 + 1] + -1 = [0] + -1$$

$$x_2 + (1 + -1) = 0 + -1$$

$$x_2 + 0 = -1$$

$$x_2 = -1$$

$$x_2 = -1$$

Case II

AI(4.5a), SPE(4.15)

APA(4.2a)

OOA(2.1)

AId(4.4b)

ONeg(3.1b)

D

Dependencies:example 15.1-20151012-190708

■

10.2 Completing The Square

Completing the square is an algebraic algorithm used to find the solutions of quadratic equations of the form, $ax^2 + bx + c = 0$. Essentially, we want to manipulate this equation such that $x = \text{some value}$. To gain some understanding of how this algorithm works, we will consider each step individually. Let's begin with a quadratic equation in the general form: $ax^2 + bx + c = 0$.

Since we are trying to manipulate the equation $ax^2 + bx + c = 0$ such that $x = \text{some value}$, we first want the coefficient factor a to be equal to 1. This is done by multiplying both expressions by the multiplicative inverse of a (step 1) followed by simplifying both expressions.

$$\begin{aligned} \frac{1}{a}[ax^2 + bx + c] &= \frac{1}{a}[0] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{a} \cdot ax^2 + \frac{1}{a} \cdot bx + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{DPE(4.8a)} \\ \frac{1}{a} \cdot a \cdot x^2 + \frac{1}{a} \cdot b \cdot x + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{JTC(3.5)} \\ x^2 + \frac{b}{a} \cdot x + \frac{c}{a} &= 0 && \text{OOM(2.3)} \\ x^2 + \frac{b}{a}x + \frac{c}{a} &= 0 && \text{CTJ(3.6)} \end{aligned}$$

We now have three terms in the left hand expression where the first two terms have at least one variable factor, x . The problem is that the third term is a constant and we want $x = \text{some value}$. This next step is to get rid of the $\frac{c}{a}$ term, which can be done by using the additive inverse followed by simplifying both expressions.

$$\begin{aligned} \left[x^2 + \frac{b}{a}x + \frac{c}{a} \right] + -\frac{c}{a} &= [0] + -\frac{c}{a} && \text{SPE(4.15) + AI(4.5a)} \\ x^2 + \frac{b}{a}x + \frac{c}{a} + -\frac{c}{a} &= 0 + -\frac{c}{a} && \text{APA(4.2a)} \\ x^2 + \frac{b}{a}x &= -\frac{c}{a} && \text{OOA(2.1)} \end{aligned}$$

The next step is called completing the square. The idea is to add a *NeW* constant, k , to the left-hand expression, $x^2 + \frac{b}{a}x + k$, such that the quadratic expression can then be factored as two identical factors, $(x + m)(x + m) = (x + m)^2$, where $k = m \cdot m$. Notice that since we are adding a constant term, k , to the left-hand expression, then we must also add this constant, k , to the right-hand expression, $x^2 + \frac{b}{a}x + k = -\frac{c}{a} + k$. To determine the values of both m and k we should refer to the organization of the two factors that make up the product of a quadratic expression, $x^2 + \frac{b}{a}x + k = (x + m)^2$.

Since both factors of this new quadratic expression are the same, both terms that make up the middle term, must also be the same. We know that $mx + mx = \frac{b}{a}x$, so we should be able to determine the value of m from this equation. If we can determine the value of m , then we can determine the value of k .

$$\begin{aligned} \frac{b}{a}x &= mx + mx \\ &= 2mx && \text{OOA(2.1)} \end{aligned}$$

Solving for m ,

$$\begin{aligned}2mx &= \frac{b}{a}x \\2 \cdot m \cdot x &= \frac{b}{a} \cdot x && \text{JTC(3.5)} \\ \frac{1}{2} [2 \cdot m \cdot x] &= \frac{1}{2} \left[\frac{b}{a} \cdot x \right] && \text{SPE(4.15) + MI(4.10a)} \\m \cdot x &= \frac{b}{2a} \cdot x && \text{OOM(2.3)} \\[m \cdot x] \frac{1}{x} &= \left[\frac{b}{2a} \cdot x \right] \frac{1}{x} && \text{SPE(4.15) + MI(4.10a)} \\m &= \frac{b}{2a} && \text{OOM(2.3)}\end{aligned}$$

$\left[x^2 + \frac{b}{a}x \right] + \left(\frac{b}{2a} \right)^2 = \left[-\frac{c}{a} \right] + \left(\frac{b}{2a} \right)^2$	SPE(4.15)
$x^2 + \frac{b}{a}x + \left(\frac{b}{2a} \right)^2 = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	APA(4.2b)
$\left(x + \frac{b}{2a} \right) \left(x + \frac{b}{2a} \right) = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	DPF(4.3b)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	PoTF(3.9)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{(b)^2}{(2a)^2}$	PoQPo(5.5a)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{b^2}{2^2 a^2}$	PoPrPo
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{b^2}{4a^2}$	OOE
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} \cdot \frac{4a}{4a} + \frac{b^2}{4a^2}$	MId
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c \cdot 4 \cdot a}{a \cdot 4 \cdot a} + \frac{b^2}{4a^2}$	JTC
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4 \cdot a \cdot c}{4 \cdot a \cdot a} + \frac{b^2}{4a^2}$	CPM
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4 \cdot a \cdot c}{4 \cdot a^2} + \frac{b^2}{4a^2}$	PrCBPo
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4ac}{4a^2} + \frac{b^2}{4a^2}$	CTJ
$\left(x + \frac{b}{2a} \right)^2 = \frac{-4ac + b^2}{4a^2}$	CD
$\left(x + \frac{b}{2a} \right)^2 = \frac{b^2 - 4ac}{4a^2}$	CPA
$\left[\left(x + \frac{b}{2a} \right)^2 \right]^{\frac{1}{2}} = \pm \left[\frac{b^2 - 4ac}{4a^2} \right]^{\frac{1}{2}}$	PoI

$$x + \frac{b}{2a} = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}} \quad \text{PoPrPo}$$

$$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{\sqrt{4a^2}} \quad \text{PoQPo}$$

$$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{4^{\frac{1}{2}}a} \quad \text{PoPrPo}$$

$$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a} \quad \text{OOE}$$

$$\left[x + \frac{b}{2a} \right] + -\frac{b}{2a} = \left[\pm \frac{\sqrt{b^2 - 4ac}}{2a} \right] + -\frac{b}{2a} \quad \text{AI}$$

$$x + \frac{b}{2a} + -\frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a} + -\frac{b}{2a} \quad \text{APA}$$

$$x + \frac{b}{2a} + -\frac{b}{2a} = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a} \quad \text{CPA}$$


$$x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a} \quad \text{OOA}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{CD}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{DOS}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{ETR}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{ONeg}$$



Functions

11. Functions

Property 11.0.1 – Function Value Argument Substitution (FVAS).

$$f(a) \quad (11.1a)$$

Example 11.1 – id:20151012-201647.

Find $f(2)$ given $f(x) = 2x^3 - 3x^2 - 12x + 1$

(S) _____

Solution:

$$\begin{aligned} f(x) &= 2x^3 + -3x^2 + -12x + 1 && \text{DOS(4.12a)} \\ f(2) &= 2[2]^3 + -3[2]^2 + -12[2] + 1 && \text{SPE(4.15)} \\ f(2) &= 2(8) + -3(4) + -12(2) + 1 && \text{OOE(2.5)} \\ f(2) &= 16 + -12 + -24 + 1 && \text{OOM(2.3)} \\ f(2) &= -19 && \text{OOA(2.1)} \end{aligned}$$

(D) _____

Dependencies:example 15.1-20151012-190708

Example 11.2 – id:20151012-203549.

Find $f(-1)$ given $f(x) = 2x^3 - 3x^2 - 12x + 1$

(S) _____

Solution:

$$f(x) = 2x^3 + -3x^2 + -12x + 1 \quad \text{DOS(4.12a)}$$

$$f(-1) = 2[-1]^3 + -3[-1]^2 + -12[-1] + 1 \quad \text{SPE(4.15)}$$

$$f(-1) = 2(-1) + -3(1) + -12(-1) + 1 \quad \text{OOE(2.5)}$$

$$f(-1) = -2 + -3 + 12 + 1 \quad \text{OOM(2.3)}$$

$$f(-1) = 8 \quad \text{OOA(2.1)}$$

(D) _____

Dependencies:example 15.1-20151012-190708

11.1 Inverse Functions

11.2 Inverses

Property 11.2.1 – Cosine Inverse (ArcCos).

$$\cos^{-1}(\cos \theta) = \theta \quad (11.2a)$$

Property 11.2.2 – Sine Inverse (ArcSin).

$$\sin^{-1}(\sin \theta) = \theta \quad (11.3a)$$

Property 11.2.3 – Tangent Inverse (ArcTan).

$$\tan^{-1}(\tan \theta) = \theta \quad (11.4a)$$

Property 11.2.4 – Exponential Inverse (EI).

$$\log_a(a^x) = x \quad (11.5a)$$

Property 11.2.5 – Logarithmic Inverse (LI).

$$a^{\log_a x} = x \quad (11.6a)$$

Property 11.2.6 – Power Inverse (Pol).

$$(b^m)^{\frac{1}{m}} = b \quad (11.7a)$$

Differential Calculus

12	Derivative by First Principles	79
12.1	Limit of the Difference Quotient	
13	Derivative Rules	81
13.1	Derivative of a Monomial Functions	
13.2	Derivative of Polynomial Functions	
13.3	Derivative of a Quotient	
13.4	Derivative of a Rational Function	
14	Equations of Tangent & Secant Lines	91
14.1	Essential Questions	
14.2	Finding the Equation of the Tangent Line	
15	First Derivative Test	95
16	Curve Sketching	97
17	Second Derivative Test	99
18	Curve Sketching	101
	Bibliography	103
	Bibliography	103
	Websites	
	Articles	
	Index	105

12. Derivative by First Principles

12.1 Limit of the Difference Quotient

Definition 12.1.1 – Derivative. The derivative of a function $f(x)$ with respect to the variable x is defined as

$$f'(x) \equiv \lim_{\Delta x \rightarrow 0} \underbrace{\frac{f(x + \Delta x) - f(x)}{\Delta x}}_{\text{Difference Quotient}} \quad (12.1)$$




Figure 12.1: [2]

Example 12.1 – id:20141219-212546.

Differentiate the function $f(x) = 5$

(S)

Solution:

$$f(x) = 5x^0 \quad \text{PoID(4.13a)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5[x + \Delta x]^0 - 5[x]^0}{\Delta x} \quad \text{SPE(4.15)&DBFP(12.1)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5(1) - 5(1)}{\Delta x} \quad \text{PoID(4.13b)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} 0 \quad \text{OOM(2.3)}$$

$$f'(x) = 0$$

■

13. Derivative Rules

13.1 Derivative of a Monomial Functions

Example 13.1 – id:20141124-153017.

Differentiate $f(x) = -3$

(S) _____

Solution:

$$f'(x) = [-3]'$$

SPE(4.15)

$$f'(x) = 0$$

DC(6.1)

(D) _____

Dependencies:

example 13.3-20141124-152503

■

Example 13.2 – id:20141124-141850.

Differentiate $f(x) = x^2$

(S) _____

Solution:

$$f'(x) = [x^2]'$$

SPE(4.15)

$$f'(x) = 2x^{2-1}$$

DPo(6.5)

$$f'(x) = 2x^1$$

OOA(2.1)

$$f'(x) = 2x$$

MId(4.9b)

(S) _____

Less Steps Solution:

$$f'(x) = 2x$$

DPo(6.5)

(D) _____

Dependencies:

example 13.2-20141124-141850

example 13.3-20141124-152503

■

13.2 Derivative of Polynomial Functions**Example 13.3 – id:20141124-152503.**Differentiate $f(x) = x^2 - 3$

(S) _____

Solution:

$$\begin{aligned}
 f(x) &= x^2 + -3 && \text{DOS(4.12a)} \\
 f'(x) &= [x^2 + -3]' && \text{SPE(4.15)} \\
 f'(x) &= [x^2]' + [-3]' && \text{DS(6.7)} \\
 f'(x) &= [x^2]' + 0 && \text{DC(6.1)} \\
 f'(x) &= [x^2]' && \text{AId(4.4a)} \\
 f'(x) &= 2x && \text{DPo(6.5) goto 13.2}
 \end{aligned}$$

(S) _____

Less Steps Solution:

$$f(x) = 2x^2 \quad \text{DPo(6.5)&DC(6.5)}$$

(D) _____

Dependencies:

example 13.14-20141124-205219

■

Example 13.4 – id:20151015-155838.Differentiate $y = x^3 + 1$

(S) _____

Solution:

$$\begin{aligned}
 [y]' &= [x^3 + 1] && \text{SPE(4.15)} \\
 y' &= [x^3]' + [1]' && \text{DS(6.7)} \\
 y' &= 3x^2 + [1]' && \text{DPo(6.5)} \\
 y' &= 3x^2 + 0 && \text{DC(6.1)} \\
 y' &= 3x^2 && \text{MId(4.9b)}
 \end{aligned}$$

D

Dependencies:example 13.16-20151015-153507

Example 13.5 – id:20151011-195002.Differentiate $f(x) = 2x^2 + 3x + 7$ **S****Solution:**

$$\begin{aligned}
 [f'(x)]' &= [2x^2 + 3x + 7]' && \text{SPE(4.15)} \\
 f'(x) &= [2x^2]' + [3x]' + [7]' && \text{DS(6.7)} \\
 f'(x) &= 2[x^2]' + 3[x]' + [7]' && \text{DCM(6.3)} \\
 f'(x) &= 2[x^2]' + 3[x]' + 0 && \text{DC(6.1)} \\
 f'(x) &= 2[x^2]' + 3[x]' && \text{AId(4.4a)} \\
 f'(x) &= 2(2x) + 3 && \text{DPo(6.5)} \\
 f'(x) &= 4x + 3 && \text{OOM(2.3)}
 \end{aligned}$$

D

Dependencies:example 14.1-20151011-154209

Example 13.6 – id:20141128-151834.Differentiate $f(x) = 3x^2 - 6x + 4$ **S**

Solution:

$$\begin{aligned}
 f(x) &= 3x^2 + \neg 6x + 4 && \text{DOS(4.12a)} \\
 f'(x) &= [\mathbf{3x^2} + \mathbf{\neg 6x} + \mathbf{4}]' && \text{SPE(4.15)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' + [\mathbf{4}]' && \text{DS(6.7)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' + 0 && \text{DC(6.1)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' && \text{AId(4.4a)} \\
 f'(x) &= 3[\mathbf{x^2}]' + 6[\mathbf{x}]' && \text{DCM(6.3)} \\
 f'(x) &= 3(2x) + 6(1) && \text{DPo(6.5)} \\
 f'(x) &= 6x + 6 && \text{OOM(2.3)}
 \end{aligned}$$

(S)

$$f'(x) = 6x + 6 \quad \text{DS(6.7)}$$

Example 13.7 – id:20141209-144203.

Differentiate $f(x) = x^2(2x + 4)$

(S)

Solution:

$$\begin{aligned}
 f'(x) &= [\mathbf{x^2}(2x + 4)]' && \text{SPE(4.15)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2[\mathbf{2x} + \mathbf{4}]' && \text{DPr(6.11)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2[\mathbf{2x}] + [\mathbf{4}]' && \text{DS(6.7)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2 \cdot 2[\mathbf{x}] + [\mathbf{4}]' && \text{DCM(6.3)} \\
 f'(x) &= 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + [\mathbf{4}]' && \text{DPo(6.5)} \\
 f'(x) &= 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0 && \text{DC(6.1)} \\
 f'(x) &= 6x^2 + 8x && \text{simplify goto 7.33}
 \end{aligned}$$

Example 13.8 – id:20141209-142321.

Differentiate $f(x) = x^2 \cos(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [x^2 \cos(x)]' && \text{SPE(4.15)} \\ f'(x) &= [x^2]' \cos(x) + x^2[\cos(x)]' && \text{DPr(6.11)} \\ f'(x) &= 2x \cos(x) + x^2[\cos(x)]' && \text{DPo(6.5)} \\ f'(x) &= 2x \cos(x) + x^2(-1 \sin(x)) && \text{DCos(6.19)} \\ f'(x) &= 2x \cos(x) - x^2 \sin x && \text{OOM(2.3)} \end{aligned}$$

■

Example 13.9 – id:20150910-115935.

Differentiate $f(x) = \sin(x) \cos(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [\sin(x) \cos(x)]' && \text{SPE(4.15)} \\ f'(x) &= [\sin(x)]' \cos(x) + \sin(x)[\cos(x)]' && \text{DPr(6.11)} \\ f'(x) &= \cos(x) \cos(x) + \sin(x)[\cos(x)]' && \text{DSin(6.17)} \\ f'(x) &= \cos(x) \cos(x) + \sin(x)(-\sin(x)) && \text{DCos(6.19)} \\ f'(x) &= \cos^2(x) - \sin^2(x) && \text{simplify goto ??} \end{aligned}$$

■

Example 13.10 – id:20141209-151354.

Differentiate $f(x) = \sin(x) \sin(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [\sin(x) \sin(x)]' && \text{SPE(4.15)} \\ f'(x) &= [\sin(x)]' \sin(x) + \sin(x)[\sin(x)]' && \text{DPr(6.11)} \\ f'(x) &= \cos(x) \sin(x) + \sin(x) \cos(x) && \text{DSin(6.17)} \\ f'(x) &= \cos(x) \sin(x) + \cos(x) \sin(x) && \text{CPM(4.6)} \\ f'(x) &= 2 \cos(x) \sin(x) && \text{OOA(2.1)} \end{aligned}$$

■

Example 13.11 – id:20141124-203850.

Differentiate $y = \ln(3x)$

(S)

Solution:

After identifying that $y = \ln(3x)$ is a composite function, we let $u = 3x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(6.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = 3x$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = 3$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(6.16)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot \frac{du}{dx} \quad \text{DNL(6.31)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot 3 \quad \text{DPo(6.5)}$$

$$\frac{dy}{dx} = \frac{1}{3x} \cdot 3$$

$$\frac{dy}{dx} = \frac{3}{3x} \quad \text{OOM(2.3)}$$

$$\frac{dy}{dx} = \frac{1}{x}$$

(D)

Dependencies:

example 13.14-20141124-205219

Example 13.12 – id:20141128-160248.

Differentiate $y = \ln(3x^2 - 6x + 4)$

(S)

Solution: After identifying that $y = \ln(3x^2 - 6x + 4)$ is a composite function, we let $u = 3x^2 - 6x + 4$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(6.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = 3x^2 - 6x + 4$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = 6x - 6 \quad \text{goto 13.6}$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(6.16)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot \frac{du}{dx} \quad \text{DNL(6.31)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot (6x - 6) \quad \text{DPo(6.5)}$$

$$\frac{dy}{dx} = \frac{1}{3x^2 - 6x + 4} \cdot (6x - 6)$$

$$\frac{dy}{dx} = \frac{6x - 6}{3x^2 - 6x + 4} \quad \text{OOM(2.3)}$$

Example 13.13 – id:20141128-155506.

Differentiate $y = \ln(\cos x)$

(S)

Solution: After identifying that $y = \ln(\cos x)$ is a composite function, we let $u = \cos x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(6.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = \cos x$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = -\sin x$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\text{d}y}{\text{d}u} \cdot \frac{\text{d}u}{\text{d}x} && \text{DComp(6.16)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot \frac{\text{d}u}{\text{d}x} && \text{DNL(6.31)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot -\sin x && \text{DPo(6.5)} \\ \frac{dy}{dx} &= \frac{1}{\cos x} \cdot -\sin x \\ \frac{dy}{dx} &= \frac{-\sin x}{\cos x} && \text{OOM(2.3)} \\ \frac{dy}{dx} &= -\tan x\end{aligned}$$

■

Example 13.14 – id:20141124-205219.Differentiate $y = (x^2 - 1) \ln(3x)$

(S)

Solution:

$$\begin{aligned}y' &= [x^2 - 3]' \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' && \text{DPr(6.11)} \\ y' &= 2x \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' && \text{differentiate goto 13.3} \\ y' &= 2x \cdot \ln x + (x^2 - 3) \cdot \frac{1}{x} && \text{differentiate goto 13.11} \\ y' &= 2x \cdot \ln x + \frac{x^2 - 3}{x} && \text{OOM(2.3)} \\ y' &= 2x^2 \ln x + \frac{x^2 - 3}{x} && \text{JTC(3.5)} \\ y' &= \frac{2x^2 \ln x + (x^2 - 3)}{x} && \text{OOA(2.1)}\end{aligned}$$

■

13.3 Derivative of a Quotient**Example 13.15 – id:20151015-165037.**Differentiate $y = \frac{\sin x}{e^x}$

(S)

Solution:

$$\frac{d[y]}{dx} = \frac{d\left[\frac{\sin x}{e^x}\right]}{dx} \quad \text{SPE(4.15)}$$

$$\frac{dy}{dx} = \frac{\frac{d}{dx}[\sin x]e^x - \sin x \frac{d}{dx}[e^x]}{[e^x]^2} \quad \text{DQ(??)}$$

$$\frac{dy}{dx} = \frac{\frac{d}{dx}[\sin x]e^x - \sin x \frac{d}{dx}[e^x]}{e^{2x}} \quad \text{PoPo(5.1a)}$$

$$\frac{dy}{dx} = \frac{\cos x e^x - \sin x \frac{d}{dx}[e^x]}{e^{2x}} \quad \text{DSin(6.18)}$$

$$\frac{dy}{dx} = \frac{\cos x e^x - \sin x e^x}{e^{2x}} \quad \text{DExp(6.34)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^{2x}} \quad \text{DPF(4.3b)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^x e^x} \quad \text{PrCBPo(5.3b)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^x e^x} \quad \text{MId(4.9a)}$$

$$\frac{dy}{dx} = \frac{\cos x - \sin x}{e^x}$$

13.4 Derivative of a Rational Function

Example 13.16 – id:20151015-153507.Differentiate $f(x) = \frac{x^3 + 1}{x^2}$

(S)

Solution:

Method 1: Derivative of a quotient

$$[f(x)]' = \left[\frac{x^3 + 1}{x^2} \right]' \quad \text{SPE(4.15)}$$

$$f'(x) = \frac{[x^3 + 1]'(x^2) + -1((x^3 + 1)[x^2]')}{[x^2]^2} \quad \text{DQ(??)}$$

$$f'(x) = \frac{(3x^2)(x^2) - 1((x^3 + 1)[x^2]')}{[x^2]^2} \quad \text{differentiate goto 13.4}$$

$$f'(x) = \frac{(3x^2)(x^2) - 1((x^3 + 1)(2x))}{[x^2]^2} \quad \text{differentiate goto 13.2 } f'(x) = \frac{3x^3 - 2x^2 - 2}{x^3} \quad \text{Simplif}$$

14. Equations of Tangent & Secant Lines

14.1 Essential Questions

Essential Questions 14.1

1. How do we find the equation of the tangent line of a given function at the point $P(a, b)$?
2. How do we find the equation of the tangent line of a given function at $x = a$?

14.2 Finding the Equation of the Tangent Line




Figure 14.1: Finding the Equation of a Tangent Line Workflow

For a given function, $f(x)$, our goal is to find the equation of the tangent line at the point $P(a, b)$, which can be expressed in slope-intercept form as:

$$y = f'(a)x + k$$

Where $f'(a)$ is the value of the derivative, slope, at $x = a$ and k is the y -intercept. We therefore need to:

1. Find the derivative of the function $f(x)$: $f'(x)$
2. Find the value of the derivative at $x = a$: $f'(a)$
3. Find the value of the y -intercept: $k = -f'(a) + b$:
4. If the ordinate b is not explicitly given, then find $f(a) = b$

Example 14.1 – id:20151011-154209.

Find the equation of the line tangent to the curve of the function $f(x) = 2x^2 + 3x + 7$ at the point $P(2, 21)$.

(S)

Solution:

Find the derivative of $f(x)$

$$f'(x) = 4x + 3 \text{ goto 13.5}$$

Evaluate the derivative at $x = 2$

$$f'(2) = 4[2] + 3$$

SPE(4.15)

$$f'(2) = 8 + 3$$

OOM(2.3)

$$f'(2) = 11$$

OOA(2.1)

Find the y -intercept, k , of the equation of the tangent line.

$$y = f'(x)x + k$$

$$[21] = [11][2] + k \quad \text{SPE(4.15)}$$

$$21 = 22 + k$$

OOM(2.3)

$$\neg 22 + [21] = \neg 22 + [22 + k]$$

SPE+AI

$$\neg 22 + 21 = (\neg 22 + 22) + k$$

APA(4.2a)

$$\neg 1 = 0 + k$$

OOA(2.1)

$$\neg 1 = k$$

AId(4.4a)

$$\neg 1 = k$$

ONeg(3.1b)

$$k = -1$$

SyPE(4.16a)

The equation of the tangent line is

$$y = f'(2)x + k$$

$$y = 11x - 1 \quad \text{SPE(4.15)}$$

Example 14.2 – id:20151015-171630.

Given $f(x) = e^{3x}$, find the the equation of the tangent line L to the curve f at the point $(2, e^2)$.

(S)

Solution:

Find the derivative of $f(x)$

$$\begin{aligned}
 [f(x)]' &= [e^{3x}]' && \text{SPE(4.15)} \\
 f'(x) &= [3x]'[e^{3x}]' && \text{DCF(6.15)} \\
 f'(x) &= 3[x]'[e^{3x}]' && \text{DCM(6.3)} \\
 f'(x) &= 3 \cdot 1[e^{3x}]' && \text{DPo(6.5)} \\
 f'(x) &= 3[e^{3x}]' && \text{OOM(2.3)} \\
 f'(x) &= 3e^{3x} && \text{DExp(6.33)}
 \end{aligned}$$

Evaluate the derivative at $x = 2$

$$\begin{aligned}
 f'(2) &= 3e^{3[2]} && \text{SPE(4.15)} \\
 f'(2) &= 3e^6 && \text{OOM(2.3)}
 \end{aligned}$$

Find the y -intercept, k , of the equation of the tangent line.

$$\begin{aligned}
 y &= f'(x)x + k \\
 [e^2] &= [3e^6][2] + k && \text{SPE(4.15)} \\
 e^2 &= 3 \cdot e^6 \cdot 2 + k && \text{JTC(3.5)} \\
 e^2 &= 3 \cdot 2 \cdot e^6 + k && \text{CPM(4.6)} \\
 e^2 &= 6 \cdot e^6 + k && \text{OOM(2.3)} \\
 e^2 &= 6e^6 + k && \text{CTJ(3.6)} \\
 -6e^6 + [e^2] &= -6e^6 + [6e^6 + k] && \text{AI(4.5a), SPE(4.15)} \\
 e^2(-6e^4 + 1) &= -6e^6 + [6e^6 + k] && \text{DPF(4.3b)} \\
 -6e^6 + e^2 &= (-6e^6 + 6e^6) + k && \text{APA(4.2a)} \\
 e^2(-6e^4 + 1) &= 0 + k && \text{OOA(2.1)} \\
 e^2(-6e^4 + 1) &= k && \text{AId(4.4b)} \\
 e^2(-6e^4 + 1) &= k && \text{ONeg(3.1b)}
 \end{aligned}$$

The equation of the tangent line, L is

$$\begin{aligned}
 y &= f'(2)x + k \\
 y &= 3e^6x + e^2(-6e^4 + 1) && \text{SPE(4.15)}
 \end{aligned}$$

15. First Derivative Test

Example 15.1 – id:20151012-190708.

Given the function $f(x) = 2x^3 - 3x^2 - 12x + 1$ find the critical points, classify the critical points and find the intervals of increasing/decreasing.

(S)

Solution:

Differentiate the function:

$$f(x) = 2x^3 + -3x^2 + -12x + 1 \quad \text{DOS(4.12a)}$$

$$[f(x)]' = [2x^2 + -3x^2 + -12x + 1]' \quad \text{SPE(4.15)}$$

$$f'(x) = [2x^3]' + [-3x^2]' + [-12x]' + [1]' \quad \text{DS(6.7)}$$

$$f'(x) = 2[x^3]' + -3[x^2]' + -12[x]' + [1]' \quad \text{DCM(6.3)}$$

$$f'(x) = 2(3x^2) + -3(2x) + -12(1) + [1]' \quad \text{DPo(6.5)}$$

$$f'(x) = 6x^2 + -6x + -12 + [1]' \quad \text{OOM(2.3)}$$

$$f'(x) = 6x^2 + -6x + -12 + 0 \quad \text{DC(6.1)}$$

$$f'(x) = 6x^2 + -6x + -12 \quad \text{AId(4.4a)}$$

Solving the equation $6x^2 + -6x + -12 = 0$ to find the x value(s) of the critical points, goto 10.2, we find that $x = 2$ and $x = -1$ are critical values of the function.

Since we are looking for critical **points**, we need to find the ordinates, y -values, of the critical points by evaluating the function for the given critical x -values.

$$f(2) = -19 \text{ goto 11.1}$$

$$f(-1) = 8 \text{ goto 11.2}$$


Therefore the critical points are $(2, -19)$ and $(-1, 8)$.

The first derivative test can be used to determine the intervals of increasing/decreasing and consequently we will be able to classify the critical points. Since we are only interested in the values of the derivative, it will be easier to use the factored form of the derivative, $f'(x)=6(x-2)(x+1)$

1st Derivative Test Table

x	$x < -1$	$x = -1$	$-1 < x < 2$	$x = 2$	$x > 2$
$f'(x)$	+	0	-	0	+
$f(x)$	↗	→	↘	→	↗
C.P.		Max		Min	

From the first derivative table we can classify the critical point $(2, -19)$ as a local maximum and $(-1, 8)$ as a local minimum.



16. Curve Sketching

16.0.1 Finding the vertex of a quadratic function using differentiation.

We can find the vertex of a quadratic function, $f(x)$ using differentiation by:

1. Differentiate the function: Find $f'(x)$.
2. Set the derivative equal to zero: $f'(x) = 0$.
3. Find the abscissa of the vertex by solving the equation $f'(x) = 0$ for x to find the critical x value: $x = k$.
4. Find the ordinate of the vertex by substituting the value of critical value $x = k$ into the function $f(x)$: Evaluate $f(k)$

Example 16.1 – id:20151008-110208.

Find the vertex of the quadratic function, $f(x) = ax^2 + bx + c$, using differentiation.

(S)

Solution:

1. Find the derivative of $f(x)$

$$[f(x)]' = [ax^2 + bx + c]' \quad \text{SPE(4.15)}$$

$$f'(x) = [ax^2]' + [bx]' + [c]' \quad \text{DS(6.7)}$$

$$f'(x) = a[x^2]' + b[x]' + [c]' \quad \text{DCM(6.3)}$$

$$f'(x) = 2 \cdot a \cdot x + b \cdot 1 + [c]' \quad \text{DPo(6.5)}$$

$$f'(x) = 2 \cdot a \cdot x + 1 \cdot b + [c]' \quad \text{CPM(4.6)}$$

$$f'(x) = 2ax + 1b + [c]' \quad \text{CTJ(3.6)}$$

$$f'(x) = 2ax + b + [c]' \quad \text{MId(4.9b)}$$

$$f'(x) = 2ax + b + 0 \quad \text{DC(6.1)}$$

$$f'(x) = 2ax + b \quad \text{AId(4.4b)}$$

2. Set the derivative equal to zero and solve for x .

$$f'(x) = 0$$

$$2ax + b = 0$$

$$x = -\frac{b}{2a} \text{ goto 9.9}$$

The abscissa of the vertex is $x = -\frac{b}{2a}$.

3. Find the ordinate of the vertex by substituting the argument $x = -\frac{b}{2a}$ into $f(x)$

Example 16.2 – id:20150923-152515.

Find the vertex of the parabola $y = x^2 - 2x - 6$ using differentiation.

(S)

1. Differentiate the function.

$$f(x) = x^2 - 2x - 6$$

$$f(x) = x^2 + \neg 2x + \neg 6 \quad \text{DOS(4.12a)}$$

$$[f(x)]' = [x^2 + \neg 2x + \neg 6]' \quad \text{SPE(4.15)}$$

$$f'(x) = [x^2]' + [\neg 2x]' + [\neg 6]' \quad \text{DS(6.7)}$$

$$f'(x) = [x^2]' + \neg 2[x]' + [\neg 6]' \quad \text{DCM(6.3)}$$

$$f'(x) = 2x + \neg 2 + [\neg 6]' \quad \text{DPo(6.5)}$$

$$f'(x) = 2x + \neg 2 + 0 \quad \text{DC(6.1)}$$

$$f'(x) = 2x + \neg 2 \quad \text{AId(4.4b)}$$

$$f'(x) = 2x - 2 \quad \text{DOS(4.12b)}$$

- 2 and 3. Set the derivative equal to zero and solve for x

$$2x - 2 = 0$$

$$x = 1$$

4. Find the value of $f(1)$

$$f(x) = x^2 - 2x - 6$$

$$f(1) = [1]^2 - 2[1] - 6 \quad \text{SPE(4.15)}$$

$$f(1) = -7 \quad \text{Evaluate}$$

The vertex of this parabola is the point $(1, -7)$

■

17. Second Derivative Test

18. Curve Sketching




Figure 18.1: Overview of Curve Sketching

Bibliography

Books

Website

- [] *Indeterminate (variable)*. [http://en.wikipedia.org/wiki/Indeterminate_\(variable\)](http://en.wikipedia.org/wiki/Indeterminate_(variable)) (cited on page 33).
- [] *Polynomial*. <http://en.wikipedia.org/wiki/Polynomial> (cited on pages 33, 58).

Articles

Index

A

Additive Identity	15
Additive Inverse	16

C

Coefficient.....	31
Common Denominator	9
Cosine Inverse	74

D

Definition of Subtraction.....	17
Degree of the Indeterminate	31
Degree of the Univariate Polynomial	56
Derivative	77
DerivativeRules	
Derivative of a Composite Function	26
Derivative of a Constant	25
Derivative of a Constant Multiple ..	25
Derivative of a Difference.....	25
Derivative of a Logarithm	27
Derivative of a Natural Exponential	28
Derivative of a Natural Logarithm .	27
Derivative of a Power.....	25
Derivative of a Product.....	26
Derivative of a Quotient	26
Derivative of a Sum	25
Derivative of Cosecant.....	27
Derivative of Cosine	26
Derivative of Cotangent	27
Derivative of Secant	27
Derivative of Sine	26
Derivative of Tangent.....	26
Derivative Structural Rules	25
Logarithm Derivative Structural Rules	
27	
Monomial Derivative Rules.....	25

Trigonometric Derivative Structural Rules 26

Derivatives

Derivative of an Exponential.....	27
Dyadic Operations	9

E

Example

20141027-075159.....	34
20141105-161225.....	54
20141106-150622	36
20141106-151138	40
20141106-152020.....	37
20141106-152339	48
20141106-154547	40
20141107-121834	49
20141107-131748	65
20141108-173613	39
20141108-191616.....	41
20141108-194156	38
20141108-194431	38
20141108-194709.....	41
20141109-090809	42
20141109-091015	43
20141109-092448	43
20141109-092651.....	44
20141109-092910.....	44
20141109-093220	45
20141109-093419	45
20141109-094928	46
20141109-095151	46
20141109-095536.....	47
20141109-095842	48
20141109-133008.....	51
20141109-133316.....	51
20141109-140659	52
20141109-141019	53
20141109-141347	53
20141111-215726	62
20141111-222931	59

20141120-202042	33
20141120-202842	39
20141120-203846	33
20141121-093439	33
20141121-093747	32
20141121-184652	36
20141121-185558.....	34
20141121-190857	35
20141121-193636	35
20141124-141850	79
20141124-152503	80
20141124-153017	79
20141124-203850	83
20141124-205219	86
20141128-151834.....	81
20141128-155506	85
20141128-160248.....	84
20141206-101107	60
20141206-101632	60
20141206-102142	59
20141206-102404.....	61
20141206-102723	62
20141206-104404.....	61
20141209-142321	82
20141209-144203	82
20141209-145211	50
20141209-151354	83
20141219-212546.....	77
20150910-115935	83
20150923-152515	96
20151008-110208	95
20151011-154209	90
20151011-195002.....	81
20151012-190708	93
20151012-192313	65
20151012-201647	73
20151012-203549	73
20151015-104754	63
20151015-153507.....	87
20151015-155838	80
20151015-165037	86
20151015-171630	90
Exponential Inverse	74
 F	
Fraction Operation of Addition	10
Function Value Argument Substitution	73
Functions	
Inverses	74
 G	
Greatest Common Divisor	7
 I	
Identities	
Power Identities.....	19, 20
Improper Fraction	7
Indeterminate	31
 L	
Like Terms	32
Logarithmic Inverse	74
LogarithmIdentity	
Logarithm Power of a Power.....	20
Logarithm Product of Common Base Powers	20
Logarithm Quotient of Common Base Powers	20
Logartihms	
Logarithm Exponent Visible	12
Logarithm to Power	12
 M	
Multiplicative Identity	16
Multiplicative Inverse	16
 N	
Notation	
Center-Dot to Juxtaposition.....	12
Derivative Notation	12
Euler's first derivative	13
Euler's nth derivative	14
Euler's second derivative	13
Juxtapostion to Center-Dot	12
LaGrange's Evaluate derivative....	13
LaGrange's first derivative.....	13
LaGrange's nth derivative	13
LaGrange's second derivative.....	13
Leibniz's Evaluate derivative	13

Leibniz's first derivative	13
Leibniz's nth derivative	13
Leibniz's second derivative	13
Logarithm Notation	12
Multiplication Center-Dot	11
Multiplication Juxtaposition	11
Multiplication Notation	11
Multiplication Times	11
Negation Notation.....	11
Power Notation	12
Number System	
Integers	7
Natural Numbers.....	7
Rational Numbers	7
Number Systems	7

O

Operation	
Operation of Addition.....	9
Operation of Exponentiation.....	9
Operation of Multiplication.....	9
Operation of Negation.....	11

P

Positive Integers	7
Power Identity.....	17
Power Inverse	74
Powers	
Factor to Power.....	12
Power Negative Exponent	12
Power of a Power	19
Power of a Product of Powers	19
Power of a Quotient of Powers	19
Power to Factor	12
Power To Logarithm.....	12
Product Common Base Powers	19
Quotient Common Base Powers	19
Radical to Power	12
Prime Numbers.....	7
Proper Fraction	7
Properties	
Summary of Field Properties	15
Properties of Addition.....	15
Property	
Associative Property of Addition ..	15

Associative Property of Multiplication	
16	
Commutative Property of Addition	15
Commutative Property of Multiplication	16
Distributive Property Factoring	15, 16
Inverses	74
Properties of Equality	17
Properties of Inequality	18
Properties of Multiplication	16
Properties of Powers.....	17
Properties of Subtraction	17
Zero Product.....	17

R

Reflexive Property of Equality	17
Relatively Prime	8

S

Simplifying Degree 0 Univariate Monomials	31
Sine Inverse	74
Solving Quadratic Equations	
Completing the Square	66
Power Inverse	59, 65
Substitution Property of Equality	17
Substitution Property of Inequality	18
Symmetric Property of Equality	17

T

Tangent Inverse	74
Transitive Property of Equality	17
Transitive Property of Inequality	18
TrigonometricIdentity	
Cosine Difference of Angles Identity	21
Cosine Double Angle Identity	21
Cosine Sum of Angles Identity	21
Sine Difference of Angles Identity ..	21
Sine Double Angle Identity	21
Sine Sum of Angles Identity	21
Tangent Difference of Angles Identity	
22	
Tangent Double Angle Identity	21
Tangent Sum of Angles Identity ..	21

Trigonometric Cotangent Identity . 21
Trigonometric Pythagorean Identities
 20
Trigonometric Reciprocal Identities 20
Trigonometric Tangent Identity 20

U

Univariate Monomial 31
Univariate Polynomial Expression 56

Z

Zero Factor Property 18