

Mathematics Notebook

by Mark Olson : v42.org

Updated: October 16, 2015

Contents

I	Reference
1	Numbers 9
1.1	Number Systems 9
1.2	Prime Numbers 9
1.2.1	Listing of Prime Numbers 2-997 10
2	Operations 11
2.1	Dyadic Operations 11
3	Notation 13
3.1	Negation Notation 13
3.2	Multiplication Notation 13
3.3	Power Notation 14
3.4	Logarithm Notation 14
3.5	Derivative Notation 14
4	Properties 17
4.1	Summary of Field Properties 17
4.2	Properties of Addition 17
4.3	Properties of Multiplication 18
4.4	Properties of Subtraction 19
4.5	Properties of Powers 19
4.6	Properties of Equality 19
4.7	Properties of Inequality 20
5	Identities 21
5.1	Power Identities 21
5.2	Logarithm Identities 22
5.3	Trigonometric Identities 22
6	Limit Properties 27
6.1	Algebraic Limit Theorem 27

7	Calculus Rules	29
7.1	Monomial Derivative Rules	29
7.2	Derivative Structural Rules	29
7.3	Trigonometric Derivative Rules	30
7.4	Logarithm Derivative Rules	31
7.5	Exponential Derivative Rules	31

II

Algebra

8	Simplifying Univariate Polynomials	35
8.1	Simplifying Degree 0 Univariate Monomials	35
8.2	Simplifying Degree 1 Univariate Monomials	37
8.3	Simplifying Degree 2 Univariate Monomials	43
8.4	Simplifying Degree 3 Univariate Monomials	46
8.5	Simplifying Degree 1 Univariate Binomials	46
8.6	Simplifying Degree 2 Univariate Binomials	53
8.7	Simplifying Degree 2 Univariate Trinomials	55
8.8	Simplifying Degree n Univariate Polynomials	59
9	Simplifying Univariate Monomials	61
9.1		61
10	Factoring Univariate Trinomials	63
11	Solving Linear Equations	65
11.1	Power Inverse	65
12	Solving Quadratic Equations	71
12.1	Power Inverse	71
12.2	Completing The Square	72

III

Functions

13	Functions	79
13.1	Inverse Functions	80
13.2	Inverses	80

IV

Differential Calculus

14	Derivative by First Principles	83
14.1	Limit of the Difference Quotient	83

15	Derivative Rules	85
15.1	Derivative of a Monomial Functions	85
15.2	Derivative of Polynomial Functions	86
15.3	Derivative of a Quotient	92
15.4	Derivative of a Rational Function	93
16	Equations of Tangent & Secant Lines	95
16.1	Essential Questions	95
16.2	Finding the Equation of the Tangent Line	95
17	First Derivative Test	99
18	Curve Sketching	101
18.0.1	Finding the vertex of a quadratic function using differentiation.	101
19	Second Derivative Test	103
20	Curve Sketching	105
	Bibliography	107
	Bibliography	107
	Websites	107
	Articles	107
	Index	109

Reference

1	Numbers	9
1.1	Number Systems	
1.2	Prime Numbers	
2	Operations	11
2.1	Dyadic Operations	
3	Notation	13
3.1	Negation Notation	
3.2	Multiplication Notation	
3.3	Power Notation	
3.4	Logarithm Notation	
3.5	Derivative Notation	
4	Properties	17
4.1	Summary of Field Properties	
4.2	Properties of Addition	
4.3	Properties of Multiplication	
4.4	Properties of Subtraction	
4.5	Properties of Powers	
4.6	Properties of Equality	
4.7	Properties of Inequality	
5	Identities	21
5.1	Power Identities	
5.2	Logarithm Identities	
5.3	Trigonometric Identities	
6	Limit Properties	27
6.1	Algebraic Limit Theorem	
7	Calculus Rules	29
7.1	Monomial Derivative Rules	
7.2	Derivative Structural Rules	
7.3	Trigonometric Derivative Rules	
7.4	Logarithm Derivative Rules	
7.5	Exponential Derivative Rules	

1. Numbers

1.1 Number Systems

Definition 1.1.1 – Natural Numbers.

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

- (R) It is not uncommon for zero to be excluded from the natural numbers. In fact, some exclude zero from the natural numbers and then describe the set of natural numbers that include zero the whole numbers.

$$\mathbb{W} = \{0, 1, 2, 3, \dots\}$$

For the purposes of these notes, zero will be included within the set of natural numbers.

Definition 1.1.2 – Integers.

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Definition 1.1.3 – Positive Integers.

$$\mathbb{Z}^+ = \{1, 2, 3, \dots\}$$

Definition 1.1.4 – Rational Numbers.

$$\mathbb{Q} = \{m/n \mid m, n \in \mathbb{Z}, n \neq 0\}$$

Definition 1.1.5 – Proper Fraction. Given $m < n$, then the fraction m/n is called **proper**.

Definition 1.1.6 – Improper Fraction. Given $m > n$, then the fraction m/n is called **improper**.

1.2 Prime Numbers

Definition 1.2.1 – Greatest Common Divisor. Suppose that m and n are positive integers. The greatest common divisor is the largest divisor (factor) common to both m and n .

Definition 1.2.2 – Relatively Prime. Two integers m and n are relatively prime to each other, $m \perp n$, if they share no common positive integer divisors (factors) except 1.

$$m \perp n \text{ if } \gcd(m, n) = 1.$$

1.2.1 Listing of Prime Numbers 2-997

2	3	5	7	11	13	17	19	23	29	31	37
41	43	47	53	59	61	67	71	73	79	83	89
97	101	103	107	109	113	127	131	137	139	149	151
157	163	167	173	179	181	191	193	197	199	211	223
227	229	233	239	241	251	257	263	269	271	277	281
283	293	307	311	313	317	331	337	347	349	353	359
367	373	379	383	389	397	401	409	419	421	431	433
439	443	449	457	461	463	467	479	487	491	499	503
509	521	523	541	547	557	563	569	571	577	587	593
599	601	607	613	617	619	631	641	643	647	653	659
661	673	677	683	691	701	709	719	727	733	739	743
751	757	761	769	773	787	797	809	811	821	823	827
829	839	853	857	859	863	877	881	883	887	907	911
919	929	937	941	947	953	967	971	977	983	991	997

2. Operations

2.1 Dyadic Operations

Definition 2.1.1 – Operation of Addition (OOA).

$$\underbrace{\begin{array}{c} a \\ \text{Augend} \end{array}}_{\text{Sum}} + \underbrace{\begin{array}{c} b \\ \text{Addend} \end{array}}_{\text{Sum}} \quad (2.1)$$

More generally,

$$\underbrace{\begin{array}{c} a \\ \text{Summand} \end{array}}_{\text{Sum}} + \underbrace{\begin{array}{c} b \\ \text{Summand} \end{array}}_{\text{Sum}} \quad (2.2)$$

Definition 2.1.2 – Operation of Multiplication (OOM).

$$\underbrace{\begin{array}{c} a \\ \text{Multiplicand} \end{array}}_{\text{Product}} \times \underbrace{\begin{array}{c} b \\ \text{Multiplier} \end{array}}_{\text{Product}} \quad (2.3)$$

More generally,

$$\underbrace{\begin{array}{c} a \\ \text{Factor} \end{array}}_{\text{Product}} \times \underbrace{\begin{array}{c} b \\ \text{Factor} \end{array}}_{\text{Product}} \quad (2.4)$$

Definition 2.1.3 – Operation of Exponentiation (OOE).

$$\underbrace{\begin{array}{c} b \\ \text{base} \end{array}}_{\text{Power}}^m \quad (2.5)$$

Definition 2.1.4 – Common Denominator (CD).

$$\frac{a}{b} + \frac{c}{b} = \frac{a+c}{b} \quad (2.6a)$$

$$\frac{a+c}{b} = \frac{a}{b} + \frac{c}{b} \quad (2.6b)$$

Rule 2.1.1 – Fraction Operation of Addition (FOOA).

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd} \quad (2.7a)$$

$$\frac{ad + bc}{bd} = \frac{a}{b} + \frac{c}{d} \quad (2.7b)$$

3. Notation

3.1 Negation Notation

Notation 3.1.1 – Operation of Negation (ONeg).

$$\neg a = \neg a \quad (3.1a)$$

$$\neg a = -a \quad (3.1b)$$

I have used a different symbol, \neg , as the prefix negation operator only to differentiate it from the minus sign infix operator symbol, $-$, which is also used as the infix operator for the dyadic operation of subtraction. I will refer to this change of symbol as ONeg. This is used only as a teaching tool and should not be confused with the logic negation operator. Another advantage of using this symbol is that it reduces the number of delimiters used in an expression for example, $\neg a$ versus $(-a)$.

- Negative five: -5
- Negative five: $\neg 5$
- Four minus five: $4 - 5$
- Four minus negative five: $4 - \neg 5$
- Four minus negative five: $4 - (-5)$
- Four minus negative five: $4 - \neg 5$
- Negative four minus five: $-4 - 5$
- Negative four minus five: $\neg 4 - 5$

3.2 Multiplication Notation

Notation 3.2.1 – Multiplication Center-Dot (MC).

$$a \cdot b \quad (3.2)$$

Notation 3.2.2 – Multiplication Juxtaposition (MJ).

$$ab, a(b), (a)b, (a)(b), a[b], [a]b, [a][b] \quad (3.3)$$

Notation 3.2.3 – Multiplication Times (MT).

$$a \times b \quad (3.4)$$

Notation 3.2.4 – Juxtaposition to Center-Dot (JTC).

$$ab = a \cdot b \quad (3.5)$$

Notation 3.2.5 – Center-Dot to Justaposition (CTJ).

$$a \cdot b = ab \quad (3.6)$$

3.3 Power Notation

Notation 3.3.1 – Power Exponent Negative Exponent (PoNegE).

$$b^{-k} = \frac{1}{b^k} \quad (3.7)$$

$$\frac{1}{b^k} = b^{-k} \quad (3.8)$$

Notation 3.3.2 – Power To Factor (PoTF).

$$a^n = a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_n \quad (3.9)$$

Notation 3.3.3 – Power To Logarithm (PoTL).

$$y = b^x \Rightarrow x = \log_b y \quad (3.10)$$

Notation 3.3.4 – Factor To Power (FTPo).

$$a_1 \cdot a_2 \cdot \dots \cdot a_{n-1} \cdot a_n = a^n \quad (3.11)$$

Notation 3.3.5 – Radical To Power (RTPo).

$$\sqrt[m]{b^n} = b^{\frac{n}{m}} \quad (3.12)$$

3.4 Logarithm Notation

Notation 3.4.1 – Logarithm Exponent Visible (LEV).

$$\log_b y \Rightarrow \log_b y = x \quad (3.13)$$

Notation 3.4.2 – Logarithm to Power (LTPo).

$$x = \log_b y \Rightarrow y = b^x \quad (3.14)$$

3.5 Derivative Notation

Notation 3.5.1 – Leibniz's first derivative.

$$\frac{dy}{dx} = \frac{d[f(x)]}{dx} = \frac{d}{dx}[f(x)] \quad (3.15)$$

Notation 3.5.2 – Leibniz's second derivative.

$$\frac{d^2y}{dx^2} \quad (3.16)$$

Notation 3.5.3 – Leibniz's nth derivative.

$$\frac{d^n y}{dx^n} \quad (3.17)$$

Notation 3.5.4 – Leibniz's Evaluate derivative.

$$\left. \frac{dy}{dx} \right|_{x=a} = \frac{dy}{dx}(a) \quad (3.18)$$

Notation 3.5.5 – LaGrange's first derivative.

$$f'(x) \quad (3.19)$$

Notation 3.5.6 – LaGrange's second derivative.

$$f''(x) \quad (3.20)$$

Notation 3.5.7 – LaGrange's nth derivative.

$$f^{(n)}(x) \quad (3.21)$$

Notation 3.5.8 – LaGrange's Evaluate derivative.

$$f'(a) \quad (3.22)$$

Notation 3.5.9 – Euler's first derivative.

$$Df = D_x f \quad (3.23)$$

Notation 3.5.10 – Euler's second derivative.

$$D^2 f = D_x^2 f \quad (3.24)$$

Notation 3.5.11 – Euler's nth derivative.

$$D^n f = D_x^n \quad (3.25)$$

4. Properties

4.1 Summary of Field Properties

Name	Addition	Multiplication
Commutative	$a + b = b + a$	$a \cdot b = b \cdot a$
Associative	$(a + b) + c = a + (b + c)$	$(a \cdot b) \cdot c = a \cdot (b \cdot c)$
Distributive	$a(b + c) = ab + ac$	$(a + b)c = ac + bc$
Identity	$a + \mathbf{0} = a = \mathbf{0} + a$	$a \cdot \mathbf{1} = a = \mathbf{1} \cdot a$
Inverse	$a + (-a) = 0 = (-a) + a$	$a \cdot a^{-1} = 1 = a^{-1} \cdot a$

Table 4.1: Summary of the Field Properties

4.2 Properties of Addition

Property 4.2.1 – Commutative Property of Addition (CPA).

$$ab = ba \quad (4.1)$$

Property 4.2.2 – Associative Property of Addition (APA).

$$a + b + c = (a + b) + c \quad (4.2a)$$

$$a + b + c = a + (b + c) \quad (4.2b)$$

Property 4.2.3 – Distributive Property Factoring (DPF).

$$ba + ca = (b + c)a \quad (4.3a)$$

$$ab + ac = a(b + c) \quad (4.3b)$$

Property 4.2.4 – Additive Identity (AId).

$$a + \mathbf{0} = a \quad (4.4a)$$

$$a = a + \mathbf{0} \quad (4.4b)$$

Property 4.2.5 – Additive Inverse (AI).

$$a + (-a) = 0 \quad (4.5a)$$

4.3 Properties of Multiplication

Property 4.3.1 – Commutative Property of Multiplication (CPM).

$$a \cdot b = b \cdot a \quad (4.6)$$

Property 4.3.2 – Associative Property of Multiplication (APM).

$$a \cdot b \cdot c = (a \cdot b) \cdot c \quad (4.7a)$$

$$a \cdot b \cdot c = a \cdot (b \cdot c) \quad (4.7b)$$

Property 4.3.3 – Distributive Property Expanding (DPE).

$$a(b + c) = ab + ac \quad (4.8a)$$

$$(b + c)a = ba + ca \quad (4.8b)$$

Property 4.3.4 – Multiplicative Identity (Mid).

$$1a = a \quad (4.9a)$$

$$a = 1a \quad (4.9b)$$

- R** If the coefficient of a univariate monomial is the multiplicative identity 4.9a, 1, then it is not shown in its canonical form.

$$\begin{aligned} C_k x^k &= C_k x^k \\ &= 1x^k \\ &= x^k \end{aligned}$$

Property 4.3.5 – Multiplicative Inverse (MI).

$$a \cdot \frac{1}{a} = 1 \quad (4.10a)$$

$$a \cdot a^{-1} = 1 \quad (4.10b)$$

Property 4.3.6 – Zero Product (ZPr).

$$\text{if } a \cdot b = 0, \text{ then } a = 0 \text{ or } b = 0 \quad (4.11a)$$

4.4 Properties of Subtraction

Definition 4.4.1 – Definition of Subtraction (DOS).

$$a - b = a + \neg b \quad (4.12a)$$

$$a + \neg b = a - b \quad (4.12b)$$

4.5 Properties of Powers

Property 4.5.1 – Power Inverse (Pold).

$$1 = b^0 \quad (4.13a)$$

$$b^0 = 1 \quad (4.13b)$$

4.6 Properties of Equality

Property 4.6.1 – Reflexive Property of Equality (RPE).

$$a = a \quad (4.14a)$$

Property 4.6.2 – Substitution Property of Equality (SPE).

Given $a = b$, then

$$E(a) = E(b) \quad (4.15)$$

$E(x)$ represents any expression.

Property 4.6.3 – Symmetric Property of Equality (SyPE).

$$a = b \quad \text{then} \quad b = a \quad (4.16a)$$

Property 4.6.4 – Transitive Property of Equality (TPE).

$$\text{if } a = b \quad \text{and} \quad b = c \quad \text{then} \quad a = c \quad (4.17a)$$

Property 4.6.5 – Zero Factor Property (ZFP).

$$\text{if } a \cdot b = 0 \text{ then } a = 0 \text{ or } b = 0 \quad (4.18a)$$

4.7 Properties of Inequality

Property 4.7.1 – Substitution Property of Inequality (SPIn).

$$a < b \text{ then } a + c < b + c \quad (4.19a)$$

$$a < b \text{ and } c > 0, \text{ then } ca < cb \quad (4.19b)$$

$$a < b \text{ and } c < 0, \text{ then } ca > cb \quad (4.19c)$$

Property 4.7.2 – Transitive Property of Inequality (TPIn).

$$\text{if } a < b \text{ and } b < c \text{ then } a > c \quad (4.20a)$$

5. Identities

5.1 Power Identities

Identity 5.1.1 – Power of Power (PoPo).

$$(b^m)^k = b^{m \cdot k} \quad (5.1a)$$

$$b^{m \cdot k} = (b^m)^k \quad (5.1b)$$

Identity 5.1.2 – Power of a Product (PoPr).

$$(a \cdot b)^k = a^k \cdot b^k \quad (5.2a)$$

$$a^k \cdot b^k = (a \cdot b)^k \quad (5.2b)$$

Identity 5.1.3 – Product Common Base Powers (PrCBPo).

$$b^m \cdot b^n = b^{m+n} \quad (5.3a)$$

$$b^{m+n} = b^m \cdot b^n \quad (5.3b)$$

Identity 5.1.4 – Quotient Common Base Powers (QCBPo).

$$\frac{b^m}{b^n} = b^{m-n} \quad (5.4a)$$

$$b^{m-n} = \frac{b^m}{b^n} \quad (5.4b)$$

Identity 5.1.5 – Power of a Quotient of Powers (PoQPo).

$$\left(\frac{a^m}{b^n}\right)^k = \frac{a^{m \cdot k}}{b^{n \cdot k}} \quad (5.5a)$$

$$\frac{a^{m \cdot k}}{b^{n \cdot k}} = \left(\frac{a^m}{b^n}\right)^k \quad (5.5b)$$

Identity 5.1.6 – Power of a Product of Powers (PoPrPo).

$$(a^m \cdot b^n)^k = a^{m \cdot k} \cdot b^{n \cdot k} \quad (5.6a)$$

$$a^{m \cdot k} \cdot b^{n \cdot k} = (a^m \cdot b^n)^k \quad (5.6b)$$

5.2 Logarithm Identities

Identity 5.2.1 – Logarithm Power of a Power (LPoPo).

$$\log_b x^n = n \log_b x \quad (5.7a)$$

$$n \log_b x = \log_b x^n \quad (5.7b)$$

Identity 5.2.2 – Logarithm Product of Common Base Powers (LPrCBPo).

$$\log_b(mn) = \log_b m + \log_b n \quad (5.8a)$$

$$\log_b m + \log_b n = \log_b(mn) \quad (5.8b)$$

Identity 5.2.3 – Logarithm Quotient of Common Base Powers (LQCBPo).

$$\log_b \left(\frac{m}{n} \right) = \log_b m - \log_b n \quad (5.9a)$$

$$\log_b m - \log_b n = \log_b \left(\frac{m}{n} \right) \quad (5.9b)$$

5.3 Trigonometric Identities

Identity 5.3.1 – Trigonometric Reciprocal Identities (TRId).

$$\sin \theta = \frac{1}{\csc \theta} \quad (5.10a)$$

$$\cos \theta = \frac{1}{\sec \theta} \quad (5.10b)$$

$$\cot \theta = \frac{1}{\tan \theta} \quad (5.10c)$$

$$\csc \theta = \frac{1}{\sin \theta} \quad (5.10d)$$

$$\sec \theta = \frac{1}{\cos \theta} \quad (5.10e)$$

$$\tan \theta = \frac{1}{\cot \theta} \quad (5.10f)$$

Identity 5.3.2 – Trigonometric Pythagorean Identities (TPythagId).

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (5.11a)$$

$$\sec^2 \theta = \tan^2 \theta + 1 \quad (5.11b)$$

$$\csc^2 \theta = 1 + \cot^2 \theta \quad (5.11c)$$

Identity 5.3.3 – Trigonometric Tangent Identity (TanId).

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad (5.12a)$$

Identity 5.3.4 – Trigonometric Cotangent Identity (CotId).

$$\cot \theta = \frac{\cos \theta}{\sin \theta} \quad (5.13a)$$

Identity 5.3.5 – Sine Double Angle Identity (SinDAId).

$$\sin 2\theta = 2 \sin \theta \cos \theta \quad (5.14a)$$

Identity 5.3.6 – Cosine Double Angle Identity (CosDAId).

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta \quad (5.15a)$$

$$= 1 - 2 \sin^2 \theta \quad (5.15b)$$

$$= 2 \cos^2 \theta - 1 \quad (5.15c)$$

Identity 5.3.7 – Tangent Double Angle Identity (TanDAId).

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} \quad (5.16a)$$

Identity 5.3.8 – Sine Sum of Angles Identity (SinSAId).

$$\sin(\theta + \phi) = \sin \theta \cos \phi + \cos \theta \sin \phi \quad (5.17a)$$

Identity 5.3.9 – Sine Difference of Angles Identity (SinDiffAId).

$$\sin(\theta - \phi) = \sin \theta \cos \phi - \cos \theta \sin \phi \quad (5.18a)$$

Identity 5.3.10 – Cosine Sum of Angles Identity (CosSAId).

$$\cos(\theta + \phi) = \cos \theta \cos \phi - \sin \theta \sin \phi \quad (5.19a)$$

Identity 5.3.11 – Cosine Difference of Angles Identity (CosDiffAId).

$$\cos(\theta - \phi) = \cos \theta \cos \phi + \sin \theta \sin \phi \quad (5.20a)$$

Identity 5.3.12 – Tangent Sum of Angles Identity (TanSAId).

$$\tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} \quad (5.21a)$$

Identity 5.3.13 – Tangent Difference of Angles Identity (TanDiffAld).

$$\tan(\theta - \phi) = \frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi} \quad (5.22a)$$




Figure 5.1: Simplifying Expressions Workflow:

■ Property, ■ Operation, ■ Notation, ■ Powers, ■ Delimiters, ■ Process, ■ Not Used

6. Limit Properties

6.1 Algebraic Limit Theorem

Rule 6.1.1 – Algebraic Limit Theorem of a Constant (ALTC). If $g(x) = a$, where a is a constant, then

$$\lim_{x \rightarrow c} [A] = A \quad (6.1)$$

Rule 6.1.2 – Algebraic Limit Theorem of a Sum (ALTS). If both the limits $\lim_{x \rightarrow c} g(x) = L_1$ and $\lim_{x \rightarrow c} h(x) = L_2$ exist, then

$$\lim_{x \rightarrow c} [g(x) + h(x)] = \lim_{x \rightarrow c} g(x) + \lim_{x \rightarrow c} h(x) \quad (6.2)$$

Rule 6.1.3 – Algebraic Limit Theorem of a Difference (ALTD). If both the limits $\lim_{x \rightarrow c} g(x) = L_1$ and $\lim_{x \rightarrow c} h(x) = L_2$ exist, then

$$\lim_{x \rightarrow c} [g(x) - h(x)] = \lim_{x \rightarrow c} g(x) - \lim_{x \rightarrow c} h(x) \quad (6.3)$$

Rule 6.1.4 – Algebraic Limit Theorem of a Product (ALTPr). If both the limits $\lim_{x \rightarrow c} g(x) = L_1$ and $\lim_{x \rightarrow c} h(x) = L_2$ exist, then

$$\lim_{x \rightarrow c} [g(x) \cdot h(x)] = \lim_{x \rightarrow c} g(x) \cdot \lim_{x \rightarrow c} h(x) \quad (6.4)$$

Rule 6.1.5 – Algebraic Limit Theorem of a Quotient (ALTQ). If both the limits $\lim_{x \rightarrow c} g(x) = L_1$ and $\lim_{x \rightarrow c} h(x) = L_2$ exist and $L_2 \neq 0$, then

$$\lim_{x \rightarrow c} \left[\frac{g(x)}{h(x)} \right] = \frac{\lim_{x \rightarrow c} g(x)}{\lim_{x \rightarrow c} h(x)} \quad (6.5)$$

7. Calculus Rules

7.1 Monomial Derivative Rules

Rule 7.1.1 – Derivative of a Constant (DC).

$$[c]' = 0 \quad (7.1)$$

$$\frac{d}{dx} [c] = 0 \quad (7.2)$$

Rule 7.1.2 – Derivative of a Constant Multiple (DCM).

$$[cf(x)]' = c[f(x)]' \quad (7.3)$$

$$\frac{d}{dx} [cf(x)] = c \frac{d}{dx} [f(x)] \quad (7.4)$$

Rule 7.1.3 – Derivative of a Power (DPo).

$$[x^n]' = nx^{n-1} \quad (7.5)$$

$$\frac{d}{dx} [x^n] = nx^{n-1} \quad (7.6)$$

7.2 Derivative Structural Rules

Rule 7.2.1 – Derivative of a Sum (DS).

$$[f(x) + g(x)]' = f'(x) + g'(x) \quad (7.7)$$

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} [f(x)] + \frac{d}{dx} [g(x)] \quad (7.8)$$

Rule 7.2.2 – Derivative of a Difference (DD).

$$[f(x) - g(x)]' = f'(x) - g'(x) \quad (7.9)$$

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} [f(x)] - \frac{d}{dx} [g(x)] \quad (7.10)$$

Rule 7.2.3 – Derivative of a Product (DPr).

$$[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x) \quad (7.11)$$

$$\frac{d}{dx} [f(x)g(x)] = \frac{d}{dx} [f(x)]g(x) + f(x)\frac{d}{dx} [g(x)] \quad (7.12)$$

Rule 7.2.4 – Derivative of a Quotient (DQ).

$$\left[\frac{f(x)}{g(x)} \right]' = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2} \quad (7.13)$$

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{\frac{d}{dx} [f(x)] g(x) - f(x) \frac{d}{dx} [g(x)]}{[g(x)]^2} \quad (7.14)$$

Rule 7.2.5 – Derivative of a Composite Function (DComp).

$$[f(g(x))]' = [g(x)]' [f(g(x))]' \quad (7.15)$$

$$\frac{d}{dx} [f(g(x))] = \frac{d}{dx} [g(x)] \frac{d}{dx} [f(g(x))] \quad (7.16)$$

7.3 Trigonometric Derivative Rules

Rule 7.3.1 – Derivative of Sine (DSin).

$$[\sin(x)]' = \cos(x) \quad (7.17)$$

$$\frac{d}{dx} [\sin(x)] = \cos(x) \quad (7.18)$$

Rule 7.3.2 – Derivative of Cosine (DCos).

$$[\cos(x)]' = -\sin(x) \quad (7.19)$$

$$\frac{d}{dx} [\cos(x)] = -\sin(x) \quad (7.20)$$

Rule 7.3.3 – Derivative of Tangent (DTan).

$$[\tan x]' = \sec^2 \quad (7.21)$$

$$\frac{d}{dx} [\tan x] = \sec^2 \quad (7.22)$$

Rule 7.3.4 – Derivative of Cosecant (DCsc).

$$[\csc x]' = -\csc x \cot x \quad (7.23)$$

$$\frac{d}{dx} [\csc x] = -\csc x \cot x \quad (7.24)$$

Rule 7.3.5 – Derivative of Secant (DSec).

$$[\sec x]' = \sec x \tan x \quad (7.25)$$

$$\frac{d}{dx} [\sec x] = \sec x \tan x \quad (7.26)$$

Rule 7.3.6 – Derivative of Cotangent (DCot).

$$[\cot x]' = -\csc^2 x \quad (7.27)$$

$$\frac{d}{dx} [\cot x] = -\csc^2 x \quad (7.28)$$

7.4 Logarithm Derivative Rules

Rule 7.4.1 – Derivative of a Logarithm (DL).

$$[\log_a x]' = \frac{1}{x \ln a} \quad (7.29)$$

$$\frac{d}{dx} [\log_a x] = \frac{1}{x \ln a} \quad (7.30)$$

Rule 7.4.2 – Derivative of a Natural Logarithm (DNL).

$$[\ln x]' = \frac{1}{x} \quad (7.31)$$

$$\frac{d}{dx} [\ln x] = \frac{1}{x} \quad (7.32)$$

7.5 Exponential Derivative Rules

Rule 7.5.1 – Derivative of an Exponential(DExp).


$$[a^x]' = a^x \ln a \quad (7.33)$$

$$\frac{d}{dx} [a^x] = a^x \ln a \quad (7.34)$$

Rule 7.5.2 – Derivative of a Natural Exponential(DNExp).

$$[e^x]' = e^x \quad (7.35)$$

$$\frac{d}{dx} [e^x] = e^x \quad (7.36)$$



Algebra

8	Simplifying Univariate Polynomials	35
8.1	Simplifying Degree 0 Univariate Monomials	
8.2	Simplifying Degree 1 Univariate Monomials	
8.3	Simplifying Degree 2 Univariate Monomials	
8.4	Simplifying Degree 3 Univariate Monomials	
8.5	Simplifying Degree 1 Univariate Binomials	
8.6	Simplifying Degree 2 Univariate Binomials	
8.7	Simplifying Degree 2 Univariate Trinomials	
8.8	Simplifying Degree n Univariate Polynomials	
9	Simplifying Univariate Monomials	61
9.1		
10	Factoring Univariate Trinomials	63
11	Solving Linear Equations	65
11.1	Power Inverse	
12	Solving Quadratic Equations	71
12.1	Power Inverse	
12.2	Completing The Square	

8. Simplifying Univariate Polynomials

The definition of a univariate polynomial expression is based on the expanded canonical form of some polynomial expression. It might be that the original expression might not be in the expanded canonical form, so a process called **simplifying by expanding** will be introduced to manipulate the expression such that it can be written in its expanded canonical form.

This process of simplifying by expanding polynomial expressions will be developed to the extent that it will be used to simplify multivariate polynomials. We will start by simplifying univariate monomial expressions.

8.1 Simplifying Degree 0 Univariate Monomials

Definition 8.1.1 – Indeterminate.

x

An indeterminate is a symbol that is treated as a variable, but does not stand for anything else but itself and is used as a placeholder.

- it does **not** designate a constant or a parameter
- it is **not** an unknown that could be solved for
- it is **not** a variable designating a function argument

[1]

Definition 8.1.2 – Degree of the Indeterminate.

x^k

The exponent of an indeterminate power, k is called the degree of the indeterminate.

Polynomial [3]

Definition 8.1.3 – Coefficient.

Cx^k

A coefficient, C is a real number multiplicative factor.

Definition 8.1.4 – Univariate Monomial.

$C_k x^k$

A univariate monomial is made up of two factors. The first factor of a monomial, C_k , is the **coefficient**. The second factor of each monomial, x^k , is an indeterminate raised to a non-negative integer power k .

Degree 0 univariate polynomial expressions are made up of univariate monomials, C_0 , called **constants**. The power identity is an indeterminate raised to a power of 0 has a value of 1. Thus, $x^0 = 1$ and results in the monomial $C_0 \cdot 1$. The canonical form of a product does not show the multiplicative identity factor, so what remains of this monomial product is only the coefficient factor C_0 and from now on will be referred to as a **constant**.

Degree 0 univariate polynomial expressions are usually a monomial in their canonical form if C_0 is a non-zero real number. The exception is if $C_0 = 0$, the additive identity, then the result is the zero polynomial, which can be considered a degree -1 polynomial.

The expression can be manipulated into its monomial canonical form by simplifying the expression. Simplifying the expression can be defined as evaluating the expression by following order of operations, which is the same as evaluating an arithmetic expression.

Definition 8.1.5 – Univariate Like Terms.

$$C_1x^k = C_2x^k$$

Two or more univariate monomials are defined as having like terms if each monomial has the same term, which will be the same indeterminate raised to the same positive integer power.

Sometimes the word **term** is used to describe monomials (including both the coefficient and the term), which may be confusing when trying to define like terms. For this reason, we will refer to the summands of a polynomial as monomials.

The monomials $5x^1$ and $3x^1$ can be described as having like terms because they share the common term x^1 . One could also say that $5x^1$ and $3x^1$ are like terms by definition and consequently giving the reader an impression that $5x^1$ and $3x^1$ are terms themselves.



- A degree 1 indeterminate does not display the multiplicative identity in the exponent when its in canonical form.

Example 8.1 – id:20141121-093747.

Express $13x^0$ in canonical form.



Solution:

8.2 Simplifying Degree 1 Univariate Monomials

Example 8.2 – id:20141120-202042.

Express $5x^1$ in canonical form

(S) _____

Solution:

$$5x \quad \text{MId}(4.9b)$$

■

Example 8.3 – id:20141121-093439.

Express $7x^1 + 5$ in canonical form.

(S) _____

Solution:

$$7x + 5 \quad \text{MId}(4.9b)$$

■

If the constant monomial is 0, the additive identity, then the canonical form of a degree 1 univariate polynomial is a degree 1 monomial.

Example 8.4 – id:20141120-203846.

Simplify by expanding $6x + 7x$

(S) _____

Solution:

Notice that the indeterminate of each monomial is of degree 1; however, the exponent 1 is not shown. The monomials $6x$ and $7x$ have a like term of x .

$$\begin{array}{ll} (6 + 7)x & \text{DPF}(4.3a) \\ 13x & \text{OOA}(2.1) \end{array}$$

Notice that the sum of two monomials that have like terms can be found by adding the coefficients of the monomials. The distributive property in the factoring direction provides some insight to why we can add the coefficients of monomials that have like terms.

S**Less Steps Solution:**

$$13x \quad \text{OOA(2.1)}$$

■

Example 8.5 – id:20141027-075159.Simplify by expanding $7 \text{ cm} + 8 \text{ cm}$ **S****Solution:**

$$(7 + 8) \text{ cm} \quad \text{DPF(4.3a)}$$

$$15 \text{ cm} \quad \text{OOA(2.1)}$$

■

R

Remember, if a monomial does not have a coefficient factor, then it's implied that the coefficient factor is 1, the multiplicative identity, and consequently its not explicitly shown.

Example 8.6 – id:20141121-185558.Simplify by expanding $x + 5x$ **S****Solution:**

It can be useful when simplifying expressions to make the multiplicative identity (MId) factor explicit.

$$\begin{array}{ll} 1x + 5x & \text{MId(4.9a)} \\ (1 + 5)x & \text{DPF(4.3a)} \\ 6x & \text{OOA(2.1)} \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll} 1x + 5x & \text{MId(4.9a)} \\ 6x & \text{OOA(2.1)} \end{array}$$

As one becomes more experienced, there is no reason to make the multiplicative identity coefficient explicit.

(S) _____

Less Steps Solution:

$$6x \quad \text{OOA(2.1)}$$

■

Example 8.7 – id:20141121-190857.

Simplify by expanding $8x - 6x$

(S) _____

Solution:

$$\begin{array}{ll} 8x + -6x & \text{DOS(4.12a)} \\ (8 + -6)x & \text{DPF(4.3a)} \\ 2x & \text{OOA(2.1)} \end{array}$$

■

Less Steps Solution:

$$\begin{array}{ll} 8x + -6x & \text{DOS(4.12a)} \\ 2x & \text{OOA(2.1)} \end{array}$$

■

Example 8.8 – id:20141121-193636.

Simplify by expanding $3x - 5x$

(S) _____

Solution:

$$\begin{array}{ll} 3x + -5x & \text{DOS(4.12a)} \\ (3 + -5)x & \text{DPF(4.3a)} \\ -2x & \text{OOA(2.1)} \\ -2x & \text{ONeg(3.1b)} \end{array}$$

■

Less Steps Solution:

$$\begin{array}{ll} 3x + -5x & \text{DOS(4.12a)} \\ -2x & \text{OOA(2.1)} \\ -2x & \text{ONeg(3.1b)} \end{array}$$

(S)

Less Steps Solution:

$$-2x \quad \text{OOA(2.1)}$$

■

Example 8.9 – id:20141106-150622.Simplify by expanding $13x - x$

(S)

Solution:

$$\begin{array}{ll} 13x - 1x & \text{MId(4.9a)} \\ 13x + -1x & \text{DOS(4.12a)} \\ (13 + -1)x & \text{DPF(4.3a)} \\ 12x & \text{OOA(2.1)} \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll} 13x + -x & \text{DOS(4.12a)} \\ 12x & \text{OOA(2.1)} \end{array}$$

■

It is possible for a univariate monomial to have more than two terms in its non-canonical form. The associative property of addition will be used to help simplify these expressions.

Example 8.10 – id:20141121-184652.Simplify by expanding $3x + 7x + 8x$

(S)

$(3x + 7x) + 8x$	APA(4.2a)
$(3 + 7)x + 8x$	DPF(4.3a)
$10x + 8x$	OOA(2.1)
$(10 + 8)x$	DPF(4.3a)
$18x$	OOA(2.1)

S**Less Steps Solution:**

$(3x + 7x) + 8x$	APA(4.2a)
$10x + 8x$	OOA(2.1)
$18x$	OOA(2.1)

You might have noticed that this expression could be simplified in one step by adding the coefficient of the three monomials $3x$, $7x$ and $8x$, which have the like term x .

S**Less Steps Solution:**

$18x$	OOA(2.1)
-------	----------

Example 8.11 – id:20141106-152020.

Simplify by expanding $4x - 2x - x$

S**Solution:**

$4x - 2x - 1x$	MId(4.9a)
$4x + -2x + -1x$	DOS(4.12a)
$(4 + -2)x + -1x$	DPF(4.3a)
$2x + -1x$	OOA(2.1)
$(2 + -1)x$	DPF(4.3a)
$1x$	OOA(2.1)
x	MId(4.9b)

S

Less Steps Solution:

$$\begin{array}{ll} 4x + -2x + -x & \text{DOS(4.12a)} \\ x & \text{OOA(2.1)} \end{array}$$

■

Example 8.12 – id:20141108-194431.Simplify by expanding $-3 \cdot 7x - 2x \cdot 4$

(S)

Solution:

$$\begin{array}{ll} -3 \cdot 7x - 2x \cdot 4 & \text{ONeg(3.1a)} \\ -3 \cdot 7x + -2x \cdot 4 & \text{DOS(4.12a)} \\ -3 \cdot 7 \cdot x + -2 \cdot x \cdot 4 & \text{JTC(3.5)} \\ -3 \cdot 7 \cdot x + -2 \cdot 4 \cdot x & \text{CPM(4.6)} \\ (-3 \cdot 7) \cdot x + (-2 \cdot 4) \cdot x & \text{APM(4.7a)} \\ -21 \cdot x + -8 \cdot x & \text{OOM(2.3)} \\ -21x + -8x & \text{CTJ(3.6)} \\ (-21 + -8)x & \text{DPF(4.3a)} \\ -29x & \text{OOA(2.1)} \\ -29x & \text{ONeg(3.1b)} \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll} -3 \cdot 7x + -2x \cdot 4 & \text{DOS(4.12a)} \\ -3 \cdot 7 \cdot x + -2 \cdot 4 \cdot x & \text{CPM(4.6)} \\ -21x + -8x & \text{OOM(2.3)} \\ -29x & \text{OOA(2.1)} \end{array}$$

■

Example 8.13 – id:20141108-194156.Simplify by expanding $3 \cdot 5x + 3x \cdot 4$

(S)

Solution:

$3 \cdot 5 \cdot x + 3 \cdot x \cdot 4$	JTC(3.5)
$3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x$	CPM(4.6)
$(3 \cdot 5) \cdot x + (3 \cdot 4) \cdot x$	APM(4.7a)
$15 \cdot x + 12 \cdot x$	OOM(2.3)
$15x + 12x$	CTJ(3.6)
$(15 + 12)x$	DPF(4.3a)
$27x$	OOA(2.1)

S**Less Steps Solution:**

$3 \cdot 5 \cdot x + 3 \cdot 4 \cdot x$	CPM(4.6)
$15x + 12x$	OOM(2.3)
$27x$	OOA(2.1)

Example 8.14 – id:20141108-173613.Simplify by expanding $8x \cdot 5$ **S****Solution:**

$8 \cdot x \cdot 5$	JTC(3.5)
$8 \cdot 5 \cdot x$	CPM(4.6)
$(8 \cdot 5) \cdot x$	APM(4.7a)
$40 \cdot x$	OOM(2.3)
$40x$	CTJ(3.6)

8.3 Simplifying Degree 2 Univariate Monomials

Example 8.15 – id:20141120-202842.Express by expanding $1x^2$ in canonical form.**S**

Solution:

$$x^2 \quad \text{MId(4.9b)}$$

■

Example 8.16 – id:20141106-151138.Simplify by expanding $4x^2 + 12x^2$

(S) _____

Solution:

$$\begin{aligned} (4 + 12)x^2 & \quad \text{DPF(4.3a)} \\ 16x^2 & \quad \text{OOA(2.1)} \end{aligned}$$

(S) _____

Less Steps Solution:

$$16x^2 \quad \text{OOA(2.1)}$$

■

Example 8.17 – id:20141106-154547.Simplify by expanding $x^2 - x + x^2 + x$

(S) _____

Solution:

$$\begin{aligned} 1x^2 - 1x + 1x^2 + 1x & \quad \text{MId(4.9a)} \\ 1x^2 + \neg 1x + 1x^2 + 1x & \quad \text{DOS(4.12a)} \\ 1x^2 + 1x^2 + \neg 1x + 1x & \quad \text{CPA(4.1)} \\ (1 + 1)x^2 + (\neg 1 + 1)x & \quad \text{DPF(4.3a)} \\ 2x^2 + 0x & \quad \text{OOA(2.1)} \\ 2x^2 & \quad \text{MId(4.9b)} \end{aligned}$$

(S) _____

■

Less Steps Solution:

$$\begin{array}{ll} x^2 + \neg x + x^2 + x & \text{DOS(4.12a)} \\ x^2 + x^2 + \neg x + x & \text{CPA(4.1)} \\ 2x^2 & \text{OOA(2.1)} \end{array}$$

■

Example 8.18 – id:20141108-194709.Simplify by expanding $-2x^4x - x \cdot -x^3$

(S)

Solution:

$$\begin{array}{ll} -2x^4x - 1x \cdot 1x^3 & \text{MId(4.9a)} \\ -2x^4x - 1x \cdot 1x^3 & \text{ONeg(3.1a)} \\ -2 \cdot x \cdot 4 \cdot x + \neg 1 \cdot x \cdot 1 \cdot x \cdot 3 & \text{JTC(3.5)} \\ -2 \cdot 4 \cdot x \cdot x + \neg 1 \cdot \neg 1 \cdot 3 \cdot x \cdot x & \text{CPM(4.6)} \\ -2 \cdot 4 \cdot x^2 + \neg 1 \cdot \neg 1 \cdot 3 \cdot x^2 & \text{PrCBPo(5.3a)} \\ (-2 \cdot 4) \cdot x^2 + (\neg 1 \cdot \neg 1 \cdot 3)x^2 & \text{APM(4.7a)} \\ -8x^2 + 3x^2 & \text{OOM(2.3)} \\ (-8 + 3)x^2 & \text{DPF(4.3a)} \\ -5x^2 & \text{OOA(2.1)} \\ -5x^2 & \text{ONeg(3.1b)} \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll} -2x^4x + \neg x \cdot -x^3 & \text{DOS(4.12a)} \\ -2 \cdot 4 \cdot x \cdot x + 3 \cdot \neg x \cdot \neg x & \text{CPM(4.6)} \\ -2 \cdot 4 \cdot x^2 + 3 \cdot x^2 & \text{PrCBPo(5.3a)} \\ -8x^2 + 3x^2 & \text{OOM(2.3)} \\ -5x^2 & \text{OOA(2.1)} \end{array}$$

■

Example 8.19 – id:20141108-191616.Simplify by expanding $-5x \cdot 4x$

(S)

Solution:

$-5x \cdot 4x$	ONeg(3.1a)
$-5 \cdot x \cdot 4 \cdot x$	JTC(3.5)
$-5 \cdot 4 \cdot x \cdot x$	CPM(4.6)
$-5 \cdot 4 \cdot x^2$	PrCBPo(5.3a)
$(-5 \cdot 4) \cdot x^2$	APM(4.7a)
$-20 \cdot x^2$	OOM(2.3)
$-20x^2$	CTJ(3.6)
$-20x^2$	ONeg(3.1b)

(S)

Less Steps Solution:

$-5 \cdot 4 \cdot x \cdot x$	CPM(4.6)
$-5 \cdot 4 \cdot x^2$	PrCBPo(5.3a)
$-20x^2$	OOM(2.3)

■

8.4 Simplifying Degree 3 Univariate Monomials

8.5 Simplifying Degree 1 Univariate Binomials

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + \textcolor{red}{C_1 x^1} + \underbrace{C_0}_{\textcolor{red}{C_0 x^0}}$$

Degree 1 univariate polynomial expressions can be expressed with at most two different terms and consequently this expression in its canonical form has at most two monomial summands – called a binomial.

Example 8.20 – id:20141109-090809.

Simplify by expanding $5(x + 4)$

(S)

Solution:

$5(1x + 4)$	MId(4.9a)
$5 \cdot 1x + 5 \cdot 5$	DPF(4.3a)
$5 \cdot 1 \cdot x + 5 \cdot 5$	JTC(3.5)
$5 \cdot x + 25$	OOM(2.3)
$5x + 25$	CTJ(3.6)

S**Less Steps Solution:**

$$5x + 20$$

DPE(4.8a)

■

Example 8.21 – id:20141109-091015.Simplify by expanding $5(3x - 9)$ **S****Solution:**

$$5(3x + -9)$$

DOS(4.12a)

$$5 \cdot 3x + 5 \cdot -9$$

DPE(4.8a)

$$5 \cdot 3 \cdot x + 5 \cdot -9$$

JTC(3.5)

$$15 \cdot x + -45$$

OOM(2.3)

$$15x + -45$$

CTJ(3.6)

$$15x - 45$$

DOS(4.12b)

S**Less Steps Solution:**

$$5(3x + -9)$$

DOS(4.12a)

$$15x + -40$$

DPE(4.8a)

$$15x - 40$$

DOS(4.12b)

■

Example 8.22 – id:20141109-092448.Simplify by expanding $-(5x + 7)$ **S**

Solution:

$\neg 1(5x + 7)$	MId(4.9a)
$\neg 1 \cdot 5x + \neg 1 \cdot 7$	DPE(4.8a)
$\neg 1 \cdot 5 \cdot x + \neg 1 \cdot 7$	JTC(3.5)
$\neg 5 \cdot x + \neg 7$	OOM(2.3)
$\neg 5x + \neg 7$	CTJ(3.6)
$\neg 5x - 7$	DOS(4.12b)
$-5x - 7$	ONeg(3.1b)

(S)

Less Steps Solution:

$$-5x - 7 \quad \text{DPE(4.8a)}$$

Example 8.23 – id:20141109-092651.

Simplify by expanding $-13(7x - 9)$

(S)

Solution:

$\neg 13(7x + \neg 9)$	DOS(4.12a)
$\neg 13 \cdot 7x + \neg 13 \cdot \neg 9$	DPE(4.8a)
$\neg 13 \cdot 7 \cdot x + \neg 13 \cdot \neg 9$	JTC(3.5)
$\neg 91 \cdot x + 117$	OOM(2.3)
$\neg 91x + 117$	CTJ(3.6)
$-91x + 117$	ONeg(3.1b)

(S)

Less Steps Solution:

$$\begin{aligned} -13(7x + \neg 9) & \quad \text{DOS(4.12a)} \\ -91x + 117 & \quad \text{DPE(4.8a)} \end{aligned}$$

Example 8.24 – id:20141109-092910.

Simplify by expanding $a(x + b)$, where $a, b \in \mathbb{Z}$

(S)

Solution:

$$\begin{array}{ll} a(1x + b) & \text{MId(4.9a)} \\ a \cdot 1x + a \cdot b & \text{DPE(4.8a)} \\ a \cdot 1 \cdot x + a \cdot b & \text{JTC(3.5)} \\ 1 \cdot a \cdot x + a \cdot b & \text{CPM(4.6)} \\ 1ax + ab & \text{JTC(3.5)} \\ ax + ab & \text{MId(4.9b)} \end{array}$$

(S)

Less Steps Solution:

$$ax + ab \quad \text{DPE(4.8a)}$$

■

Example 8.25 – id:20141109-093220.

Simplify by expanding $5(x + 2) + 4$

(S)

Solution:

$$\begin{array}{ll} 5(1x + 2) + 4 & \text{MId(4.9a)} \\ 5 \cdot 1x + 5 \cdot 2 + 4 & \text{DPE(4.8a)} \\ 5 \cdot 1 \cdot x + 5 \cdot 2 + 4 & \text{JTC(3.5)} \\ 5 \cdot x + 10 + 4 & \text{OOM(2.3)} \\ 5x + 10 + 4 & \text{CTJ(3.6)} \\ 5x + 14 & \text{OOA(2.1)} \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll} 5x + 10 + 4 & \text{DPE(4.8a)} \\ 5x + 14 & \text{OOA(2.1)} \end{array}$$

■

Example 8.26 – id:20141109-093419.

Simplify by expanding $7x + 5(4x + 8)$

(S) _____

Solution:

$$\begin{array}{ll}
 7x + 5 \cdot 4x + 5 \cdot 8 & \text{DPE(4.8a)} \\
 7 \cdot x + 5 \cdot 4 \cdot x + 5 \cdot 8 & \text{JTC(3.5)} \\
 7 \cdot x + 20 \cdot x + 40 & \text{OOM(2.3)} \\
 7x + 20x + 40 & \text{CTJ(3.6)} \\
 (7 + 20)x + 40 & \text{DPF(4.3a)} \\
 27x + 40 & \text{OOA(2.1)}
 \end{array}$$

■

Example 8.27 – id:20141109-094928.

Simplify by expanding $4(3x + 4) + x + 6$

(S) _____

Solution:

$$\begin{array}{ll}
 4(3x + 4) + 1x + 6 & \text{MId(4.9a)} \\
 4 \cdot 3x + 4 \cdot 4 + 1x + 6 & \text{DPE(4.8a)} \\
 4 \cdot 3 \cdot x + 4 \cdot 4 + 1 \cdot x + 6 & \text{JTC(3.5)} \\
 12 \cdot x + 16 + 1 \cdot x + 6 & \text{OOM(2.3)} \\
 12x + 16 + 1x + 6 & \text{CTJ(3.6)} \\
 12 + 1x + 16 + 6 & \text{CPA(4.1)} \\
 (12 + 1)x + 16 + 6 & \text{DPF(4.3a)} \\
 13x + 22 & \text{OOA(2.1)}
 \end{array}$$

(S) _____

Less Steps Solution:

$$\begin{array}{ll}
 12x + 16 + x + 6 & \text{DPE(4.8a)} \\
 12x + x + 16 + 6 & \text{CPA(4.1)} \\
 13x + 22 & \text{OOA(2.1)}
 \end{array}$$

■

Example 8.28 – id:20141109-095151.

Simplify by expanding $5(x - 4) + 3x - 5$

(S)

Solution:

$5(1x - 4) + 3x - 5$	MId(4.9a)
$5(1x + \neg 4) + 3x + \neg 5$	DOS(4.12a)
$5 \cdot 1x + 5 \cdot \neg 4 + 3x + \neg 5$	DPE(4.8a)
$5 \cdot 1 \cdot x + 5 \cdot \neg 4 + 3 \cdot x + \neg 5$	JTC(3.5)
$5 \cdot x + \neg 20 + 3 \cdot x + \neg 5$	OOM(2.3)
$5x + \neg 20 + 3x + \neg 5$	JTC(3.5)
$5x + 3x + \neg 20 + \neg 5$	CPA(4.1)
$(5 + 3)x + \neg 20 + \neg 5$	DPF(4.3a)
$8x + \neg 25$	OOA(2.1)
$8x - 25$	DOS(4.12b)

(S)

Less Steps Solution:

$5(x + \neg 4) + 3x + \neg 5$	DOS(4.12a)
$5x + \neg 20 + 3x + \neg 5$	DPE(4.8a)
$5x + 3x + \neg 20 + \neg 5$	CPA(4.1)
$8x + \neg 25$	OOA(2.1)
$8x - 25$	DOS(4.12b)

Example 8.29 – id:20141109-095536.

Simplify by expanding $8x - 5 - 4(x - 3)$

(S)

Solution:

$8x - 5 - 4(1x - 3)$	MId(4.9a)
$8x + -5 + -4(1x + -3)$	DOS(4.12a)
$8x + -5 + -4 \cdot 1x + -4 \cdot -3$	DPE(4.8a)
$8 \cdot x + -5 + -4 \cdot 1 \cdot x + -4 \cdot -3$	JTC(3.5)
$8 \cdot x + -5 + -4 \cdot x + 12$	OOM(2.3)
$8x + -5 + -4x + 12$	CTJ(3.6)
$8x + -4x + -5 + 12$	CPA(4.1)
$(8 + -4)x + -5 + 12$	DPF(4.3a)
$4x + 7$	OOA(2.1)

S

Less Steps Solution:

$8x + -5 + -4(x + -3)$	DOS(4.12a)
$8x + -5 + -4x + 12$	DPE(4.8a)
$8x + -4x + -5 + 12$	CPA(4.1)
$4x + 7$	OOA(2.1)

Example 8.30 – id:20141109-095842.

Simplify by expanding $5(x + 3) + 3(x + 2)$

S

Solution:

$5 \cdot x + 5 \cdot 3 + 3 \cdot x + 3 \cdot 2$	DPE(4.8a)
$5 \cdot x + 15 + 3 \cdot x + 6$	OOM(2.3)
$5x + 15 + 3x + 6$	CTJ(3.6)
$5x + 3x + 15 + 6$	CPA(4.1)
$(3 + 5)x + 15 + 6$	DPF(4.3a)
$8x + 21$	OOA(2.1)

S

Less Steps Solution:

$$\begin{array}{ll} 5x + 15 + 3x + 6 & \text{DPE(4.8a)} \\ 5x + 3x + 15 + 6 & \text{CPA(4.1)} \\ 8x + 21 & \text{OOA(2.1)} \end{array}$$

■

8.6 Simplifying Degree 2 Univariate Binomials**Example 8.31 – id:20141106-152339.**

Simplify by expanding $3x^2 + 2x + 5x^2 + 4x$

(S) _____

Solution:

$$\begin{array}{ll} 3x^2 + 5x^2 + 2x + 4x & \text{CPA(4.1)} \\ (3 + 5)x^2 + (2 + 4)x & \text{DPF(4.3a)} \\ 8x^2 + 6x & \text{OOA(2.1)} \end{array}$$

If needed we could continue and express it in the simplified factored form using the distributive property

$$(4x + 3)2x \quad \text{DPF(4.3a)}$$

(S) _____

Less Steps Solution:

$$\begin{array}{ll} 3x^2 + 5x^2 + 2x + 4x & \text{CPA(4.1)} \\ 8x^2 + 6x & \text{OOA(2.1)} \end{array}$$

■

Example 8.32 – id:20141107-121834.

Simplify by expanding $(\sqrt{9 - x^2})^2$

(S) _____

Solution:

$$\begin{aligned}
 & (\sqrt{9 - 1x^2})^2 && \text{MId(4.9a)} \\
 & (\sqrt{9 + -1x^2})^2 && \text{DOS(4.12a)} \\
 & [(9 + -x^2)^{\frac{1}{2}}]^2 && \text{RTPo(3.12)} \\
 & 9 + -1x^2 && \text{PoPo(5.1a)} \\
 & -1x^2 + 9 && \text{CPA(4.1)} \\
 & -x^2 + 9 && \text{MId(4.9a)} \\
 & -x^2 + 9 && \text{ONeg(3.1b)}
 \end{aligned}$$

(S)**Less Steps Solution:**

$$9 - x^2 \quad \text{PoPo(5.1a)}$$

It might be easier to view this using a variable substitution for the radicand, $9 - x^2$. Let $k = 9 + -1x^2$.

$$\begin{aligned}
 & (\sqrt{k})^2 && \text{MId(4.9a)} \\
 & (\sqrt{k})^2 && \text{DOS(4.12a)} \\
 & [(k)^{\frac{1}{2}}]^2 && \text{RTPo(3.12)} \\
 & k && \text{PoPo(5.1a)} \\
 & 9 + -1x^2 && \text{CPA(4.1)} \\
 & -1x^2 + 9 && \text{CPA(4.1)} \\
 & -x^2 + 9 && \text{MId(4.9a)} \\
 & -x^2 + 9 && \text{ONeg(3.1b)}
 \end{aligned}$$

(D)

Dependencies:example ??-20141105-144223

Example 8.33 – id:20141209-145211.Simplify by expanding $2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0$ **(S)**

Solution:

$$\begin{aligned}
 & 2x \cdot 2x + 2x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0 && \text{DPE(4.8a)} \\
 & 2 \cdot x \cdot 2 \cdot x + 2 \cdot x \cdot 4 + x^2 \cdot 2 \cdot 1 + 0 && \text{JTC(3.5)} \\
 & 2 \cdot 2 \cdot x \cdot x + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0 && \text{CPM(4.6)} \\
 & 2 \cdot 2 \cdot x^2 + 2 \cdot 4 \cdot x + 2 \cdot 1 \cdot x^2 + 0 && \text{PrCBPo(5.3a)} \\
 & 4 \cdot x^2 + 8 \cdot x + 2 \cdot x^2 + 0 && \text{OOM(2.3)} \\
 & 4x^2 + 8x + 2x^2 + 0 && \text{CTJ(3.6)} \\
 & 4x^2 + 2x^2 + 8x && \text{APA(4.2a)} \\
 & (4 + 2)x^2 + 8x && \text{DPF(4.3b)} \\
 & 6x^2 + 8x && \text{OOA(2.1)}
 \end{aligned}$$

D

Dependencies:example 15.7-20141209-144203



8.7 Simplifying Degree 2 Univariate Trinomials

Example 8.34 – id:20141109-133008.Simplify by expanding $(x + 5)(x - 8)$ **S****Solution:**

$$\begin{aligned}
 & (1x + 5)(1x - 8) && \text{MId(4.9a)} \\
 & (1x + 5)(1x + \neg 8) && \text{DOS(4.12a)} \\
 & 1x(1x + \neg 8) + 5(1x + \neg 8) && \text{DPE(4.8b)} \\
 & 1x \cdot 1x + 1x \cdot \neg 8 + 5 \cdot 1x + 5 \cdot \neg 8 && \text{DPE(4.8a)} \\
 & 1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot \neg 8 + 5 \cdot 1 \cdot x + 5 \cdot \neg 8 && \text{JTC(3.5)} \\
 & 1 \cdot 1 \cdot x \cdot x + \neg 8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + \neg 8 \cdot 5 && \text{CPM(4.6)} \\
 & 1 \cdot 1 \cdot x^2 + \neg 8 \cdot 1 \cdot x + 1 \cdot 5 \cdot x + \neg 8 \cdot 5 && \text{PrCBPo(5.3a)} \\
 & 1 \cdot x^2 + \neg 8 \cdot x + 5 \cdot x + \neg 40 && \text{OOM(2.3)} \\
 & 1x^2 + \neg 8x + 5x + \neg 40 && \text{CTJ(3.6)} \\
 & 1x^2 + \neg 3x + \neg 40 && \text{OOA(2.1)} \\
 & 1x^2 - 3x - 40 && \text{DOS(4.12b)} \\
 & x^2 - 3x - 40 && \text{MId(4.9a)}
 \end{aligned}$$

S

Less Steps Solution:

$$\begin{array}{ll}
 (x + 5)(x + -8) & \text{DOS(4.12a)} \\
 x(x + -8) + 5(x + -8) & \text{DPE(4.8b)} \\
 x^2 + -8x + 5x + -40 & \text{DPE(4.8a)} \\
 x^2 - 3x - 40 & \text{OOA(2.1)}
 \end{array}$$

Example 8.35 – id:20141109-133316.

Simplify by expanding $(x + a)(x + b)$, where $a, b \in \mathbb{Z}$

(S)

Solution:

$$\begin{array}{ll}
 (1x + a)(1x + b) & \text{MId(4.9a)} \\
 1x(1x + b) + a(1x + b) & \text{DPE(4.8b)} \\
 1x \cdot 1x + 1x \cdot b + a \cdot 1x + a \cdot b & \text{DPE(4.8a)} \\
 1 \cdot x \cdot 1 \cdot x + 1 \cdot x \cdot b + a \cdot 1 \cdot x + a \cdot b & \text{JTC(3.5)} \\
 1 \cdot 1 \cdot x \cdot x + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{CPM(4.6)} \\
 1 \cdot 1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{PrCBPo(5.3a)} \\
 1 \cdot x^2 + 1 \cdot b \cdot x + 1 \cdot a \cdot x + a \cdot b & \text{OOM(2.3)} \\
 1x^2 + 1bx + 1ax + ab & \text{CTJ(3.6)} \\
 1x^2 + (1b + 1a)x + ab & \text{DPF(4.3a)} \\
 x^2 + (b + a)x + ab & \text{MId(4.9b)}
 \end{array}$$

(S)

Less Steps Solution:

$$\begin{array}{ll}
 x(x + b) + a(x + b) & \text{DPE(4.8b)} \\
 x^2 + (b + a)x + ab & \text{DPE(4.8a)}
 \end{array}$$

Example 8.36 – id:20141109-140659.

Simplify by expanding $(2x + 3)(5x + 13)$

(S)

Solution:

$$\begin{aligned}
 & 2x(5x + 13) + 3(5x + 13) && \text{DPE(4.8b)} \\
 & 2x \cdot 5x + 2x \cdot 13 + 3 \cdot 5x + 3 \cdot 13 && \text{DPE(4.8a)} \\
 & 2 \cdot x \cdot 5 \cdot x + 2 \cdot x \cdot 13 + 3 \cdot 5 \cdot x + 3 \cdot 13 && \text{JTC(3.5)} \\
 & 2 \cdot 5 \cdot x \cdot x + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13 && \text{CPM(4.6)} \\
 & 2 \cdot 5 \cdot x^2 + 2 \cdot 13 \cdot x + 5 \cdot 3 \cdot x + 3 \cdot 13 && \text{PrCBPo(5.3a)} \\
 & 10 \cdot x^2 + 26 \cdot x + 16 \cdot x + 39 && \text{OOM(2.3)} \\
 & 10x^2 + 26x + 15x + 39 && \text{CTJ(3.6)} \\
 & 10x^2 + 41x + 39 && \text{OOA(2.1)}
 \end{aligned}$$

(S)**Less Steps Solution:**

$$\begin{aligned}
 & 2x(5x + 13) + 3(5x + 13) && \text{DPE(4.8b)} \\
 & 10x^2 + 26x + 15x + 39 && \text{DPE(4.8a)} \\
 & 10x^2 + 41x + 39 && \text{OOA(2.1)}
 \end{aligned}$$

Example 8.37 – id:20141109-141019.Simplify by expanding $(-3x - 5)(7x + 8)$ **(S)****Solution:**

$$\begin{aligned}
 & (-3x - 5)(7x + 8) && \text{ONeg(3.1a)} \\
 & (-3x + -5)(7x + 8) && \text{DOS(4.12a)} \\
 & -3x(7x + 8) + -5(7x + 8) && \text{DPE(4.8b)} \\
 & -3x \cdot 7x + -3x \cdot 8 + -5 \cdot 7x + -5 \cdot 8 && \text{DPE(4.8a)} \\
 & -3 \cdot x \cdot 7 \cdot x + -3 \cdot x \cdot 8 + -5 \cdot 7 \cdot x + -5 \cdot 8 && \text{JTC(3.5)} \\
 & -3 \cdot 7 \cdot x \cdot x + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 && \text{CPM(4.6)} \\
 & -3 \cdot 7 \cdot x^2 + -3 \cdot 8 \cdot x + -5 \cdot 7 \cdot x + -5 \cdot 8 && \text{PrCBPo(5.3a)} \\
 & -21 \cdot x^2 + -24 \cdot x + -35 \cdot x + -40 && \text{OOM(2.3)} \\
 & -21x^2 + -24x + -35x + -40 && \text{CTJ(3.6)} \\
 & -21x^2 + -59x + -40 && \text{OOA(2.1)} \\
 & -21x^2 - 59x - 40 && \text{DOS(4.12b)} \\
 & -21x^2 - 59x - 40 && \text{ONeg(3.1b)}
 \end{aligned}$$

S**Less Steps Solution:**

$$\begin{array}{ll}
 (-3x + -5)(7x + 8) & \text{DOS(4.12a)} \\
 -3x(7x + 8) + -5(7x + 8) & \text{DPE(4.8b)} \\
 -21x^2 + -24x + -35x + -40 & \text{CTJ(4.8a)} \\
 -21x^2 - 59x - 40 & \text{OOA(2.1)}
 \end{array}$$

■

Example 8.38 – id:20141109-141347.Simplify by expanding $(ax + b)(cx + d)$, where $a, b, c, d \in \mathbb{Z}$ **S****Solution:**

$$\begin{array}{ll}
 ax(cx + d) + b(cx + d) & \text{DPE(4.8b)} \\
 ax \cdot cx + ax \cdot d + b \cdot cx + b \cdot d & \text{DPE(4.8a)} \\
 a \cdot x \cdot c \cdot x + a \cdot x \cdot d + b \cdot c \cdot x + b \cdot d & \text{JTC(3.5)} \\
 a \cdot c \cdot x \cdot x + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d & \text{CPM(4.6)} \\
 a \cdot c \cdot x^2 + a \cdot d \cdot x + b \cdot c \cdot x + b \cdot d & \text{PrCBPo(5.3a)} \\
 acx^2 + adx + bcx + bd & \text{CTJ(3.6)} \\
 acx^2 + (ad + bc)x + bd & \text{DPF(4.3a)}
 \end{array}$$

S**Less Steps Solution:**

$$\begin{array}{ll}
 ax(cx + d) + b(cx + d) & \text{DPE(4.8b)} \\
 acx^2 + (ad + bc)x + bd & \text{DPE(4.8a)}
 \end{array}$$

■

Example 8.39 – id:20141105-161225.Simplify by expanding $\left(2 - \frac{x}{2}\right)^2$.**S**

Solution:

$$\begin{aligned}
 & \left(2 - \frac{1x}{2}\right)^2 && \text{MId(4.9a)} \\
 & \left(2 + -\frac{1x}{2}\right)^2 && \text{DOS(4.12a)} \\
 & \left(2 + -\frac{1x}{2}\right)\left(2 + -\frac{1x}{2}\right) && \text{PoTF(3.9)} \\
 & 2\left(2 + -\frac{1x}{2}\right) + -\frac{1x}{2}\left(2 + -\frac{1x}{2}\right) && \text{DPE(4.8b)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1x}{2} + -\frac{1x}{2} \cdot 2 + -\frac{1x}{2} \cdot -\frac{1x}{2} && \text{DPE(4.8a)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot x \cdot 2 + -\frac{1}{2} \cdot x \cdot -\frac{1}{2} \cdot x && \text{JTC(3.5)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x \cdot x && \text{CPM(4.6)} \\
 & 2 \cdot 2 + 2 \cdot -\frac{1}{2} \cdot x + -\frac{1}{2} \cdot 2 \cdot x + -\frac{1}{2} \cdot -\frac{1}{2} \cdot x^2 && \text{PrCBPo(5.3a)} \\
 & 4 + -1 \cdot x + -1 \cdot x + \frac{1}{4} \cdot x^2 && \text{OOM(2.3)} \\
 & 4 + -1x + -1x + \frac{1}{4}x^2 && \text{CTJ(3.6)} \\
 & \frac{1}{4}x^2 + -1x + -1x + 4 && \text{CPA(4.1)} \\
 & \frac{1}{4}x^2 + -2x + 4 && \text{OOA(2.1)} \\
 & \frac{1}{4}x^2 - 2x + 4 && \text{DOS(4.12b)}
 \end{aligned}$$

(S)**Less Steps Solution:**

$$\begin{aligned}
 & 7x + 20x + 40 && \text{DPE(4.8a)} \\
 & 27x + 40 && \text{OOA(2.1)}
 \end{aligned}$$

8.8 Simplifying Degree n Univariate Polynomials

Definition 8.8.1 – Univariate Polynomial Expression.

$$\sum_{k=0}^n C_k x^n = C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0} \quad (8.1)$$

A univariate polynomial in an indeterminate x is an expression made up of one or more summands of the form $C_k x^k$, which are called monomials. The first factor of each monomial, C_k , is a numerical factor called the **coefficient** where $C_k \in$. The second factor

of each monomial, x^k , is an indeterminate raised to a non-negative integer power i .

Polynomial [3]

Definition 8.8.2 – Degree of the Univariate Polynomial.

$$C_n x^n + C_{n-1} x^{n-1} + \cdots + C_k x^k + \cdots + C_2 x^2 + C_1 x^1 + \underbrace{C_0}_{C_0 x^0}$$

The degree of the univariate polynomial is determined by the monomial with the largest degree of the indeterminate.

9. Simplifying Univariate Monomials

9.1

10. Factoring Univariate Trinomials

Example 10.1 – id:20151015-184212.

Simplify by factoring $x^2 + 7x + 12$

(S) _____


Solution:

Using the distributive property organizer

The factors of $(x^2 + 7x + 12)$ are $(x + 3)(x + 4)$

$$1x^2 + 7x + 12$$

MId(4.9a)



$$(1x + 3)(1x + 4)$$

$$(x + 3)(x + 4)$$

MId(4.9b)

Example 10.2 – id:20151016-063338.

Simplify by factoring $x^2 + x - 2$

(S) _____

Solution:

Using the distributive property organizer


The factors of $(x^2 + x - 2)$ are $(\textcolor{red}{x} + \textcolor{blue}{2})(\textcolor{red}{x} - \textcolor{blue}{1})$

$$1x^2 + 1x - 2$$

AId(4.4a)

$$1x^2 + 1x + \neg 2$$

DOS(4.12a)



$$(1x + 2)(1x + \neg 1)$$

DOS(4.12b)

$$(x + 2)(x - 1)$$

MId(4.9b)

■

11. Solving Linear Equations

11.1 Power Inverse

Example 11.1 – id:20141206-102142.

Solve the equation $x + a = b$ for x

(S) _____

Solution:

$$\begin{array}{ll} [x + a] + \neg a = [b] + \neg a & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x + (a + \neg a) = b + \neg a & \text{APA(4.2b)} \\ x + 0 = b + \neg a & \text{OOA(2.1)} \\ x = b + \neg a & \text{AId(4.4b)} \\ x = b - a & \text{DOS(4.12b)} \end{array}$$

■

Example 11.2 – id:20141111-222931.

Solve the equations $x + 8 = 0$

(S) _____

Solution:

$$\begin{array}{ll} [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x + (8 + \neg 8) = 0 + \neg 8 & \text{APA(4.2b)} \\ x + 0 = \neg 8 & \text{OOA(2.1)} \\ x = \neg 8 & \text{AId(4.4b)} \\ x = -8 & \text{ONeg(3.1b)} \end{array}$$

(S) _____

Less Steps Solution:

$$\begin{array}{ll} [x + 8] + \neg 8 = [0] + \neg 8 & \text{SPE(4.15)} + \text{AI(4.5a)} \\ x = -8 & \text{OOA(2.1)} \end{array}$$

D

Dependencies:
example ??-20141111-190212

Example 11.3 – id:20141206-101632.

Solve the equation $x + 4 = 7$

S

Solution:

$$\begin{array}{ll} [x + 4] + \neg 4 = [7] + \neg 4 & \text{SPE(4.15) + AI(4.5a)} \\ x + (4 + \neg 4) = 7 + \neg 4 & \text{APA(4.2b)} \\ x + 0 = 3 & \text{OOA(2.1)} \\ x = 3 & \text{AId(4.4b)} \end{array}$$

S

Less Steps Solution:

$$\begin{array}{ll} [x] + 4 + \neg 4 = [7] + \neg 4 & \text{SPE(4.15) + AI(4.5a)} \\ x = 3 & \text{OOA(2.1)} \end{array}$$

Example 11.4 – id:20141206-101107.

Solve the equation $x - 8 = 15$ for x

S

Solution:

$$\begin{array}{ll} x + \neg 8 = 15 & \text{DOS(4.12a)} \\ [x + \neg 8] + 8 = [15] + 8 & \text{SPE(4.15) + AI(4.5a)} \\ x + (\neg 8 + 8) = 15 + 8 & \text{APA(4.2b)} \\ x + 0 = 23 & \text{OOA(2.1)} \\ x = 23 & \text{AId(4.4b)} \end{array}$$

S

Less Steps Solution:

$$\begin{aligned} [x + \cancel{-8}] + 8 &= [15] + 8 && \text{SPE(4.15) + AI(4.5a)} \\ x &= 23 && \text{OOA(2.1)} \end{aligned}$$

■

Example 11.5 – id:20141206-102404.

Solve the equation $5x = 9$ for x .

(S)

Solution:

$$\begin{aligned} \frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{5} \cdot [5 \cdot x] &= \frac{1}{5} \cdot 9 && \text{JTC(3.5)} \\ \left(\frac{1}{5} \cdot 5\right) \cdot x &= \frac{1}{5} \cdot 9 && \text{APM(4.7b)} \\ 1 \cdot x &= \frac{9}{5} && \text{OOM(2.3)} \\ x &= \frac{9}{5} && \text{MId(4.9b)} \end{aligned}$$

(S)

Less Steps Solution:

$$\begin{aligned} \frac{1}{5}[5x] &= \frac{1}{5}[9] && \text{SPE(4.15) + MI(4.10a)} \\ x &= \frac{9}{5} && \text{OOM(2.3)} \end{aligned}$$

■

Example 11.6 – id:20141206-104404.

Solve the equation $ax = b$ for x .

(S)

Solution:

$$\begin{aligned}\frac{1}{a}[\mathbf{ax}] &= \frac{1}{a}[\mathbf{b}] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{a} \cdot (a \cdot x) &= \frac{1}{a} \cdot b && \text{JTC(3.5)} \\ \left(\frac{1}{a} \cdot a\right) \cdot x &= \frac{1}{a} \cdot b && \text{APM(4.7b)} \\ 1 \cdot x &= \frac{b}{a} && \text{OOM(2.3)} \\ x &= \frac{b}{a} && \text{MId(4.9b)}\end{aligned}$$

■

Example 11.7 – id:20141206-102723.Solve the equation $-2x = 7$ for x

(S)

Solution:

$$\begin{aligned}-2x &= 7 && \text{ONeg(3.1a)} \\ -\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.15) + MI(4.10b)} \\ -\frac{1}{2} \cdot (-2 \cdot x) &= -\frac{1}{2} \cdot 7 && \text{JTC(3.5)} \\ \left(-\frac{1}{2} \cdot -2\right) \cdot x &= -\frac{1}{2} \cdot 7 && \text{APM(4.7b)} \\ 1 \cdot x &= -\frac{7}{2} && \text{OOM(2.3)} \\ 1 \cdot x &= -\frac{7}{2} && \text{ONeg(3.1b)} \\ x &= -\frac{7}{2} && \text{MId(4.9b)}\end{aligned}$$

(S)

Less Steps Solution:

$$\begin{aligned}-\frac{1}{2}[-2x] &= -\frac{1}{2}[7] && \text{SPE(4.15) + MI(4.10b)} \\ x &= -\frac{7}{2} && \text{OOM(2.3)}\end{aligned}$$

■

Example 11.8 – id:20141111-215726.

Solve the equation $2x + 5 = 0$ for x

(S) _____

Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.15) + AI(4.5a)} \\
 2x + (5 + \neg 5) &= 0 + \neg 5 && \text{APA(4.2a)} \\
 2x + 0 &= \neg 5 && \text{OOA(2.1)} \\
 2x &= \neg 5 && \text{AId(4.4a)} \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.15) + MI(4.10a)} \\
 \frac{1}{2} \cdot 2 \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{JTC(3.5)} \\
 \left(\frac{1}{2} \cdot 2\right) \cdot x &= \frac{1}{2} \cdot \neg 5 && \text{APM(4.7a)} \\
 1 \cdot x &= \frac{\neg 5}{2} && \text{OOM(2.3)} \\
 1x &= -\frac{5}{2} && \text{ONeg(3.1b)} \\
 x &= -\frac{5}{2} && \text{MId(4.9b)}
 \end{aligned}$$

(S) _____

Less Steps Solution:

$$\begin{aligned}
 [2x + 5] + \neg 5 &= [0] + \neg 5 && \text{SPE(4.15) + AI(4.5a)} \quad (11.1) \\
 2x &= \neg 5 && \text{OOA(2.1)} \quad (11.2) \\
 \frac{1}{2}[2x] &= \frac{1}{2}[\neg 5] && \text{SPE(4.15) + MI(4.10a)} \quad (11.3) \\
 x &= -\frac{5}{2} && \text{OOM(2.3)} \quad (11.4)
 \end{aligned}$$

(D) _____

Dependencies:

example ??-20141111-192213

■

Example 11.9 – id:20151015-104754.

Solve the equation $2ax + b = 0$ for x .

(S) _____

Solution:

$$\begin{aligned}
 [2ax + b] + \neg b &= [0] + \neg b && \text{AI(4.5a), SPE(4.15)} \\
 2ax + (b + \neg b) &= 0 + \neg b && \text{APA(4.2b)} \\
 2ax + 0 &= 0 + \neg b && \text{OOA(2.1)} \\
 2ax &= \neg b && \text{AId(4.4a)} \\
 \frac{1}{2a} [2ax] &= \frac{1}{2a} [\neg b] && \text{MI(4.5a), SPE(4.15)} \\
 \frac{1}{2a} \cdot (2ax) &= \frac{1}{2a} \cdot (\neg b) && \text{JTC(3.5)} \\
 \left(\frac{1}{2a} \cdot 2a\right) \cdot x &= \frac{1}{2a} \cdot \neg b && \text{APA(4.2a)} \\
 \left(\frac{1}{2a} \cdot \frac{2a}{1}\right) x &= \frac{1}{2a} \cdot \frac{\text{neg } b}{1} && \text{MId(4.9a)} \\
 \frac{2a}{2a} \cdot x &= \frac{\neg b}{2a} && \text{OOM(2.3)} \\
 1 \cdot x &= \frac{\neg b}{2a} x && = \frac{\neg b}{2a} && \text{MId(4.9b)} \\
 x &= -\frac{b}{2a} && \text{ONeg(3.1b)}
 \end{aligned}$$

(D)

Dependencies:example 18.1-20151008-110208

12. Solving Quadratic Equations

12.1 Power Inverse

Example 12.1 – id:20141107-131748.

Solve the equation $2 - x^2 = 0$ for x

(S) _____

Solution:

$$2 - 1x^2 = 0 \quad \text{MId(4.9a)}$$

$$2 + -1x^2 = 0 \quad \text{DOS(4.12a)}$$

$$[2 + -1x^2] + 1x^2 = [0] + 1x^2 \quad \text{SPE(4.15) + AI(4.5a)}$$

$$2 + (-1x^2 + 1x^2) = 0 + 1x^2 \quad \text{APA(4.2a)}$$

$$2 + 0 = 0 + 1x^2 \quad \text{OOA(2.1)}$$

$$2 = 1x^2 \quad \text{AIId(4.4a)}$$

$$2 = x^2 \quad \text{MId(4.9b)}$$

$$\pm [2]^{\frac{1}{2}} = [x^2]^{\frac{1}{2}} \quad \text{PoPo(5.1a)}$$

$$\pm 2^{\frac{1}{2}} = x \quad \text{PoTR(??)}$$

$$\pm \sqrt{2} = x \quad \text{SyPE(4.16a)}$$

Example 12.2 – id:20151012-192313.

Solve the equation $6x^2 - 6x - 12 = 0$


(S) _____

Solution:

$$6x^2 + -6x + -12 = 0 \quad \text{DOS(4.12a)}$$

$$6(1x^2 - 1x + -2) = 0 \quad \text{DPF(4.3b)}$$

Using the factor organizer,



Solving the linear equations using ZPr (4.11a)

$$x_1 + -2 = 0$$

Case I

$$[x_1 + -2] + 2 = [0] + -2$$

AI(4.5a), SPE(4.15)

$$x_1 + (-2 + 2) = 0 + -2$$

APA(4.2a)

$$x_1 + 0 = -2$$

OOA(2.1)

$$x_1 = -2$$

AId(4.4b)

$$x_1 = -2$$

ONeg(3.1b)

$$x_2 + 1 = 0$$

Case II

$$[x_2 + 1] + -1 = [0] + -1$$

AI(4.5a), SPE(4.15)

$$x_2 + (1 + -1) = 0 + -1$$

APA(4.2a)

$$x_2 + 0 = -2$$

OOA(2.1)

$$x_2 = -2$$

AId(4.4b)

$$x_2 = -2$$

ONeg(3.1b)

D

Dependencies:example 17.1-20151012-190708

12.2 Completing The Square

Completing the square is an algebraic algorithm used to find the solutions of quadratic equations of the form, $ax^2 + bx + c = 0$. Essentially, we want to manipulate this equation such that $x = \text{some value}$. To gain some understanding of how this algorithm works, we will consider each step individually. Let's begin with a quadratic equation in the general form: $ax^2 + bx + c = 0$.

Since we are trying to manipulate the equation $ax^2 + bx + c = 0$ such that $x = \text{some value}$, we first want the coefficient factor a to be equal to 1. This is done by multiplying both expressions by the multiplicative inverse of a (step 1) followed by simplifying both expressions.

$$\begin{aligned} \frac{1}{a}[ax^2 + bx + c] &= \frac{1}{a}[0] && \text{SPE(4.15) + MI(4.10a)} \\ \frac{1}{a} \cdot ax^2 + \frac{1}{a} \cdot bx + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{DPE(4.8a)} \\ \frac{1}{a} \cdot a \cdot x^2 + \frac{1}{a} \cdot b \cdot x + \frac{1}{a} \cdot c &= \frac{1}{a}[0] && \text{JTC(3.5)} \\ x^2 + \frac{b}{a} \cdot x + \frac{c}{a} &= 0 && \text{OOM(2.3)} \\ x^2 + \frac{b}{a}x + \frac{c}{a} &= 0 && \text{CTJ(3.6)} \end{aligned}$$

We now have three terms in the left hand expression where the first two terms have at least one variable factor, x . The problem is that the third term is a constant and we want $x = \text{some value}$. This next step is to get rid of the $\frac{c}{a}$ term, which can be done by using the additive inverse followed by simplifying both expressions.

$$\begin{aligned} \left[x^2 + \frac{b}{a}x + \frac{c}{a} \right] + -\frac{c}{a} &= [0] + -\frac{c}{a} && \text{SPE(4.15) + AI(4.5a)} \\ x^2 + \frac{b}{a}x + \frac{c}{a} + -\frac{c}{a} &= 0 + -\frac{c}{a} && \text{APA(4.2a)} \\ x^2 + \frac{b}{a}x &= -\frac{c}{a} && \text{OOA(2.1)} \end{aligned}$$

The next step is called completing the square. The idea is to add a *NeW* constant, k , to the left-hand expression, $x^2 + \frac{b}{a}x + k$, such that the quadratic expression can then be factored as two identical factors, $(x + m)(x + m) = (x + m)^2$, where $k = m \cdot m$. Notice that since we are adding a constant term, k , to the left-hand expression, then we must also add this constant, k , to the right-hand expression, $x^2 + \frac{b}{a}x + k = -\frac{c}{a} + k$. To determine the values of both m and k we should refer to the organization of the two factors that make up the product of a quadratic expression, $x^2 + \frac{b}{a}x + k = (x + m)^2$.

Since both factors of this new quadratic expression are the same, both terms that make up the middle term, must also be the same. We know that $mx + mx = \frac{b}{a}x$, so we should be able to determine the value of m from this equation. If we can determine the value of m , then we can determine the value of k .

$$\begin{aligned} \frac{b}{a}x &= mx + mx \\ &= 2mx && \text{OOA(2.1)} \end{aligned}$$

Solving for m ,

$$2mx = \frac{b}{a}x$$

$$2 \cdot m \cdot x = \frac{b}{a} \cdot x$$

JTC(3.5)

$$\frac{1}{2} [2 \cdot m \cdot x] = \frac{1}{2} \left[\frac{b}{a} \cdot x \right]$$

SPE(4.15) + MI(4.10a)

$$m \cdot x = \frac{b}{2a} \cdot x$$

OOM(2.3)

$$[m \cdot x] \frac{1}{x} = \left[\frac{b}{2a} \cdot x \right] \frac{1}{x}$$


SPE(4.15) + MI(4.10a)

$$m = \frac{b}{2a}$$

OOM(2.3)

$\left[x^2 + \frac{b}{a}x \right] + \left(\frac{b}{2a} \right)^2 = \left[-\frac{c}{a} \right] + \left(\frac{b}{2a} \right)^2$	SPE(4.15)
$x^2 + \frac{b}{a}x + \left(\frac{b}{2a} \right)^2 = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	APA(4.2b)
$\left(x + \frac{b}{2a} \right) \left(x + \frac{b}{2a} \right) = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	DPF(4.3b)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \left(\frac{b}{2a} \right)^2$	PoTF(3.9)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{(b)^2}{(2a)^2}$	PoQPo(5.5a)
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{b^2}{2^2 a^2}$	PoPrPo
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} + \frac{b^2}{4a^2}$	OOE
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c}{a} \cdot \frac{4a}{4a} + \frac{b^2}{4a^2}$	MId
$\left(x + \frac{b}{2a} \right)^2 = -\frac{c \cdot 4 \cdot a}{a \cdot 4 \cdot a} + \frac{b^2}{4a^2}$	JTC
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4 \cdot a \cdot c}{4 \cdot a \cdot a} + \frac{b^2}{4a^2}$	CPM
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4 \cdot a \cdot c}{4 \cdot a^2} + \frac{b^2}{4a^2}$	PrCBPo
$\left(x + \frac{b}{2a} \right)^2 = -\frac{4ac}{4a^2} + \frac{b^2}{4a^2}$	CTJ
$\left(x + \frac{b}{2a} \right)^2 = \frac{-4ac + b^2}{4a^2}$	CD
$\left(x + \frac{b}{2a} \right)^2 = \frac{b^2 - 4ac}{4a^2}$	CPA
$\left[\left(x + \frac{b}{2a} \right)^2 \right]^{\frac{1}{2}} = \pm \left[\frac{b^2 - 4ac}{4a^2} \right]^{\frac{1}{2}}$	PoI

$x + \frac{b}{2a} = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}}$	PoPrPo
$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{\sqrt{4a^2}}$	PoQPo
$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2^{\frac{1}{2}}a}$	PoPrPo
$x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a}$	OOE
$\left[x + \frac{b}{2a}\right] + \neg\frac{b}{2a} = \left[\pm \frac{\sqrt{b^2 - 4ac}}{2a}\right] + \neg\frac{b}{2a}$	AI
$x + \frac{b}{2a} + \neg\frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a} + \neg\frac{b}{2a}$	APA
$x + \frac{b}{2a} + \neg\frac{b}{2a} = \neg\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$	CPA
$x = \neg\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$	OOA
$x = \frac{\neg b \pm \sqrt{b^2 - 4ac}}{2a}$	CD
$x = \frac{\neg b \pm \sqrt{b^2 - 4ac}}{2a}$	DOS
$x = \frac{\neg b \pm \sqrt{b^2 - 4ac}}{2a}$	ETR
$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$	ONeg



Functions

13. Functions

Property 13.0.1 – Function Value Argument Substitution (FVAS).

$$f(a) \quad (13.1a)$$

Example 13.1 – id:20151012-201647.

Find $f(2)$ given $f(x) = 2x^3 - 3x^2 - 12x + 1$

(S) _____

Solution:

$$f(x) = 2x^3 + -3x^2 + -12x + 1 \quad \text{DOS}(4.12a)$$

$$f(2) = 2[2]^3 + -3[2]^2 + -12[2] + 1 \quad \text{SPE}(4.15)$$

$$f(2) = 2(8) + -3(4) + -12(2) + 1 \quad \text{OOE}(2.5)$$

$$f(2) = 16 + -12 + -24 + 1 \quad \text{OOM}(2.3)$$

$$f(2) = -19 \quad \text{OOA}(2.1)$$

(D) _____

Dependencies:example 17.1-20151012-190708

Example 13.2 – id:20151012-203549.

Find $f(-1)$ given $f(x) = 2x^3 - 3x^2 - 12x + 1$

(S) _____

Solution:

$$f(x) = 2x^3 + -3x^2 + -12x + 1 \quad \text{DOS(4.12a)}$$

$$f(-1) = 2[-1]^3 + -3[-1]^2 + -12[-1] + 1 \quad \text{SPE(4.15)}$$

$$f(-1) = 2(-1) + -3(1) + -12(-1) + 1 \quad \text{OOE(2.5)}$$

$$f(-1) = -2 + -3 + 12 + 1 \quad \text{OOM(2.3)}$$

$$f(-1) = 8 \quad \text{OOA(2.1)}$$

(D) _____

Dependencies:example 17.1-20151012-190708

■

13.1 Inverse Functions

13.2 Inverses

Property 13.2.1 – Cosine Inverse (ArcCos).

$$\cos^{-1}(\cos \theta) = \theta \quad (13.2a)$$

Property 13.2.2 – Sine Inverse (ArcSin).

$$\sin^{-1}(\sin \theta) = \theta \quad (13.3a)$$

Property 13.2.3 – Tangent Inverse (ArcTan).

$$\tan^{-1}(\tan \theta) = \theta \quad (13.4a)$$

Property 13.2.4 – Exponential Inverse (EI).

$$\log_a(a^x) = x \quad (13.5a)$$

Property 13.2.5 – Logarithmic Inverse (LI).

$$a^{\log_a x} = x \quad (13.6a)$$

Property 13.2.6 – Power Inverse (Pol).

$$(b^m)^{\frac{1}{m}} = b \quad (13.7a)$$

Differential Calculus

14	Derivative by First Principles	83
14.1	Limit of the Difference Quotient	
15	Derivative Rules	85
15.1	Derivative of a Monomial Functions	
15.2	Derivative of Polynomial Functions	
15.3	Derivative of a Quotient	
15.4	Derivative of a Rational Function	
16	Equations of Tangent & Secant Lines	95
16.1	Essential Questions	
16.2	Finding the Equation of the Tangent Line	
17	First Derivative Test	99
18	Curve Sketching	101
19	Second Derivative Test	103
20	Curve Sketching	105
	Bibliography	107
	Bibliography	107
	Websites	
	Articles	
	Index	109

14. Derivative by First Principles

14.1 Limit of the Difference Quotient

Definition 14.1.1 – Derivative. The derivative of a function $f(x)$ with respect to the variable x is defined as

$$f'(x) \equiv \lim_{\Delta x \rightarrow 0} \underbrace{\frac{f(x + \Delta x) - f(x)}{\Delta x}}_{\text{Difference Quotient}} \quad (14.1)$$




Figure 14.1: [2]

Example 14.1 – id:20141219-212546.

Differentiate the function $f(x) = 5$

(S)

Solution:

$$f(x) = 5x^0 \quad \text{PoID(4.13a)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5[x + \Delta x]^0 - 5[x]^0}{\Delta x} \quad \text{SPE(4.15)&DBFP(14.1)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{5(1) - 5(1)}{\Delta x} \quad \text{PoID(4.13b)}$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} 0 \quad \text{OOM(2.3)}$$

$$f'(x) = 0$$

■

15. Derivative Rules

15.1 Derivative of a Monomial Functions

Example 15.1 – id:20141124-153017.

Differentiate $f(x) = -3$

(S) _____

Solution:

$$f'(x) = [-3]'$$

SPE(4.15)

$$f'(x) = 0$$

DC(7.1)

(D) _____

Dependencies:

example 15.3-20141124-152503

■

Example 15.2 – id:20141124-141850.

Differentiate $f(x) = x^2$

(S) _____

Solution:

$$f'(x) = [x^2]'$$

SPE(4.15)

$$f'(x) = 2x^{2-1}$$

DPo(7.5)

$$f'(x) = 2x^1$$

OOA(2.1)

$$f'(x) = 2x$$

MId(4.9b)

(S) _____

Less Steps Solution:

$$f'(x) = 2x$$

DPo(7.5)

(D) _____

Dependencies:

example 15.2-20141124-141850

example 15.3-20141124-152503

■

15.2 Derivative of Polynomial Functions

Example 15.3 – id:20141124-152503.Differentiate $f(x) = x^2 - 3$

(S) _____

Solution:

$$\begin{aligned}
 f(x) &= x^2 + -3 && \text{DOS(4.12a)} \\
 f'(x) &= [x^2 + -3]' && \text{SPE(4.15)} \\
 f'(x) &= [x^2]' + [-3]' && \text{DS(7.7)} \\
 f'(x) &= [x^2]' + 0 && \text{DC(7.1)} \\
 f'(x) &= [x^2]' && \text{AId(4.4a)} \\
 f'(x) &= 2x && \text{DPo(7.5) goto 15.2}
 \end{aligned}$$

(S) _____

Less Steps Solution:

$$f(x) = 2x^2 \quad \text{DPo(7.5)&DC(7.5)}$$

(D) _____

Dependencies:

example 15.14-20141124-205219

■

Example 15.4 – id:20151015-155838.Differentiate $y = x^3 + 1$

(S) _____

Solution:

$$\begin{aligned} [y]' &= [x^3 + 1] && \text{SPE(4.15)} \\ y' &= [x^3]' + [1]' && \text{DS(7.7)} \\ y' &= 3x^2 + [1]' && \text{DPo(7.5)} \\ y' &= 3x^2 + 0 && \text{DC(7.1)} \\ y' &= 3x^2 && \text{MId(4.9b)} \end{aligned}$$

(D) _____

Dependencies:example 15.16-20151015-153507

■

Example 15.5 – id:20151011-195002.

Differentiate $f(x) = 2x^2 + 3x + 7$

(S) _____

Solution:

$$\begin{aligned} [f'(x)]' &= [2x^2 + 3x + 7]' && \text{SPE(4.15)} \\ f'(x) &= [2x^2]' + [3x]' + [7]' && \text{DS(7.7)} \\ f'(x) &= 2[x^2]' + 3[x]' + [7]' && \text{DCM(7.3)} \\ f'(x) &= 2[x^2]' + 3[x]' + 0 && \text{DC(7.1)} \\ f'(x) &= 2[x^2]' + 3[x]' && \text{AId(4.4a)} \\ f'(x) &= 2(2x) + 3 && \text{DPo(7.5)} \\ f'(x) &= 4x + 3 && \text{OOM(2.3)} \end{aligned}$$

(D) _____

Dependencies:example 16.1-20151011-154209

■

Example 15.6 – id:20141128-151834.

Differentiate $f(x) = 3x^2 - 6x + 4$

(S) _____

Solution:

$$\begin{aligned}
 f(x) &= 3x^2 + \neg 6x + 4 && \text{DOS(4.12a)} \\
 f'(x) &= [\mathbf{3x^2} + \mathbf{\neg 6x} + \mathbf{4}]' && \text{SPE(4.15)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' + [\mathbf{4}]' && \text{DS(7.7)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' + 0 && \text{DC(7.1)} \\
 f'(x) &= [\mathbf{3x^2}]' + [\mathbf{6x}]' && \text{AId(4.4a)} \\
 f'(x) &= 3[\mathbf{x^2}]' + 6[\mathbf{x}]' && \text{DCM(7.3)} \\
 f'(x) &= 3(2x) + 6(1) && \text{DPo(7.5)} \\
 f'(x) &= 6x + 6 && \text{OOM(2.3)}
 \end{aligned}$$

(S)

$$f'(x) = 6x + 6 \quad \text{DS(7.7)}$$

Example 15.7 – id:20141209-144203.

Differentiate $f(x) = x^2(2x + 4)$

(S)

Solution:

$$\begin{aligned}
 f'(x) &= [\mathbf{x^2}(2x + 4)]' && \text{SPE(4.15)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2[\mathbf{2x} + \mathbf{4}]' && \text{DPr(7.11)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2[\mathbf{2x}] + [\mathbf{4}]' && \text{DS(7.7)} \\
 f'(x) &= [\mathbf{x^2}]'(2x + 4) + x^2 \cdot 2[\mathbf{x}] + [\mathbf{4}]' && \text{DCM(7.3)} \\
 f'(x) &= 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + [\mathbf{4}]' && \text{DPo(7.5)} \\
 f'(x) &= 2x(2x + 4) + x^2 \cdot 2 \cdot 1 + 0 && \text{DC(7.1)} \\
 f'(x) &= 6x^2 + 8x && \text{simplify goto 8.33}
 \end{aligned}$$

Example 15.8 – id:20141209-142321.

Differentiate $f(x) = x^2 \cos(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [x^2 \cos(x)]' && \text{SPE(4.15)} \\ f'(x) &= [x^2]' \cos(x) + x^2[\cos(x)]' && \text{DPr(7.11)} \\ f'(x) &= 2x \cos(x) + x^2[\cos(x)]' && \text{DPo(7.5)} \\ f'(x) &= 2x \cos(x) + x^2(-1 \sin(x)) && \text{DCos(7.19)} \\ f'(x) &= 2x \cos(x) - x^2 \sin x && \text{OOM(2.3)} \end{aligned}$$

■

Example 15.9 – id:20150910-115935.

Differentiate $f(x) = \sin(x) \cos(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [\sin(x) \cos(x)]' && \text{SPE(4.15)} \\ f'(x) &= [\sin(x)]' \cos(x) + \sin(x)[\cos(x)]' && \text{DPr(7.11)} \\ f'(x) &= \cos(x) \cos(x) + \sin(x)[\cos(x)]' && \text{DSin(7.17)} \\ f'(x) &= \cos(x) \cos(x) + \sin(x)(-\sin(x)) && \text{DCos(7.19)} \\ f'(x) &= \cos^2(x) - \sin^2(x) && \text{simplify goto ??} \end{aligned}$$

■

Example 15.10 – id:20141209-151354.

Differentiate $f(x) = \sin(x) \sin(x)$

(S)

Solution:

$$\begin{aligned} f'(x) &= [\sin(x) \sin(x)]' && \text{SPE(4.15)} \\ f'(x) &= [\sin(x)]' \sin(x) + \sin(x)[\sin(x)]' && \text{DPr(7.11)} \\ f'(x) &= \cos(x) \sin(x) + \sin(x) \cos(x) && \text{DSin(7.17)} \\ f'(x) &= \cos(x) \sin(x) + \cos(x) \sin(x) && \text{CPM(4.6)} \\ f'(x) &= 2 \cos(x) \sin(x) && \text{OOA(2.1)} \end{aligned}$$

■

Example 15.11 – id:20141124-203850.

Differentiate $y = \ln(3x)$

(S)

Solution:

After identifying that $y = \ln(3x)$ is a composite function, we let $u = 3x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = 3x$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = 3$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.16)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot \frac{du}{dx} \quad \text{DNL(7.31)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot 3 \quad \text{DPo(7.5)}$$

$$\frac{dy}{dx} = \frac{1}{3x} \cdot 3$$

$$\frac{dy}{dx} = \frac{3}{3x} \quad \text{OOM(2.3)}$$

$$\frac{dy}{dx} = \frac{1}{x}$$

(D)

Dependencies:

example 15.14-20141124-205219

Example 15.12 – id:20141128-160248.

Differentiate $y = \ln(3x^2 - 6x + 4)$

(S)

Solution: After identifying that $y = \ln(3x^2 - 6x + 4)$ is a composite function, we let $u = 3x^2 - 6x + 4$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = 3x^2 - 6x + 4$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = 6x - 6 \quad \text{goto 15.6}$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.16)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot \frac{du}{dx} \quad \text{DNL(7.31)}$$

$$\frac{dy}{dx} = \frac{1}{u} \cdot (6x - 6) \quad \text{DPo(7.5)}$$

$$\frac{dy}{dx} = \frac{1}{3x^2 - 6x + 4} \cdot (6x - 6)$$

$$\frac{dy}{dx} = \frac{6x - 6}{3x^2 - 6x + 4} \quad \text{OOM(2.3)}$$

Example 15.13 – id:20141128-155506.

Differentiate $y = \ln(\cos x)$

(S)

Solution: After identifying that $y = \ln(\cos x)$ is a composite function, we let $u = \cos x$ and thus we get a new function $y = \ln(u)$.

Using chain rule,

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \quad \text{DComp(7.16)}$$

We need to find the factors $\frac{dy}{du}$ and $\frac{du}{dx}$.

$$y = \ln(u) \quad u = \cos x$$

$$\frac{dy}{du} = \frac{1}{u} \quad \frac{du}{dx} = -\sin x$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\text{d}y}{\text{d}u} \cdot \frac{\text{d}u}{\text{d}x} && \text{DComp(7.16)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot \frac{\text{d}u}{\text{d}x} && \text{DNL(7.31)} \\ \frac{dy}{dx} &= \frac{1}{u} \cdot -\sin x && \text{DPo(7.5)} \\ \frac{dy}{dx} &= \frac{1}{\cos x} \cdot -\sin x \\ \frac{dy}{dx} &= \frac{-\sin x}{\cos x} && \text{OOM(2.3)} \\ \frac{dy}{dx} &= -\tan x\end{aligned}$$

■

Example 15.14 – id:20141124-205219.Differentiate $y = (x^2 - 1) \ln(3x)$

(S)

Solution:

$$\begin{aligned}y' &= [x^2 - 3]' \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' && \text{DPr(7.11)} \\ y' &= 2x \cdot \ln x + (x^2 - 3) \cdot [\ln 3x]' && \text{differentiate goto 15.3} \\ y' &= 2x \cdot \ln x + (x^2 - 3) \cdot \frac{1}{x} && \text{differentiate goto 15.11} \\ y' &= 2x \cdot \ln x + \frac{x^2 - 3}{x} && \text{OOM(2.3)} \\ y' &= 2x^2 \ln x + \frac{x^2 - 3}{x} && \text{JTC(3.5)} \\ y' &= \frac{2x^2 \ln x + (x^2 - 3)}{x} && \text{OOA(2.1)}\end{aligned}$$

■

15.3 Derivative of a Quotient**Example 15.15 – id:20151015-165037.**Differentiate $y = \frac{\sin x}{e^x}$

(S)

Solution:

$$\frac{d[y]}{dx} = \frac{d\left[\frac{\sin x}{e^x}\right]}{dx} \quad \text{SPE(4.15)}$$

$$\frac{dy}{dx} = \frac{\frac{d}{dx}[\sin x]e^x - \sin x \frac{d}{dx}[e^x]}{[e^x]^2} \quad \text{DQ(??)}$$

$$\frac{dy}{dx} = \frac{\frac{d}{dx}[\sin x]e^x - \sin x \frac{d}{dx}[e^x]}{e^{2x}} \quad \text{PoPo(5.1a)}$$

$$\frac{dy}{dx} = \frac{\cos x e^x - \sin x \frac{d}{dx}[e^x]}{e^{2x}} \quad \text{DSin(7.18)}$$

$$\frac{dy}{dx} = \frac{\cos x e^x - \sin x e^x}{e^{2x}} \quad \text{DExp(7.34)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^{2x}} \quad \text{DPF(4.3b)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^x e^x} \quad \text{PrCBPo(5.3b)}$$

$$\frac{dy}{dx} = \frac{e^x(\cos x - \sin x)}{e^x e^x} \quad \text{MId(4.9a)}$$

$$\frac{dy}{dx} = \frac{\cos x - \sin x}{e^x} \quad \blacksquare$$

15.4 Derivative of a Rational Function

Example 15.16 – id:20151015-153507.Differentiate $f(x) = \frac{x^3 + 1}{x^2}$

(S)

Solution:

Method 1: Derivative of a quotient

$$[f(x)]' = \left[\frac{x^3 + 1}{x^2} \right]' \quad \text{SPE(4.15)}$$

$$f'(x) = \frac{[x^3 + 1]'(x^2) + -1((x^3 + 1)[x^2]')}{[x^2]^2} \quad \text{DQ(??)}$$

$$f'(x) = \frac{(3x^2)(x^2) - 1((x^3 + 1)[x^2]')}{[x^2]^2} \quad \text{differentiate goto 15.4}$$

$$f'(x) = \frac{(3x^2)(x^2) - 1((x^3 + 1)(2x))}{[x^2]^2} \quad \text{differentiate goto 15.2 } f'(x) = \frac{3x^3 - 2x^2 - 2}{x^3} \quad \text{Simplif}$$

16. Equations of Tangent & Secant Lines

16.1 Essential Questions

Essential Questions 16.1

1. How do we find the equation of the tangent line of a given function at the point $P(a, b)$?
2. How do we find the equation of the tangent line of a given function at $x = a$?

16.2 Finding the Equation of the Tangent Line




Figure 16.1: Finding the Equation of a Tangent Line Workflow

For a given function, $f(x)$, our goal is to find the equation of the tangent line at the point $P(a, b)$, which can be expressed in slope-intercept form as:

$$y = f'(a)x + k$$

Where $f'(a)$ is the value of the derivative, slope, at $x = a$ and k is the y -intercept. We therefore need to:

1. Find the derivative of the function $f(x)$: $f'(x)$
2. Find the value of the derivative at $x = a$: $f'(a)$
3. Find the value of the y -intercept: $k = -f'(a) + b$:
4. If the ordinate b is not explicitly given, then find $f(a) = b$

Example 16.1 – id:20151011-154209.

Find the equation of the line tangent to the curve of the function $f(x) = 2x^2 + 3x + 7$ at the point $P(2, 21)$.

(S)

Solution:

Find the derivative of $f(x)$

$$f'(x) = 4x + 3 \text{ goto 15.5}$$

Evaluate the derivative at $x = 2$

$$f'(2) = 4[2] + 3$$

SPE(4.15)

$$f'(2) = 8 + 3$$

OOM(2.3)

$$f'(2) = 11$$

OOA(2.1)

Find the y -intercept, k , of the equation of the tangent line.

$$y = f'(x)x + k$$

$$[21] = [11][2] + k \quad \text{SPE(4.15)}$$

$$21 = 22 + k$$

OOM(2.3)

$$\neg 22 + [21] = \neg 22 + [22 + k]$$

SPE+AI

$$\neg 22 + 21 = (\neg 22 + 22) + k$$

APA(4.2a)

$$\neg 1 = 0 + k$$

OOA(2.1)

$$\neg 1 = k$$

AId(4.4a)

$$\neg 1 = k$$

ONeg(3.1b)

$$k = -1$$

SyPE(4.16a)

The equation of the tangent line is

$$y = f'(2)x + k$$

$$y = 11x - 1 \quad \text{SPE(4.15)}$$

Example 16.2 – id:20151015-171630.

Given $f(x) = e^{3x}$, find the the equation of the tangent line L to the curve f at the point $(2, e^2)$.

(S)

Solution:

Find the derivative of $f(x)$

$[f(x)]' = [e^{3x}]'$	SPE(4.15)
$f'(x) = [3x]'[e^{3x}]'$	DCF(7.15)
$f'(x) = 3[x]'[e^{3x}]'$	DCM(7.3)
$f'(x) = 3 \cdot 1[e^{3x}]'$	DPo(7.5)
$f'(x) = 3[e^{3x}]'$	OOM(2.3)
$f'(x) = 3e^{3x}$	DExp(7.33)

Evaluate the derivative at $x = 2$

$$\begin{aligned} f'(2) &= 3e^{3[2]} && \text{SPE(4.15)} \\ f'(2) &= 3e^6 && \text{OOM(2.3)} \end{aligned}$$

Find the y -intercept, k , of the equation of the tangent line.

$$\begin{aligned} y &= f'(x)x + k \\ [e^2] &= [3e^6][2] + k && \text{SPE(4.15)} \\ e^2 &= 3 \cdot e^6 \cdot 2 + k && \text{JTC(3.5)} \\ e^2 &= 3 \cdot 2 \cdot e^6 + k && \text{CPM(4.6)} \\ e^2 &= 6 \cdot e^6 + k && \text{OOM(2.3)} \\ e^2 &= 6e^6 + k && \text{CTJ(3.6)} \\ -6e^6 + [e^2] &= -6e^6 + [6e^6 + k] && \text{AI(4.5a), SPE(4.15)} \\ e^2(-6e^4 + 1) &= -6e^6 + [6e^6 + k] && \text{DPF(4.3b)} \\ -6e^6 + e^2 &= (-6e^6 + 6e^6) + k && \text{APA(4.2a)} \\ e^2(-6e^4 + 1) &= 0 + k && \text{OOA(2.1)} \\ e^2(-6e^4 + 1) &= k && \text{AId(4.4b)} \\ e^2(-6e^4 + 1) &= k && \text{ONeg(3.1b)} \end{aligned}$$

The equation of the tangent line, L is

$$\begin{aligned} y &= f'(2)x + k \\ y &= 3e^6x + e^2(-6e^4 + 1) && \text{SPE(4.15)} \end{aligned}$$

17. First Derivative Test

Example 17.1 – id:20151012-190708.

Given the function $f(x) = 2x^3 - 3x^2 - 12x + 1$ find the critical points, classify the critical points and find the intervals of increasing/decreasing.

(S)

Solution:

Differentiate the function:

$$f(x) = 2x^3 + -3x^2 + -12x + 1 \quad \text{DOS(4.12a)}$$

$$[f(x)]' = [2x^2 + -3x^2 + -12x + 1]' \quad \text{SPE(4.15)}$$

$$f'(x) = [2x^3]' + [-3x^2]' + [-12x]' + [1]' \quad \text{DS(7.7)}$$

$$f'(x) = 2[x^3]' + -3[x^2]' + -12[x]' + [1]' \quad \text{DCM(7.3)}$$

$$f'(x) = 2(3x^2) + -3(2x) + -12(1) + [1]' \quad \text{DPo(7.5)}$$

$$f'(x) = 6x^2 + -6x + -12 + [1]' \quad \text{OOM(2.3)}$$

$$f'(x) = 6x^2 + -6x + -12 + 0 \quad \text{DC(7.1)}$$

$$f'(x) = 6x^2 + -6x + -12 \quad \text{AId(4.4a)}$$

Solving the equation $6x^2 + -6x + -12 = 0$ to find the x value(s) of the critical points, goto 12.2, we find that $x = 2$ and $x = -1$ are critical values of the function.

Since we are looking for critical **points**, we need to find the ordinates, y -values, of the critical points by evaluating the function for the given critical x -values.

$$f(2) = -19 \text{ goto 13.1}$$

$$f(-1) = 8 \text{ goto 13.2}$$


Therefore the critical points are $(2, -19)$ and $(-1, 8)$.

The first derivative test can be used to determine the intervals of increasing/decreasing and consequently we will be able to classify the critical points. Since we are only interested in the values of the derivative, it will be easier to use the factored form of the derivative, $f'(x)=6(x-2)(x+1)$

1st Derivative Test Table

x	$x < -1$	$x = -1$	$-1 < x < 2$	$x = 2$	$x > 2$
$f'(x)$	+	0	-	0	+
$f(x)$	↗	→	↘	→	↗
C.P.		Max		Min	

From the first derivative table we can classify the critical point $(2, -19)$ as a local maximum and $(-1, 8)$ as a local minimum.



18. Curve Sketching

18.0.1 Finding the vertex of a quadratic function using differentiation.

We can find the vertex of a quadratic function, $f(x)$ using differentiation by:

1. Differentiate the function: Find $f'(x)$.
2. Set the derivative equal to zero: $f'(x) = 0$.
3. Find the abscissa of the vertex by solving the equation $f'(x) = 0$ for x to find the critical x value: $x = k$.
4. Find the ordinate of the vertex by substituting the value of critical value $x = k$ into the function $f(x)$: Evaluate $f(k)$

Example 18.1 – id:20151008-110208.

Find the vertex of the quadratic function, $f(x) = ax^2 + bx + c$, using differentiation.

(S)

Solution:

1. Find the derivative of $f(x)$

$$[f(x)]' = [ax^2 + bx + c]' \quad \text{SPE(4.15)}$$

$$f'(x) = [ax^2]' + [bx]' + [c]' \quad \text{DS(7.7)}$$

$$f'(x) = a[x^2]' + b[x]' + [c]' \quad \text{DCM(7.3)}$$

$$f'(x) = 2 \cdot a \cdot x + b \cdot 1 + [c]' \quad \text{DPo(7.5)}$$

$$f'(x) = 2 \cdot a \cdot x + 1 \cdot b + [c]' \quad \text{CPM(4.6)}$$

$$f'(x) = 2ax + 1b + [c]' \quad \text{CTJ(3.6)}$$

$$f'(x) = 2ax + b + [c]' \quad \text{MId(4.9b)}$$

$$f'(x) = 2ax + b + 0 \quad \text{DC(7.1)}$$

$$f'(x) = 2ax + b \quad \text{AId(4.4b)}$$

2. Set the derivative equal to zero and solve for x .

$$f'(x) = 0$$

$$2ax + b = 0$$

$$x = -\frac{b}{2a} \text{ goto 11.9}$$

The abscissa of the vertex is $x = -\frac{b}{2a}$.

3. Find the ordinate of the vertex by substituting the argument $x = -\frac{b}{2a}$ into $f(x)$

Example 18.2 – id:20150923-152515.

Find the vertex of the parabola $y = x^2 - 2x - 6$ using differentiation.

(S)

- Differentiate the function.

$$f(x) = x^2 - 2x - 6$$

$$f(x) = x^2 + \neg 2x + \neg 6$$

$$[f(x)]' = [x^2 + \neg 2x + \neg 6]'$$

$$f'(x) = [x^2]' + [\neg 2x]' + [\neg 6]'$$

$$f'(x) = [x^2]' + \neg 2[x]' + [\neg 6]'$$

$$f'(x) = 2x + \neg 2 + [\neg 6]'$$

$$f'(x) = 2x + \neg 2 + 0$$

$$f'(x) = 2x + \neg 2$$

$$f'(x) = 2x - 2$$

DOS(4.12a)

SPE(4.15)

DS(7.7)

DCM(7.3)

DPo(7.5)

DC(7.1)

AId(4.4b)

DOS(4.12b)

- Set the derivative equal to zero and solve for x

$$2x - 2 = 0$$

$$x = 1$$

- Find the value of $f(1)$

$$f(x) = x^2 - 2x - 6$$

$$f(1) = [1]^2 - 2[1] - 6$$

$$f(1) = -7$$

SPE(4.15)

Evaluate

The vertex of this parabola is the point $(1, -7)$

■

19. Second Derivative Test

20. Curve Sketching




Figure 20.1: Overview of Curve Sketching

Bibliography

Books

Website

- [] *Indeterminate (variable)*. [http://en.wikipedia.org/wiki/Indeterminate_\(variable\)](http://en.wikipedia.org/wiki/Indeterminate_(variable)) (cited on page 35).
- [] *Polynomial*. <http://en.wikipedia.org/wiki/Polynomial> (cited on pages 35, 60).

Articles

Index

A

Additive Identity	15
Additive Inverse	16
Algebraic Limit Theorem	25
AlgebraicLimitTheorem	
Limit of a Constant	25
Limit of a Difference	25
Limit of a Product	25
Limit of a Quotient	25
Limit of a Sum	25

C

Coefficient.....	33
Common Denominator	9
Cosine Inverse	78

D

Definition of Subtraction.....	17
Degree of the Indeterminate	33
Degree of the Univariate Polynomial	58
Derivative	81
DerivativeRules	
Derivative of a Composite Function	28
Derivative of a Constant	27
Derivative of a Constant Multiple	27
Derivative of a Difference	27
Derivative of a Logarithm	29
Derivative of a Natural Exponential	30
Derivative of a Natural Logarithm	29
Derivative of a Power	27
Derivative of a Product	28
Derivative of a Quotient	28
Derivative of a Sum	27
Derivative of Cosecant	29
Derivative of Cosine	28
Derivative of Cotangent	29

Derivative of Secant	29
Derivative of Sine	28
Derivative of Tangent	28
Derivative Structural Rules	27
Logarithm Derivative Structural Rules	
29	
Monomial Derivative Rules	27
Trigonometric Derivative Structural Rules	
28	

Derivatives

Derivative of an Exponential	29
Dyadic Operations	9

E

Example	
20141027-075159	36
20141105-161225	56
20141106-150622	38
20141106-151138	42
20141106-152020	39
20141106-152339	51
20141106-154547	42
20141107-121834	51
20141107-131748	69
20141108-173613	41
20141108-191616	43
20141108-194156	40
20141108-194431	40
20141108-194709	43
20141109-090809	44
20141109-091015	45
20141109-092448	45
20141109-092651	46
20141109-092910	46
20141109-093220	47
20141109-093419	48
20141109-094928	48
20141109-095151	49
20141109-095536	49
20141109-095842	50

20141109-133008	53
20141109-133316	54
20141109-140659	54
20141109-141019	55
20141109-141347	56
20141111-215726	66
20141111-222931	63
20141120-202042	35
20141120-202842	41
20141120-203846	35
20141121-093439	35
20141121-093747	34
20141121-184652	38
20141121-185558	36
20141121-190857	37
20141121-193636	37
20141124-141850	83
20141124-152503	84
20141124-153017	83
20141124-203850	87
20141124-205219	90
20141128-151834	85
20141128-155506	89
20141128-160248	88
20141206-101107	64
20141206-101632	64
20141206-102142	63
20141206-102404	65
20141206-102723	66
20141206-104404	65
20141209-142321	86
20141209-144203	86
20141209-145211	52
20141209-151354	87
20141219-212546	81
20150910-115935	87
20150923-152515	100
20151008-110208	99
20151011-154209	94
20151011-195002	85
20151012-190708	97
20151012-192313	69
20151012-201647	77
20151012-203549	77
20151015-104754	67
20151015-153507	91
20151015-155838	84
20151015-165037	90
20151015-171630	94
20151015-184212	61
20151016-063338	61
Exponential Inverse	78
F	
Fraction Operation of Addition	10
Function Value Argument Substitution	77
Functions	
Inverses	78
G	
Greatest Common Divisor	7
I	
Identities	
Power Identities	19, 20
Improper Fraction	7
Indeterminate	33
L	
Like Terms	34
Logarithmic Inverse	78
LogarithmIdentity	
Logarithm Power of a Power	20
Logarithm Product of Common Base Powers	20
Logarithm Quotient of Common Base Powers	20
Logartihms	
Logarithm Exponent Visible	12
Logarithm to Power	12
M	
Multiplicative Identity	16
Multiplicative Inverse	16
N	
Notation	
Center-Dot to Juxtaposition	12

Derivative Notation	12
Euler's first derivative	13
Euler's nth derivative	14
Euler's second derivative	13
Juxtaposition to Center-Dot	12
LaGrange's Evaluate derivative	13
LaGrange's first derivative	13
LaGrange's nth derivative	13
LaGrange's second derivative	13
Leibniz's Evaluate derivative	13
Leibniz's first derivative	13
Leibniz's nth derivative	13
Leibniz's second derivative	13
Logarithm Notation	12
Multiplication Center-Dot	11
Multiplication Juxtaposition	11
Multiplication Notation	11
Multiplication Times	11
Negation Notation	11
Power Notation	12
Number System	
Integers	7
Natural Numbers	7
Rational Numbers	7
Number Systems	7
 O	
Operation	
Operation of Addition	9
Operation of Exponentiation	9
Operation of Multiplication	9
Operation of Negation	11
 P	
Positive Integers	7
Power Identity	17
Power Inverse	78
Powers	
Factor to Power	12
Power Negative Exponent	12
Power of a Power	19
Power of a Product of Powers	19
Power of a Quotient of Powers	19
Power to Factor	12
Power To Logarithm	12
Product Common Base Powers	19
Quotient Common Base Powers	19
Radical to Power	12
Prime Numbers	7
Proper Fraction	7
Properties	
Summary of Field Properties	15
Properties of Addition	15
Property	
Associative Property of Addition	15
Associative Property of Multiplication	16
Commutative Property of Addition	15
Commutative Property of Multiplication	16
Distributive Property Factoring	15, 16
Inverses	78
Properties of Equality	17
Properties of Inequality	18
Properties of Multiplication	16
Properties of Powers	17
Properties of Subtraction	17
Zero Product	17
 R	
Reflexive Property of Equality	17
Relatively Prime	8
 S	
Simplifying Degree 0 Univariate Monomials	33
Sine Inverse	78
Solving Quadratic Equations	
Completing the Square	70
Power Inverse	63, 69
Substitution Property of Equality	17
Substitution Property of Inequality	18
Symmetric Property of Equality	17
 T	
Tangent Inverse	78
Transitive Property of Equality	17
Transitive Property of Inequality	18
Trigonometric Identity	
Cosine Difference of Angles Identity	21

- Cosine Double Angle Identity 21
Cosine Sum of Angles Identity 21
Sine Difference of Angles Identity . 21
Sine Double Angle Identity 21
Sine Sum of Angles Identity 21
Tangent Difference of Angles Identity
 22
Tangent Double Angle Identity 21
Tangent Sum of Angles Identity ... 21
Trigonometric Cotangent Identity . 21
Trigonometric Pythagorean Identities
 20
Trigonometric Reciprocal Identities 20
Trigonometric Tangent Identity 20

U

- Univariate Monomial 33
Univariate Polynomial Expression 57

Z

- Zero Factor Property 18