

Quantitative Methods 2

Tutorial 4

Nhan La

Last week

1. Descriptive statistics

2. Normality test

- Visual assessment
- Descriptive measures: **relative comparisons**
 - Mean vs. median
 - Skewness
 - \widehat{SK} vs. 0
 - $|\widehat{SK}|$ vs. $2s_{\widehat{SK}}$
 - Kurtosis
 - \widehat{K} vs. 3
 - $|\widehat{K} - 3|$ vs. $2s_{\widehat{K}}$
- Hypothesis test: Jacque-Bera

Last week

3. Test of population's central location

- Parametric: t test
- Non-parametric
 - Sign test
 - Wilcoxon signed ranks test
 - Sample size: $n = \text{number of non-zero deviations}$
- Find confidence interval of the population mean

Last week

- Some tests need d.f: t , χ^2 , F
 - Probability distribution, and thus the estimation, of the test statistics depend on the sample size or the number of groups, or both
 - E.g., $t = \frac{\bar{X} - \mu_0}{s/\sqrt{n}}$
- Some don't: Z , Fisher's exact test

Hypothesis testing

- Choose the appropriate test based on:
 - Levels of measurement
 - Types of variable
 - Distribution of the underlying population
 - Sample size
 - Relevant population parameter (e.g., σ^2) known/unknown

Tutorial 4


1. Paired-sample test

- Parametric: Z test or t test
- Nonparametric: sign test or Wilcoxon signed rank test


2. Independent samples test

- Parametric: Z test or t test
- Nonparametric: Wilcoxon rank sum test


One population



Two populations - Numerical



Two populations - Ordinal and Nominal



Tutorial 4


Wilcoxon signed rank test: Rank of each observation given by its deviation from the population median.

Values	7	3	4	6	10	5	6	4	3	8
Deviation	1	-3	-2	0	4	-1	0	-2	-3	2
Abs. dev.	1	3	2	0	4	1	0	2	3	2
Ranks	1.5	6.5	4		8	1.5		4	6.5	4

Calculate the sums of negative and positive deviation groups

Tutorial 4

Wilcoxon rank sum test: Rank of each observation given by its position in the pooled sample



Tutorial 4

Parameter	Population	Test	EViews default reported p-values
Normality		Jacque-Bera test	One-sided
Central location	One population	Mean: Z- or t-test	Two-sided
		Median: Sign test or Wilcoxon signed rank test (using normal approx. Z-test)	Two-sided
	Two populations – Paired samples	Mean: Z- or t-test	Two-sided
		Median: Sign test or Wilcoxon signed rank test (using normal approx. Z-test)	Two-sided
	Two populations – Independent samples	Mean: Z- or t-test	Two-sided
		Median: Wilcoxon rank-sum test (using normal approx. Z-test)	Two-sided
	Three or more populations – Independent samples	Mean: F-test	One-sided
		Median: Kruskal-Wallis test	One-sided

Tutorial 4

Why always 0?

Depend on the question being asked.

Two sided test:

$$H_0: \mu_0 = \mu_1 \text{ or } H_0: \mu_0 - \mu_1 = 0;$$

$$H_A: \mu_0 \neq \mu_1 \text{ or } H_A: \mu_0 - \mu_1 \neq 0$$

Tutorial 4

Parameter	Population	Test	EViews default reported p-values
Variance	One population	t-test	One-sided, smaller value (see Week 4 lecture slides) Note: it's more straightforward to rely on the comparison between σ^2 and s^2
	Two populations	F-test	Two-sided (see Week 4 lecture slides, especially for one-sided test hypotheses and statistic formation)
Proportion	One population	Z-test	Two-sided
	Two populations	t-test (to produce approximated results)	Two-sided

Tutorial 4

Parameter	Population	Test	EViews default reported p-values
Correlation and regression	Pearson correlation coefficient	t-test	Two-sided
	Coefficient (slopes or intercept)	t-test	Two-sided
	Model significance/utility	F-test	One-sided
	Linear restriction of coefficient	One conjecture ($<$, $>$ or $=$): t-test or F-test Note: the conjecture can involve more than one parameter	t-test: Two-sided F-test: One-sided
		More than one conjecture: F-test	One-sided
	Heteroskedasticity White test	F-test	One-sided