UNIVERSITY OF CAMBRIDGE

Analysis of Stochastic Gradient Descent in Continuous Time

Jonas Latz

Department of Applied Mathematics and Theoretical Physics, University of Cambridge, UK

SIAM Conference on Mathematics of Data Science 2020

Analysis of Stochastic Gradient Descent in Continuous Time

Preprint: L. 2020: —, arXiv:2004.07177



2004.07177-2.pdf (page 1 of 29)

Analysis of Stochastic Gradient Descent in Continuous Time*

Jonas Latz

Department of Applied Mathematics and Theoretical Physics, University of Cambridge
Wilberforce Road, Cambridge, CB3 0WA, United Kingdom
j12160@cam.ac.uk

Jun 2020

Abstract

Stochastic gradient descent is an optimisation method that combines classical gradient descent with random subsampling within the target functional. In this work, we introduce the stochastic gradient process as a continuous-time representation of stochastic gradient descent.

Funding: Engineering and Physical Sciences Research Council (EPSRC), Swindon, UK

Outline

Motivation

Modelling Stochastic Gradient Descent in Continuous Time

Longtime behaviour

Practical considerations and illustrations

Conclusions

Outline

Motivation

- Stochastic Gradient Descent
- Continuous time models

Modelling Stochastic Gradient Descent in Continuous Time

Longtime behaviour

Practical considerations and illustrations

Conclusions

Optimisation problem

- Consider an optimisation problem on $X := \mathbb{R}^K$; of the form

$$\theta^* \in \operatorname{argmin}_{\theta \in X} \bar{\Phi}(\theta) := \frac{1}{N} \sum_{i=1}^{N} \Phi_i, \qquad \text{(OptProb)}$$

  with potentials $\bar{\Phi}, \Phi_i \in C^1(X; \mathbb{R}), i \in I := \{1, ..., N\}$.

- Typical in statistical, imaging, and machine learning applications:
  - $\bar{\Phi}$: misfit between a model and a (big) data set
  - Φ_i: misfit between a model and the i-th partition of the data set

Stochastic Gradient Descent

- Algorithm to solve (OptProb): Stochastic Gradient Descent (SGD) [Robbins & Monro: 1951]

Algorithm 1 Stochastic gradient descent (SGD)

1: initialise $\theta_0 \in X$ deterministically or randomly
2: define non-increasing sequence $(\eta_k)_{k=1}^{\infty} \in (0, \infty)^{\mathbb{N}}$
3: **for** $k = 1, 2, \ldots$ **do**
4: sample $\boldsymbol{i}_k \sim \mathrm{Unif}(I)$
5: $\theta_k \leftarrow \theta_{k-1} - \eta_k \nabla \Phi_{\boldsymbol{i}_k}(\theta_{k-1})$
6: **return** $(\theta_k)_{k=0}^{\infty}$

- gradient descent steps w.r.t. randomly selected Φ_i instead of $\bar{\Phi}$ $\rightarrow$ memory efficient
- $(\eta_k)_{k=1}^{\infty}$: step lengths, learning rates

Stochastic Gradient Descent

- SGD constructs a Markov chain
- Stochastic properties hardly discussed [Benaïm; 1999][Dieuleveut et al.; 2017][Hu et al.; 2019]
  - Stationary distribution?
  - Ergodicity?
  - Speed of convergence?
    $\rightarrow$ this talk
- Long-term goals
  - Construct more efficient stochastic optimisation algorithms
  - Understand random subsampling in SGD and other methods
  - Understand SGD in non-convex optimisation

Stochastic Gradient Descent in continuous time?

- Idealisation and simplification of models through continuity assumption
  - Usual modelling tool in many scientific disciplines (e.g., continuum mechanics,...)
  - Recently also used in data science, machine learning, and algorithms
    - Ensemble Kalman Inversion [Schillings & Stuart; 2017, 2018][Blömker et al.; 2019]...
    - Continuum limits of graphs [Trillos & Sanz-Alonso; 2018] and in MCMC [Kuntz et al.; 2019]
    - PDE-based image reconstruction [Rudin et al.; 1992][Schönlieb; 2015]...
  - continuous models tend to be easier to analyse: no numerical artefacts

- Past work on SGD in continuous time:
  - Idea: $\eta_k \approx 0 \Rightarrow \Phi_{\boldsymbol{i}_k} - \bar{\Phi}$ approximately Gaussian (CLT)
  - Hence, $(\theta_k)_{k=1}^{\infty}$ can be represented by a diffusion process

$$\dot{\theta}(t) = -\nabla\bar{\Phi}(\theta(t)) + \Sigma(\theta(t))\dot{\mathrm{W}}_t \quad (t \geq 0), \qquad \theta(0) = \theta_0.$$

[Hu et al.; 2019][Li et al.; 2016, 2017, 2019][Mandt et al.; 2015, 2016, 2017]

Stochastic Gradient Descent in continuous time?

- Diffusion approximation may be not very accurate
  - does not represent the discrete nature of the potential selection
  - for large η_k, the paths of $(\theta_k)_{k=1}^{\infty}$ are very different from a diffusion

- Purpose of this work:
  - Give an improved continuous-time model of SGD and show its suitability
  - Analyse the model
  - Find implications on (discrete-time) SGD

Outline

Motivation

Modelling Stochastic Gradient Descent in Continuous Time

- Idea and definition
- Comparison to SGD
- Biological interpretation

Longtime behaviour

Practical considerations and illustrations

Conclusions

Observations and fundamental idea

- $\theta_k \leftarrow \theta_{k-1} - \eta_k \nabla \Phi_{\boldsymbol{i}_k}(\theta_{k-1})$ is a forward Euler discretisation of the gradient flow

$$\dot{\theta}(t) = -\nabla \Phi_{\boldsymbol{i}_k}(\theta(t))$$

- learning rate η_k has two different meanings
  (i) η_k is the step size of the gradient flow discretisation
  (ii) η_k determines the length of the time interval with which we switch the Φ_i

Idea.

Obtain a continuous time model for SGD, by

(i) let the step size go to 0

(ii) switch the potentials in the gradient flow at a rate of $1/\eta_k$

Switching of potentials

- control the switching of the potentials by a continuous-time Markov process (CTMP) $(\boldsymbol{i}(t))_{t\geq 0}$ on $I := \{1, ..., N\}$

CTMPs 101

- $(\boldsymbol{i}(t))_{t\geq 0}$ is piecewise constant
- randomly jumps from one state to another after a random waiting time $T \sim \pi_{\mathrm{wt}}(\cdot|t_0)$, where $\pi_{\mathrm{wt}}([t,\infty)|t_0) = \exp\left(-\int_0^t \nu(u+t_0)\mathrm{d}u\right), t \geq 0.$

- two versions: $\begin{cases} \text{(i) constant learning rate } \eta_\bullet \equiv \eta > 0 \\ \text{(ii) decreasing learning rate } \eta_\bullet > 0, \text{ with } \eta_k \downarrow 0 \ (k \to \infty) \end{cases}$

Continuous-time Markov processes

Definition.

(i) Let $(\boldsymbol{i}(t))_{t\geq 0}$ be a CTMP on I with $\boldsymbol{i}(0) \sim \mathrm{Unif}(I)$ and transition rate matrix

$$A \in \mathbb{R}^{N\times N} : A_{i,j} := \begin{cases} \frac{1}{(N-1)\eta}, & \text{if } i \neq j, \\ -\frac{1}{\eta}, & \text{if } i = j. \end{cases}$$

(ii) Let $(\boldsymbol{j}(t))_{t\geq 0}$ be a CTMP on I with $\boldsymbol{j}(0) \sim \mathrm{Unif}(I)$ and transition rate matrix

$$B \in \mathbb{R}^{N\times N\times [0,\infty)} : B(t)_{i,j} := \begin{cases} \frac{1}{(N-1)H(t)}, & \text{if } i \neq j, \\ -\frac{1}{H(t)}, & \text{if } i = j, \end{cases}$$

where $(H(t))_{t\geq 0}$ is Lipschitz & interpolates $(\eta_k)_{k=1}^{\infty}$.

Stochastic gradient process

the Stochastic gradient process (SGP) is our continuous-time version of SGD

Definition

We define the Stochastic gradient process...

(i) ...with constant learning rate (SGPC) by $(\theta(t))_{t\geq 0}$, which satisfies

$$\dot{\theta}(t) = -\nabla\Phi_{i(t)}(\theta(t)) \quad (t \geq 0), \qquad \theta(0) = \theta_0.$$

(ii) ...with decreasing learning rate (SGPD) by $(\xi(t))_{t\geq 0}$, which satisfies

$$\dot{\xi}(t) = -\nabla\Phi_{j(t)}(\xi(t)) \quad (t \geq 0), \qquad \xi(0) = \xi_0.$$

$(\theta(t))_{t\geq 0}$ and $(\xi(t))_{t\geq 0}$ are almost surely well-defined, if

Assumption 2.2. *For $i \in I : \Phi_i \in C^1(X, \mathbb{R})$ and $\nabla\Phi_i$ is Lipschitz continuous.*

Stochastic gradient process



Figure: Cartoon of SGPC

Piecewise deterministic Markov processes

$(\theta(t))_{t\geq 0}$, $(\xi(t))_{t\geq 0}$ are piecewise deterministic Markov processes (PDMPs)

- 'a general class of non-diffusion stochastic models' [Davis; 1984, 1993]
- progression via deterministic dynamic (ODE) with jumps after random waiting times or when hitting a boundary [Bakhtin & Hurth; 2012][Benaïm et al.; 2012, 2015][Yin & Zhu; 2010]...
- used for stochastic modelling in engineering, computer science, and biology [Rudnicki & Tyran-Kamińska; 2017]
- used as a basis for non-reversible MCMC algorithms [Bierkens et al.; 2019][Fearnhead et al.; 2018][Power & Goldman; 2019],...

SGD vs. SGP

Gradient flow

Uniform sampling

Markov property

Learning rate

Approximation of deterministic gradient flow

SGD vs. SGP

Gradient flow

SGD update:

$$\theta_k \leftarrow \theta_{k-1} - \eta_k \nabla \Phi_{\boldsymbol{i}_k}(\theta_{k-1})$$

is a forward Euler discretisation of a gradient flow for any realisation $i \in I$

In SGP, we consider the continuous gradient flow:

$$\dot{\theta}(t) = -\nabla \Phi_i(\theta(t))$$

SGD vs. SGP

Uniform sampling

Subsampling in SGD: $\boldsymbol{i}_k \sim \mathrm{Unif}(I)$, for $k \in \mathbb{N}$.

In SGP, we have

$$\boldsymbol{i}(t), \boldsymbol{j}(t) \sim \mathrm{Unif}(I) \qquad (t \geq 0).$$

- implied by the specific choice of transition rate matrices $A, B(\cdot)$

SGD vs. SGP

Markov property

SGD generates a discrete-time Markov chain $(\theta_k, \boldsymbol{i}_k)_{k=0}^{\infty}$

- SGPC generates a continuous-time Markov process $(\theta(t), \boldsymbol{i}(t))_{t\geq 0}$
- SGPD generates a continuous-time Markov process $(\xi(t), \boldsymbol{j}(t))_{t\geq 0}$

SGD vs. SGP

Learning rate

SGD: waiting time between switches equals η_k, i.e.

- switch with rate $1/\eta_k$,

In SGP, we switch with random waiting times $T \sim \mu_{\mathrm{wt}}(\cdot|t_0)$

- look at rate of events happening at time $u \geq 0$ (= hazard function) instead:

$$\nu(u + t_0) = \lim_{d \to 0} \frac{\mathbb{P}(u \leq T \leq u + d | T \geq u)}{d}$$

- SGPC: $\nu(u + t_0) = \frac{1}{\eta}$,
- SGPD: $\nu(u + t_0) = \frac{1}{H(u+t_0)}$.

SGD vs. SGP

Approximation of deterministic gradient flow

SGD with constant learning rate $\eta \approx 0$ approximates the 'exact' gradient flow

$$\frac{\mathrm{d}\zeta}{\mathrm{d}t} = -\nabla\bar{\Phi}(\zeta(t)), \qquad \zeta(0) = \theta_0.$$

Intuition:

- Euler scheme converges $\Rightarrow$ gradient flow
- law of large numbers (LLN):

$$\theta_k = \theta_0 - \big(\eta\nabla\Phi_{\boldsymbol{i}_1}(\theta_0) + \cdots + \eta\nabla\Phi_{\boldsymbol{i}_k}(\theta_{k-1})\big) \overset{(\eta\approx 0)}{\approx} \theta_0 - \underbrace{\big(\eta\nabla\Phi_{\boldsymbol{i}_1}(\theta_0) + \cdots + \eta\nabla\Phi_{\boldsymbol{i}_k}(\theta_0)\big)}_{\overset{\text{LLN}}{\approx}\, \eta k\bar{\Phi}(\theta_0)}$$

SGD vs. SGP

SGPC, with $\eta \approx 0$, also approximates the 'exact' gradient flow

Assumption 2.6. *For any $i \in I$, let $\Phi_i \in C^2(X;\mathbb{R})$ and let $\nabla\Phi_i, \mathrm{H}\Phi_i$ be continuous and bounded on bounded subsets of X.*

Theorem 2.7.

Let $\zeta(0) = \theta(0)$ and let Assumption 2.6 hold, then $(\theta(t))_{t\geq 0} \to (\zeta(t))_{t\geq 0}$, weakly in $C^0([0,\infty);X)$, as $\eta \downarrow 0$.

Proof. Perturbed test function theory of [Kushner; 1984]. □

SGD vs. SGP

Example. Let $\Phi_1(\theta) := (\theta - 1)^2/2$ and $\Phi_2(\theta) := (\theta + 1)^2/2$. $\Rightarrow \bar{\Phi}(\theta) = (\theta^2 + 1)/2$.



Figure: Exemplary realisations of SGPC and plot of precise gradient flow. Discretisation with ode45.

Stochastic gradient descent in nature

Is there an interpretation of SGD in terms of a physical model?

(Langevin) diffusion process: molecular dynamics
SGP: bet-hedging in clonal populations

- some clonal populations (bacteria, fungi,...) live in randomly changing environments
- diversified bet-hedging:
  - organisms have different phenotypes
  - development/switching of phenotypes based on (or independent of) current environment
- [Kussell & Leibler; 2005] represent number of organisms with phenotype $i \in I$ at time $t \geq 0$ by $\theta_i(t)$, where
$$\dot{\theta}(t) = A_{\boldsymbol{i}(t)}\theta(t) \qquad (t > 0)$$
where $\boldsymbol{i}(t) \in I$ is the environment at time t and A_i represent growth in environment i.
- [Sasaki & Ellner; 1995] discuss bet-hedging strategies that are evolutionary stable

Outline

Motivation

Modelling Stochastic Gradient Descent in Continuous Time

Longtime behaviour

- Constant learning rate
- Decreasing learning rate

Practical considerations and illustrations

Conclusions

Long-time behaviour of the Stochastic Gradient Process

Study long-time behaviour of the stochastic gradient processes, i.e., study

$$\mathbb{P}(\theta(t) \in \cdot), \qquad \mathbb{P}(\xi(t) \in \cdot) \qquad\qquad (t \gg 0 \text{ very large}).$$

- existence and uniqueness of stationary measures
- convergence to stationary measures and its speed
- SGPD: convergence to $\delta(\cdot - \theta^*)$, where $\theta^* \in \operatorname{argmin}_{\theta \in X} \bar{\Phi}(\theta)$

Preliminaries

Wasserstein distance

Let $q \in (0,1]$. Consider Wasserstein distance between $\pi, \pi' \in \mathrm{Prob}(X)$:

$$d_{\mathrm{W}}(\pi, \pi') := \inf_{H \in \mathrm{Coup}(\pi,\pi')} \int_{X \times X} \min\{1, \|\theta - \theta'\|_2^q\} H(\mathrm{d}\theta, \mathrm{d}\theta'),$$

$$\mathrm{Coup}(\pi, \pi') := \{G \in \mathrm{Prob}(X^2) : \quad G(\cdot \times X) = \pi, \quad G(X \times \cdot) = \pi'\}$$

- metrises weak convergence, i.e.

$$d_{\mathrm{W}}(\pi_n, \pi) \to 0, \text{ as } n \to \infty \qquad \Leftrightarrow \qquad \pi_n \to \pi, \text{ weakly, as } n \to \infty$$

Preliminaries

Assumption 2.6. *For any $i \in I$, let $\Phi_i \in C^2(X;\mathbb{R})$ and let $\nabla\Phi_i, \mathrm{H}\Phi_i$ be continuous and bounded on bounded subsets of X.*

Assumption 3.1 *There is some $\kappa > 0$, with*

$$\langle \theta_0 - \theta_0', \nabla\Phi_i(\theta_0) - \nabla\Phi_i(\theta_0') \rangle \geq \kappa \|\theta_0 - \theta_0'\|^2 \qquad (\theta_0, \theta_0' \in X, i \in I),$$

i.e. Φ_i are strongly convex for $i \in I$.

Constant learning rate

Theorem 3.3.

Let Assumptions 2.6 and 3.1 hold. Then, $(\theta(t), \boldsymbol{i}(t))_{t>0}$ has a unique stationary measure π_{C} on $(X \times I, \mathcal{B}X \otimes 2^I)$. Moreover, there exist $\kappa', c > 0$ and $q \in (0, 1]$, with

$$d_{\mathrm{W}}(\pi_{\mathrm{C}}(\cdot \times I), \mathbb{P}(\theta(t) \in \cdot | \theta_0, i_0)) \leq c \exp(-\kappa' t) \left(1 + \sum_{i \in I} \int_X \|\theta_0 - \theta'\|^q \pi_{\mathrm{C}}(\mathrm{d}\theta' \times \{i\})\right)$$

$$(i_0 \in I, \theta_0 \in X).$$

Constant learning rate

Theorem 3.3.

Let Assumptions 2.6 and 3.1 hold. Then, $(\theta(t), \boldsymbol{i}(t))_{t>0}$ has a unique stationary measure π_{C} on $(X \times I, \mathcal{B}X \otimes 2^I)$. Moreover, there exist $\kappa', c > 0$ and $q \in (0,1]$, with

$$d_{\mathrm{W}}(\pi_{\mathrm{C}}(\cdot \times I), \mathbb{P}(\theta(t) \in \cdot | \theta_0, i_0)) \leq c \exp(-\kappa' t) \left(1 + \sum_{i \in I} \int_X \|\theta_0 - \theta'\|^q \pi_{\mathrm{C}}(\mathrm{d}\theta' \times \{i\})\right) \quad (i_0 \in I, \theta_0 \in X).$$

- d_{W} depends on q
- convergence with exponential speed
- proof based on results by [Benaïm et al.; 2012][Cloez & Hairer; 2015]
- convexity assumption can be weakened (needs Hörmander Bracket condition)
- finding an analytical expression for π_{C} is probably hard

Decreasing learning rate

Theorem 3.4.

Let Assumptions 2.6 and 3.1 hold. Then, there is a function $\alpha : [0,1) \to [0,\infty)$ that is continuous at 0 and satifies $\alpha(0) = 0$. Moreover, for any $\varepsilon > 0$, we have a unique probability measure π_ε and constants $\kappa', c > 0$, $q \in (0,1]$ such that

$$d_{\mathrm{W}}(\delta(\cdot - \theta^*), \mathbb{P}(\xi(t) \in \cdot | \xi_0, j_0)) \leq c \exp(-\kappa' t) \left(1 + \sum_{j \in I} \int_X \|\xi_0 - \xi''\|^q \pi_\varepsilon(\mathrm{d}\xi'' \times \{j\}) \right) + \alpha(\varepsilon) \quad (j_0 \in I, \xi_0 \in X).$$

Decreasing learning rate

Theorem 3.4.

Let Assumptions 2.6 and 3.1 hold. Then, there is a function $\alpha : [0,1) \to [0,\infty)$ that is continuous at 0 and satifies $\alpha(0) = 0$. Moreover, for any $\varepsilon > 0$, we have a unique probability measure π_ε and constants $\kappa', c > 0$, $q \in (0,1]$ such that

$$d_{\mathrm{W}}(\delta(\cdot - \theta^*), \mathbb{P}(\xi(t) \in \cdot | \xi_0, j_0)) \leq c \exp(-\kappa' t) \left(1 + \sum_{j \in I} \int_X \|\xi_0 - \xi''\|^q \pi_\varepsilon(\mathrm{d}\xi'' \times \{j\}) \right) + \alpha(\varepsilon) \quad (j_0 \in I, \xi_0 \in X).$$

- Get to a distribution arbitrarily close to $\delta(\cdot - \theta^*)$ at exponential speed
  $\underbrace{\text{SGPD}}_{O(\exp(-t))}$ has faster convergence rates than $\underbrace{\text{SGD}}_{O(1/\sqrt{k})}$
- proof is significantly more involved
  - $(\xi(t), \boldsymbol{j}(t))_{t \geq 0}$ is inhomogeneous in time
  - rate matrix $B(\cdot)$ degenerates, as $t \to \infty$
  - uses results from [Benaïm et al.; 2012][Cloez & Hairer; 2015][Kushner; 1984]

Example: linear least squares

Consider a linear least squares problem, i.e.

$$\bar{\Phi}(\theta_0) := \frac{1}{2} \sum_{i \in I} \| G_i \theta_0 - y_i \|_2^2 / N \qquad (\theta_0 \in X),$$

where $y_i \in Y_i := \mathbb{R}^{M_i}$ and $G_i : X \to Y_i$ is linear for $i \in I$.

- Subsampling using the partial potentials

$$\Phi_i(\theta_0) := \frac{1}{2} \| G_i \theta_0 - y_i \|_2^2 \qquad (\theta_0 \in X, i \in I).$$

- Assumptions 2.6 and 3.1 are satisfied, if

$$G_i^T G_i \text{ is strictly positive definite } (i \in I) \Leftrightarrow \dim X \leq \dim Y_i \text{ and } G_i \text{ has full rank } (i \in I).$$

Example: regularised linear least squares

Consider a regularised linear least squares problem, i.e.

$$\bar{\Phi}(\theta_0) := \frac{1}{2}\sum_{i\in I} \| G_i\theta_0 - y_i\|_2^2/N \quad + \quad \gamma\|\theta_0\|_2^2 \qquad (\theta_0 \in X),$$

where $\gamma > 0$, $y_i \in Y_i := \mathbb{R}^{M_i}$, and $G_i : X \to Y_i$ is linear for $i \in I$.

- Subsampling using the partial potentials

$$\Phi_i(\theta_0) := \frac{1}{2}\| G_i\theta_0 - y_i\|_2^2 \quad + \quad \gamma\|\theta_0\|_2^2 \qquad (\theta_0 \in X, i \in I).$$

- Assumptions 2.6 and 3.1 are satisfied.

Outline

Motivation

Modelling Stochastic Gradient Descent in Continuous Time

Longtime behaviour

Practical considerations and illustrations

- Applying SGP in practice?
- SGD, Stochastic Proximal Point, SVRG, SAG, SAGA,...?
- Illustrations

Conclusions

SGP in practice

Employ SGP in practice:

(i) discretise gradient flows $\dot{\theta}(t) = -\nabla\Phi_i(\theta(t))$, $\theta(0) = \theta_0$ for several $i \in I, \theta_0 \in X$

(ii) discretise CTMPs $(\boldsymbol{i}(t))_{t\geq 0}$, $(\boldsymbol{j}(t))_{t\geq 0}$

SGP in practice

(i) discretise gradient flows $\dot{\theta}(t) = -\nabla\Phi_i(\theta(t))$, $\theta(0) = \theta_0$ for several $i \in I, \theta_0 \in X$

- use numerical time stepping method; easy, right?
  - which time stepping method is particularly suitable for these gradient flows?
  - which time stepping method will lead to similar ergodic behaviour?
  - here: use a high-order adaptive Runge-Kutta scheme (rk45)

(ii) discretise CTMPs $(\boldsymbol{i}(t))_{t\geq 0}$, $(\boldsymbol{j}(t))_{t\geq 0}$

SGP in practice

(i) discretise gradient flows $\dot{\theta}(t) = -\nabla\Phi_i(\theta(t))$, $\theta(0) = \theta_0$ for several $i \in I, \theta_0 \in X$

(ii) discretise CTMPs $(\boldsymbol{i}(t))_{t\geq 0}$, $(\boldsymbol{j}(t))_{t\geq 0}$

- Exact sampling of $(\boldsymbol{i}(t))_{t\geq 0}$, $(\boldsymbol{j}(t))_{t\geq 0}$ using algorithm by [Gillespie; 1977]: needs to sample waiting times from $\pi_{\mathrm{wt}}(\cdot|t_0)$
  - sampling from exponential distribution in case of $(\boldsymbol{i}(t))_{t\geq 0}$
  - more complicated in case of $(\boldsymbol{j}(t))_{t\geq 0}$
- The SGD-way: use fixed waiting times and sample from $\mathrm{Unif}(I)$
  - representation is quite imprecise, but might do the job
  - continuous time modelling step backwards

SGD, Stochastic Proximal Point, SVRG, SAG, SAGA,...?

Retrieving well-known algorithms from SGP

- choose deterministic waiting times in the discretisation of the CTMP
- choose particular time stepping schemes for the gradient flows
  - forward Euler $\Rightarrow$ SGD [Robbins & Monro; 1951]
  - backward Euler $\Rightarrow$ Stochastic Proximal Point [Bertsekas; 2011]
  - forward Euler + control variate (or a multistep method?) $\Rightarrow$ SVRG [Johnson & Zhang; 2013], SAG [Schmidt et al.; 2017], SAGA [Defazio et al.; 2014]
  - higher order scheme $\Rightarrow$ higher order SGD-type method [Song et al.; 2018]

Illustrations

Constant learning rate:

- What is the stationary measure π_C?
- How does it compare to the stationary measure (well-defined?) of SGD?

Decreasing learning rate:

- How does the distribution $\mathbb{P}(\xi(t) \in \cdot)$ evolve over time?

Example 2. Let $N := 3$, i.e. $I := \{1, 2, 3\}$, and $X := \mathbb{R}$. We define the potentials

$$\Phi_1(\theta) := \frac{1}{2}(\theta + 2)^2, \quad \Phi_2(\theta) := \frac{1}{2}(\theta - 1.5)^2, \quad \Phi_3(\theta) := \frac{1}{2}(\theta - 2)^2 \quad (\theta \in X).$$

The minimiser of $\bar{\Phi} \equiv \Phi_1/3 + \Phi_2/3 + \Phi_3/3$ is $\theta^* = 0.5$.

Constant learning rate



Figure: Kernel density estimates of $\mathbb{P}(\theta(10) \in \cdot | \theta(0) = -1.5) \approx \pi_C$ (SGPC) and $\mathbb{P}(\theta_{10/\eta} \in \cdot | \theta_0 = -1.5)$ (SGD) based on $\eta \in \{1, 0.1, 0.01, 0.001\}$ using 10,000 samples each.

Decreasing learning rate

- time-dependent learning rate: $\eta'(t) := \frac{1}{100t+1}$, for $t \in [0, \infty)$
- sample the waiting times in $(\boldsymbol{j}(t))_{t \geq 0}$ using the quantile function of $\pi_{\mathrm{wt}}(\cdot | t_0)$



Decreasing learning rate



Figure: Left: kernel density estimates of $\mathbb{P}(\xi(10) \in \cdot | \xi(0) = -1.5)$ with $t \in \{1/4, 1/2, 1, 2, 4, 10\}$ using 10,000 samples each. Right: Comparison of $\mathbb{P}(\xi(10) \in \cdot | \xi(0) = -1.5)$ and $\mathbb{P}(\theta(10) \in \cdot | \theta(0) = -1.5)$, where $\eta'(10) \approx 0.001 = \eta$

Outline

Motivation

Modelling Stochastic Gradient Descent in Continuous Time

Longtime behaviour

Practical considerations and illustrations

Conclusions

Take-home messages

- we introduced SGP – a continuous-time model for SGD with non-asymptotic learning rates
- captures most properties of SGD
  - gradient flow structure, uniform subsampling, Markov property, learning rates/switching rate, approximates deterministic gradient flows
- has a biological interpretation
- SGPC converges to a unique stationary measure π_{C} at exponential speed
- SGPD converges to (a neighbourhood of) $\delta(\cdot - \theta^*)$ at exponential speed
- SGP models SGD, but also other stochastic optimisation methods

Where do we go from here?

- Develop efficient practical algorithms from SGP
  - ODE solver?
  - Large scale application
- SGD in constrained optimisation
  - impose jumps in $(\boldsymbol{i}(t))_{t\geq 0}$ and $(\boldsymbol{j}(t))_{t\geq 0}$ when hitting the boundary
- SGD in non-convex optimisation could probably explained in a similar fashion
  - learning rate acts similar as a temperature in simulated annealing
  - Hörmander bracket conditions
- introduce subsampling in other continuous-time algorithms
  - Ensemble Kalman inversion [Kovachki & Stuart; 2019][Schillings & Stuart; 2017]...
  - Continuous time methods in imaging [Bertozzi & Flenner; 2012]...
- understand statistical properties of π_{C}
  - seems related to a posterior density

UNIVERSITY OF CAMBRIDGE

Jonas Latz input/output: www.latz.io twitter: @latzplacian