

Series 1. Sep 13th 2020 (Recap IML and important concepts)

Teaching assistant: **Shaoduo Gan**
sgan@inf.ethz.ch

Problem 1 (Regression):

1. **Linear Regression** Given is a dataset of inputs $\mathbf{X} \in \mathbb{R}^{n \times d}$ and output variable $\mathbf{y} \in \mathbb{R}^n$. Recall the statistical model for linear regression in homogeneous coordinates is $\hat{y}_i = \sum_{j=1}^d x_{ij} \beta_j \implies \hat{\mathbf{y}} = \mathbf{X} \beta$.
 - (a) State the residual sum of squares error for this problem in sum and matrix notation.
 - (b) Derive the optimal $\hat{\beta}$ by minimizing the loss stated above.
2. **Ridge Regression** In Ridge Regression, an additional term $\lambda \beta^\top \beta$ is added to the residual sum of squares loss.
 - (a) Explain the term $\lambda \beta^\top \beta$ in the loss function, what is its impact on the solution? What is the role of the design parameter λ , i.e. what happens for $\lambda \rightarrow 0$ and $\lambda \rightarrow \infty$?
 - (b) Derive the optimal $\hat{\beta}$ by minimizing the ridge loss.

Problem 2 (Comprehension Questions):

1. **Overfitting** A dataset is split into training and validation set. Given are the training and the validation loss as a function of the training time for three neural networks. For every graph, ...
 - (a) ... state whether the network is underfitting, reasonable or overfitting and why.
 - (b) ... give an example of what you could do to improve the solution.




Figure 1: Loss diagrams for networks i. (left), ii. (middle) and iii. (right).

2. **Cross Validation** We want to train a Ridge Regression model and estimate its prediction error. We therefore split the available dataset into a training, validation and test set.
 - (a) Explain what the different sets are used for and why the distinction is important.

- (b) You choose to use K-fold cross validation to estimate the prediction error for different values of λ . Briefly describe the procedure and explain its advantage over a single split of the training/validation set.
 - (c) Is there a systematic tendency affecting the obtained error estimate? Discuss the impact of the number K . How would you choose K in general? What happens for $K \rightarrow 1, K \rightarrow n$?
3. **Generative vs. Discriminative Modeling** Contrast the generative and the discriminative modeling approach.
- (a) Which probability distributions are they trying to estimate? Which one is harder?
 - (b) How is the decision boundary derived in each approach? How does this affect outliers?
 - (c) What is the influence of the model specification and available data on the prediction error? Which approach would you prefer? What about unlabeled data?

Problem 3 (EM-Algorithm for Gaussian Mixtures):

Given is a dataset $\mathcal{X} = \{x_i\}_{i=1,\dots,n}, x_i \in \mathbb{R}$, that can be modeled as a mixture of k Gaussian components with parameters $\theta = \{\pi_c, \mu_c, \sigma_c^2\}_{c=1,\dots,k}$, where π_c is the prior probability that a sample is generated by mixture component c , μ_c and $\sigma_c^2 > 0$ are the mean and variance of component c . A local optimum for the parameters θ as well as assignment probabilities $P(c|x_i, \theta) = \gamma_{ic}$ can be computed using the EM algorithm.

We are now going to derive the E- and M-Step update equations that maximize the likelihood of our model.

1. Find an expression for the probability of datum x_i in our model, i.e. $P(x_i|\theta)$.
2. Compute the log likelihood of the dataset $L(\mathcal{X}|\theta) = \log(P(\mathcal{X}|\theta))$.
3. We now introduce the binary latent variable $M_{ic} \in \{0, 1\}$, where $M_{ic} = 1$ indicates that x_i is generated by component c , $M_{ic} = 0$ indicates that x_i is not generated by component c . Find an expression for the joint likelihood $P(\mathcal{X}, M|\theta)$ and $L(\mathcal{X}, M|\theta)$.
4. Next we compute the expectation over the latent variables and introduce the assignment probabilities $\gamma_{ic} := \mathbb{E}_{M|\mathcal{X}, \theta}[M_{ic}]$. Calculate an expression for $Q(\theta) = \mathbb{E}_{M|\mathcal{X}, \theta}[L(\mathcal{X}, M|\theta)]$ as a function of γ_{ic}, π_c and $P(x_i|c, \mu_c, \sigma_c^2)$.
5. **E-step** During the E-step, we keep our current estimate of θ fixed and calculate the expected assignments. Calculate the new assignment probabilities γ_{ic} as an expression of π_c and $P(x_i|c, \mu_c, \sigma_c^2)$.

Hint: Use the definition of γ_{ic} from 4.

6. **M-step** During the M-step, the assignments γ_{ic} are fixed and we optimize $\theta \in \arg \max_{\theta} Q(\theta)$.

- (a) Compute the new estimate of μ_c .

Hint: From the introduction $P(x_i|c, \mu_c, \sigma_c^2)$ is a 1-D Gaussian.

- (b) Compute the new estimate of σ_c^2 .
- (c) Compute the new estimate of π_c s.t. $\sum_{c=1}^k \pi_c = 1$.

Hint: The constrained optimization problem $\min_x f(x)$ s.t. $g(x) = 0$ can be solved by introducing the Lagrange multiplier λ : $\mathcal{L} = f(x) + \lambda g(x)$. In this problem, use $\frac{d}{dx} \mathcal{L} = 0$, then eliminate λ using the constraint $\sum_{c=1}^k \pi_c = 1$ to arrive at the optimizer.