Dirac Equation from the SL(2, $\mathbb{C}$) Pulsation of K_4 in the PDL Framework

C. Laubscher*

27th March 2026

Abstract

The Projective Dynamic Logo (PDL) framework derives physical constants and dynamical laws from four axioms on finite signed graphs. Paper D32 of the corpus established that the non-relativistic Schrödinger equation for an electron in the Coulomb field of a PDL proton follows from the $(A) \wedge (B)$ stability criterion of D29 and the Compton pulsation of K_4, without free parameter. The transition coefficient $b = -i\hbar\Delta t/(2m_e^{\mathrm{PDL}}(\Delta x)^2)$ was derived from the phase inversion $P_2 = -P_1$ imposed by condition (B). The present paper derives the relativistic extension: the Dirac equation for a PDL electron follows from the same condition (B), via an algebraic chain that requires no additional postulate.

Four results are established. *Lemma 1* identifies the half-cycle transition operator T on the two-dimensional pulsation space $\mathcal{H}_{\mathrm{cycl}} \cong \mathbb{C}^2$ from two conditions that are direct translations of the PDL pulsation axiom and condition (B) respectively. *Theorem 1* proves algebraically that $T^2 = -I_2$, with the corollary that the PDL pulsation has period 4 in $\mathcal{H}_{\mathrm{cycl}}$ (four half-cycles per complete cycle in the complex amplitude representation), establishing the spin-$\frac{1}{2}$ structure of K_4 as a theorem rather than a postulate. *Theorem 2* (uniqueness) shows that $T = -i\tau_2$ is the *unique* 2×2 operator satisfying the four PDL pulsation constraints; in particular, τ_1 is excluded by the algebraic condition $\tau_1^2 = +I_2 \neq -I_2$. *Theorem 3* proves that the Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I_4$ for the PDL-motivated matrices $\gamma^0 = \tau_3 \otimes I_2$ and $\gamma^i = i\tau_2 \otimes \sigma_i$ is a theorem of the PDL pulsation axioms, with numerical residual 0.00×10^0. The non-relativistic limit reproduces exactly the Schrödinger–PDL equation of D32. No free parameter enters at any stage.

The document is self-contained with respect to its sole external dependency: a compact verification table for condition (B) is included (Table 1), allowing the reader to verify the key ingredient independently without access to D29.

1 Introduction

The PDL programme reconstructs fundamental physics from four axioms on finite signed graphs [12, 11]. The proton is characterised by the unique integer quintuplet $(n_u, n_d, r_{\mathrm{val}}, R_{\mathrm{sea}}, R_{\mathrm{tot}}) = (24, 28, 930, 10087, 11017)$, from which the fine-structure constant $\alpha_{\mathrm{PDL}}^{-1} = \varphi^2 r_{\mathrm{val}}^2/(9\mu) = 137.022$ (where $\mu = R_{\mathrm{tot}}/R_e = 11017/6$) and the gravitational constant G_{PDL} are derived without free parameter [7, 5, 8].

Paper D29 established that a mixed triangle (e_i, e_j, p_k), coupling two electronic vertices of K_4 to one proton vertex, is stable across both half-cycles of the K_4 pulsation if and only if two conditions hold [4]: condition (A) requires initial alignment of the cross-product P_1 with the K_4 edge sign $s^{(1)}(e_i, e_j)$; condition (B) imposes the exact phase inversion $P_2 = -P_1$. This was verified exhaustively over 768 cases with zero counter-example.

Paper D32 derived the non-relativistic Schrödinger equation from condition (B) and the Compton pulsation of K_4 [14]. The key advance was the identification that condition (B) forces

*Independent researcher, Switzerland. cedric.laubscher@gmail.com

a π phase shift between the two half-cycles of every stable mixed triangle; the coherent sum over one pulsation cycle then fixes $b = -i\hbar\Delta t/(2m_e^{\mathrm{PDL}}(\Delta x)^2)$ without free parameter. Open problem OP4 of D32 identified the relativistic extension — the Dirac equation from the full $\mathrm{SL}(2,\mathbb{C})$ structure of the Compton pulsation — as the next target.

The present paper resolves OP4. The K_4 pulsation between $s^{(1)}$ and $s^{(2)} = -s^{(1)}$ defines, upon representation in $\mathcal{H}_{\mathrm{cycl}} \cong \mathbb{C}^2$, a half-cycle operator T whose algebraic properties are entirely determined by condition (B). We prove that $T^2 = -I_2$ is a theorem, that $T = -i\tau_2$ is uniquely forced among all 2×2 matrices satisfying the PDL constraints, and that the Clifford algebra then follows without additional input. The factorisation $\mathcal{H}_{\mathrm{Dirac}} \cong \mathcal{H}_{\mathrm{cycl}} \otimes \mathcal{H}_{\mathrm{spin}}$ yields a PDL-motivated Dirac equation whose non-relativistic limit is exactly D32.

Section 2 recalls the PDL prerequisites, including a self-contained verification table for condition (B). Section 3 derives T with an explicit remark on the physical origin of the sign in condition (T-b). Sections 4 through 6 establish Theorems 1–3. Section 7 derives the non-relativistic limit. Sections 8 and 9 present numerical checks and open problems.

2 PDL prerequisites

2.1 The K_4 closure and its pulsation

K_4 is the complete signed graph on four vertices $\{0,1,2,3\}$ and six edges $E(K_4) = \{\{i,j\} : i < j\}$. A sign assignment $s^{(1)} : E(K_4) \to \{+1,-1\}$ satisfying triangular coherence ($s^{(1)}_{ij} s^{(1)}_{jk} s^{(1)}_{ik} = +1$ for every triangle) exists and is unique up to the global flip. The *binary pulsation* alternates between $s^{(1)}$ and $s^{(2)} = -s^{(1)}$; both configurations are triangularly coherent, and the two-step cycle is an irreducible internal clock with angular frequency $\omega_c = m_e^{\mathrm{PDL}} c^2/\hbar$ [12]. There are exactly 8 coherent sign configurations of K_4, forming 4 pulsation pairs $\{s^{(1)}, s^{(2)}\} = \{s, -s\}$.

2.2 Condition (B) and its self-contained verification

A mixed triangle (e_i, e_j, p_k), with $e_i, e_j \in V(K_4)$ and p_k a proton vertex, is *stable* across both half-cycles if and only if the cross-product $P_c = s_\times(e_i, p_k, c) \cdot s_\times(e_j, p_k, c)$ at half-cycle $c \in \{1,2\}$ satisfies [4]:

$$\text{(A)} \quad P_1 = s^{(1)}(e_i, e_j), \tag{1}$$

$$\text{(B)} \quad P_2 = -P_1. \tag{2}$$

Table 1 provides a self-contained verification of the logical equivalence $(A) \wedge (B) \Leftrightarrow$ stable for all four sign combinations $(P_1, P_2) \in \{+1,-1\}^2$, together with the counts from the 768 exhaustive cases of D29. A reader without access to D29 can verify the sole external dependency of this paper directly from this table.

The sole relevant output of condition (B) for the present paper is the ratio $P_2/P_1 = -1$, verified in all 192 stable cases. This is the quantity that constrains the half-cycle operator T in Section 3.

2.3 The Dirac spinor factorisation

The PDL Dirac spinor space is factorised as [15]:

$$\mathcal{H}_{\mathrm{Dirac}} \;\cong\; \mathcal{H}_{\mathrm{cycl}} \otimes \mathcal{H}_{\mathrm{spin}}, \tag{3}$$

where $\mathcal{H}_{\mathrm{cycl}} \cong \mathbb{C}^2$ encodes the two half-cycles of the K_4 pulsation (positive and negative energy branches), and $\mathcal{H}_{\mathrm{spin}} \cong \mathbb{C}^2$ encodes the spin-$\frac{1}{2}$ doublet of the $(4,6)$ block. The spin-$\frac{1}{2}$ structure of $\mathcal{H}_{\mathrm{spin}}$ follows from the two logically stationary motifs of the $(4,6)$ closure, related by a discrete symmetry; Born's rule on $\mathcal{H}_{\mathrm{spin}}$ is derived in [13, 10]. The present paper derives the operator structure of $\mathcal{H}_{\mathrm{cycl}}$ from the PDL pulsation axioms; the structure of $\mathcal{H}_{\mathrm{spin}}$ is used only in the tensor product definition of γ^i (Section 6).

Table 1: Self-contained verification of the stability criterion for mixed triangles in the PDL framework. The sign $s^{(1)}(e_i, e_j)$ is fixed at $+1$ without loss of generality (the case -1 is equivalent by global sign flip). Counts are summed over all 8 coherent K_4 configurations, all 6 edges, and all $2^4 = 16$ cross-sign combinations (768 cases total; 192 stable; 0 counter-examples to the equivalence).

P_1	P_2	(A): $P_1 = s^{(1)}$	(B): $P_2 = -P_1$	$(A) \wedge (B)$	Stable
$+1$	$+1$	✓	×	×	No
$+1$	-1	✓	✓	✓	Yes
-1	$+1$	×	✓	×	No
-1	-1	×	×	×	No

$P_2/P_1 = -1$ in all 192 stable cases. $(A) \wedge (B) \Leftrightarrow$ stable: 768/768, 0 counter-examples.

3 The half-cycle operator T

3.1 Representation in $\mathcal{H}_{\text{cycl}}$)

We represent the two pulsation states in an orthonormal basis of $\mathcal{H}_{\text{cycl}}$:

$$s^{(1)} \longleftrightarrow |c_+\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \qquad s^{(2)} = -s^{(1)} \longleftrightarrow |c_-\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{4}$$

The half-cycle operator $T : \mathcal{H}_{\text{cycl}} \to \mathcal{H}_{\text{cycl}}$ advances the system by one half-cycle.

Definition 3.1 (Half-cycle operator)**.** The operator T is defined by two conditions translating the PDL pulsation axiom and condition (B) into $\mathcal{H}_{\text{cycl}}$:

$$(T\text{-}a) \quad T|c_+\rangle = |c_-\rangle, \tag{5}$$

$$(T\text{-}b) \quad T|c_-\rangle = -|c_+\rangle. \tag{6}$$

Condition (T-a) expresses the forward step $s^{(1)} \to s^{(2)}$. Condition (T-b) expresses the return step $s^{(2)} \to s^{(1)}$ with the sign inversion discussed in Remark 3.2 below.

Remark 3.2 (Physical origin of the sign in condition (T-b))**.** The sign -1 in condition (T-b) is not a convention. It is forced by the structure of the PDL pulsation combined with condition (B), as follows.

Consider the two-step cycle $s^{(1)} \to s^{(2)} \to s^{(1)}$ at the level of sign configurations. The return step $s^{(2)} \to s^{(1)}$ appears, classically, to simply restore the initial state. However, $s^{(2)} = -s^{(1)}$: the second configuration carries a global factor -1 relative to the first. When this factor is represented in $\mathcal{H}_{\text{cycl}}$, the return step cannot map $|c_-\rangle$ to $+|c_+\rangle$ without discarding the phase information accumulated during the forward step. The requirement that T *preserves* this phase information — which is the content of unitarity, condition (ii) of Theorem 5.1 — forces $T|c_-\rangle = -|c_+\rangle$.

Equivalently, Table 1 shows that $P_2/P_1 = -1$ in all stable cases. In the amplitude representation, this ratio is the phase factor acquired over one half-cycle. The product over a full cycle $(P_2/P_1) \cdot (P_1/P_2) = (-1) \cdot (-1) = +1$ confirms that the sign configuration returns to $s^{(1)}$ after two steps, while the amplitude acquires a factor $T^2|c_+\rangle = -|c_+\rangle$ (proved in Theorem 4.1). This accumulated factor -1 is the PDL origin of the spin-$\frac{1}{2}$ double-cover: the sign configuration is period-2, but the complex amplitude is period-4.

Lemma 3.3 (Explicit form of T)**.** *The unique matrix on $\mathbb{C}^2$ satisfying Definition 3.1 is:*

$$T = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = -i\tau_2, \tag{7}$$

where $\tau_2 = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix}\right)$.

Proof. Conditions (T-a) and (T-b) fix all four entries of T: the first column is $T|c_+\rangle = |c_-\rangle = (0,1)^T$ and the second column is $T|c_-\rangle = -|c_+\rangle = (-1,0)^T$. Hence $T = \left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right) = -i\tau_2$. □

4 $T^2 = -I_2$ and the period-4 corollary

Theorem 4.1 ($T^2 = -I_2$)**.** *The half-cycle operator T defined by conditions (T-a) and (T-b) satisfies $T^2 = -I_2$.*

Proof. We evaluate T^2 on both basis vectors:

$$T^2|c_+\rangle = T(T|c_+\rangle) \overset{(T\text{-}a)}{=} T|c_-\rangle \overset{(T\text{-}b)}{=} -|c_+\rangle,$$
$$T^2|c_-\rangle = T(T|c_-\rangle) \overset{(T\text{-}b)}{=} T(-|c_+\rangle) = -T|c_+\rangle \overset{(T\text{-}a)}{=} -|c_-\rangle.$$

Since T^2 acts as $-I_2$ on every basis vector of $\mathcal{H}_{\text{cycl}}$, we conclude $T^2 = -I_2$. □

Remark 4.2. Theorem 4.1 uses only the PDL pulsation axiom ($s^{(2)} = -s^{(1)}$, encoded in (T-a)) and condition (B) (encoded in (T-b)). No Clifford algebra, no spinor formalism, and no relativistic postulate is invoked.

Corollary 4.3 (Period-4 pulsation and spin-$\frac{1}{2}$ structure)**.** *The* PDL *pulsation has period 4 in $\mathcal{H}_{\text{cycl}}$: $T^4 = I_2$. A complete return to the initial complex amplitude requires four half-cycles, not two.*

Proof. $T^4 = (T^2)^2 = (-I_2)^2 = +I_2$. □

Remark 4.4 (Sign configurations versus complex amplitudes)**.** The PDL pulsation is period-2 at the level of sign configurations: $s^{(1)} \to s^{(2)} \to s^{(1)}$. The period-4 structure of Corollary 4.3 is a property of the complex amplitude representation in $\mathcal{H}_{\text{cycl}}$. After two half-cycles, the sign configuration returns to $s^{(1)}$, but the amplitude has acquired a factor $T^2 = -I_2$: only after four half-cycles does $T^4 = +I_2$ restore the full state. This is the defining algebraic property of a spin-$\frac{1}{2}$ rotor, here derived from the PDL axioms rather than postulated. The standard statement that "a spinor requires a rotation of 4π to return to itself" has here a precise combinatorial meaning: four half-cycles of the K_4 binary pulsation.

5 Uniqueness of $T = -i\tau_2$

Theorem 5.1 (Uniqueness of T)**.** *Among all operators O on $\mathbb{C}^2$, the unique one satisfying simultaneously:*

(i) $O^2 = -I_2$,

(ii) $O^\dagger O = I_2$ *(unitarity, from preservation of coherence),*

(iii) $O^\dagger = -O$ *(anti-Hermiticity, from time-reversal anti-symmetry),*

(iv) $O|c_+\rangle = |c_-\rangle$ *(forward half-cycle, condition (T-a)),*

is $O = -i\tau_2$.

Proof. Conditions (ii) and (iii) require $O \in i\,\mathfrak{su}(2)$, i.e. $O = i(n_0 I_2 + n_1\tau_1 + n_2\tau_2 + n_3\tau_3)$ for $n_\mu \in \mathbb{R}$ with $n_0^2 + |\mathbf{n}|^2 = 1$. Condition (i) gives $O^2 = -(n_0^2 + |\mathbf{n}|^2)I_2 - 2in_0(\mathbf{n}\cdot\boldsymbol{\tau}) = -I_2$, so the off-diagonal term requires $n_0 = 0$, leaving $|\mathbf{n}| = 1$. Thus $O = i(n_1\tau_1 + n_2\tau_2 + n_3\tau_3)$. Condition (iv) reads $O|c_+\rangle = |c_-\rangle$: evaluating gives $i(n_3,\, n_1 + in_2)^T = (0,1)^T$, which yields $n_3 = 0$ and $i(n_1 + in_2) = 1$, hence $n_1 = 0$ and $n_2 = -1$ (with $|\mathbf{n}| = 1$ confirmed). The unique solution is $O = -i\tau_2$. □

Table 2: Exhaustive check of the four PDL conditions for all canonical candidates. Only $-i\tau_2$ satisfies all four simultaneously. The exclusion of τ_1 via $\tau_1^2 = +I_2 \neq -I_2$ is an algebraic result independent of any convention.

Operator O	$O^2 = -I_2$	$O^\dagger O = I_2$	$O^\dagger = -O$	$O\lvert c_+\rangle = \lvert c_-\rangle$
τ_1	× $(\tau_1^2 = +I_2)$	✓	×	✓
$i\tau_1$	✓	✓	✓	×
τ_2	×	✓	×	×
$i\tau_2$	✓	✓	✓	×
$-i\tau_2$	✓	✓	✓	✓
τ_3	×	✓	×	×
$i\tau_3$	✓	✓	✓	×
iI_2	✓	✓	✓	×

Table 2 verifies this exhaustively for all eight canonical candidates in $\{I_2, \tau_1, \tau_2, \tau_3\} \times \{1, i\}$.

Remark 5.2 (Why τ_1 is excluded)**.** The operator $\tau_1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$ maps $|c_+\rangle \mapsto |c_-\rangle$ correctly (condition (iv) satisfied), but $\tau_1^2 = I_2 \neq -I_2$. It describes a period-2 oscillation in $\mathbb{C}^2$ — a classical bit-flip — not the period-4 pulsation demanded by the PDL axioms. The failure $\tau_1^2 = +I_2$ is algebraic and cannot be remedied by any choice of convention or phase. The PDL pulsation is not a classical oscillator, and τ_1 is excluded by the axioms themselves.

6 Dirac matrices and the Clifford algebra as a theorem

6.1 Construction of the PDL Dirac matrices

Definition 6.1 (PDL Dirac matrices)**.** Using the factorisation $\mathcal{H}_{\text{Dirac}} \cong \mathcal{H}_{\text{cycl}} \otimes \mathcal{H}_{\text{spin}}$ with $T = -i\tau_2$ from Theorem 5.1, the PDL Dirac matrices are:

$$\gamma^0 = \tau_3 \otimes I_2, \qquad \gamma^i = i\tau_2 \otimes \sigma_i \quad (i = 1, 2, 3), \tag{8}$$

where σ_i are the Pauli matrices acting on $\mathcal{H}_{\text{spin}}$. Note that $\gamma^i = (-T) \otimes \sigma_i$: the spatial Dirac matrices are built from the PDL half-cycle operator T, with sign fixed by the Minkowski convention $\eta^{\mu\nu} = \text{diag}(+1, -1, -1, -1)$.

Remark 6.2 (Why $\gamma^0 = \tau_3 \otimes I_2$)**.** The temporal matrix γ^0 uses τ_3 rather than $T = -i\tau_2$. This is forced by the requirement $\{\gamma^0, \gamma^i\} = 0$ (Case 3 of Theorem 6.3), which holds because τ_3 and τ_2 anti-commute for distinct Pauli indices. The operator $\tau_3 = \left(\begin{smallmatrix} +1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$ is the *pulsation sign operator*: it assigns $+1$ to the first half-cycle state $|c_+\rangle$ (positive energy branch) and -1 to the second $|c_-\rangle$ (negative energy branch), providing the correct causal structure for γ^0 in the Minkowski metric $(+, -, -, -)$.

6.2 The Clifford algebra as a PDL theorem

Theorem 6.3 (Clifford algebra from $T^2 = -I_2$)**.** *The* PDL *Dirac matrices of Definition 6.1 satisfy:*

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I_4. \tag{9}$$

Proof. We verify the three independent cases.

Case 1: $\{\gamma^i, \gamma^i\} = -2I_4$.

$$\{\gamma^i, \gamma^i\} = 2(\gamma^i)^2 = 2(i\tau_2)^2 \otimes \sigma_i^2 = 2(-I_2) \otimes I_2 = -2I_4, \tag{10}$$

where $(i\tau_2)^2 = -I_2$ is precisely $T^2 = -I_2$ from Theorem 4.1, and $\sigma_i^2 = I_2$.
Case 2: $\{\gamma^i, \gamma^j\} = 0$ *for* $i \neq j$.

$$\{\gamma^i, \gamma^j\} = (i\tau_2)^2 \otimes \{\sigma_i, \sigma_j\} = (-I_2) \otimes 0 = 0, \tag{11}$$

using $\{\sigma_i, \sigma_j\} = 2\delta_{ij} I_2 = 0$ for $i \neq j$.
Case 3: $\{\gamma^0, \gamma^i\} = 0$.

$$\{\gamma^0, \gamma^i\} = (\tau_3 \cdot i\tau_2 + i\tau_2 \cdot \tau_3) \otimes \sigma_i = \{\tau_3, i\tau_2\} \otimes \sigma_i = 0 \otimes \sigma_i = 0, \tag{12}$$

since $\{\tau_3, i\tau_2\} = i(\tau_3\tau_2 + \tau_2\tau_3) = 0$ (Pauli matrices anti-commute for distinct indices). Assembling the three cases gives equation (9). □

Remark 6.4 (The Clifford algebra is not postulated)**.** Equation (9) is here a *theorem*, not a postulate. Its key ingredient is $T^2 = -I_2$ (Theorem 4.1), which itself follows solely from the PDL axiom $s^{(2)} = -s^{(1)}$ and condition (B). In the standard derivation of the Dirac equation, the Clifford algebra is postulated as the starting point; in the PDL framework, it emerges from the combinatorial structure of K_4.

Proposition 6.5 (Hermiticity)**.** *The* PDL *Dirac matrices satisfy* $(\gamma^0)^\dagger = \gamma^0$ *and* $(\gamma^i)^\dagger = -\gamma^i$.

Proof. $(\gamma^0)^\dagger = \tau_3^\dagger \otimes I_2 = \tau_3 \otimes I_2 = \gamma^0$. For γ^i: $(i\tau_2)^\dagger = -i\tau_2^\dagger = -i\tau_2$, so $(\gamma^i)^\dagger = -i\tau_2 \otimes \sigma_i = -\gamma^i$. □

7 The PDL Dirac equation and its non-relativistic limit

7.1 The PDL Dirac equation

Using the matrices of Definition 6.1, the PDL Dirac equation for an electron in the Coulomb field of a PDL proton is:

$$\left[i\gamma^\mu \left(\partial_\mu + i\frac{q_{\mathrm{PDL}}}{\hbar c} A_\mu \right) - \frac{m_e^{\mathrm{PDL}} c}{\hbar} \right] \Psi = 0, \tag{13}$$

where m_e^{PDL} is fixed by the Compton pulsation $\hbar\omega_c = m_e^{\mathrm{PDL}} c^2$, and q_{PDL} is the PDL charge. For the hydrogenic Coulomb field: $A_0(r) = -\alpha_{\mathrm{PDL}} \hbar c/(q_{\mathrm{PDL}} r)$ and $\mathbf{A} = 0$, with $\alpha_{\mathrm{PDL}} = \mu/R_{\mathrm{surf}}^2$ from D29–D30 [4, 6].

7.2 Non-relativistic limit and consistency with D32

Proposition 7.1 (Dirac–Schrödinger consistency)**.** *In the non-relativistic limit, the large component* ϕ_+ *of the* PDL *spinor* $\Psi = (\phi_+, \phi_-)^T \mathrm{e}^{-i m_e^{\mathrm{PDL}} c^2 t/\hbar}$ *with* $\phi_- \ll \phi_+$ *satisfies:*

$$i\hbar \frac{\partial \phi_+}{\partial t} = \left[-\frac{\hbar^2}{2m_e^{\mathrm{PDL}}} \nabla^2 - \frac{\alpha_{\mathrm{PDL}} \hbar c}{r} \right] \phi_+. \tag{14}$$

*This is exactly the Schrödinger–*PDL *equation of D32 [14], with the same parameters* m_e^{PDL} *and* α_{PDL} *fixed by the proton quintuplet.*

Proof. The decoupling condition $\{\gamma^0, \gamma^i\} = 0$ (Case 3 of Theorem 6.3) guarantees that the large and small components separate in the non-relativistic regime. The block structure $\gamma^0 = \mathrm{diag}(+I_2, -I_2)$ identifies ϕ_+ as the upper two components. Eliminating ϕ_- via $\phi_- \approx \boldsymbol{\sigma} \cdot \hat{\mathbf{p}}/(2m_e^{\mathrm{PDL}} c)\, \phi_+$ and retaining leading-order terms yields equation (14), with coupling $q_{\mathrm{PDL}} A_0(r) = -\alpha_{\mathrm{PDL}} \hbar c/r$. Relativistic corrections at order $1/c^2$ generate the standard Darwin, spin–orbit, and kinematic terms [1]. □

Remark 7.2 (Internal consistency at two relativistic levels)**.** Equation (14) is identical to equation (1) of D32 [14]. The parameters m_e^{PDL} and α_{PDL} are the same in both the relativistic description (D33) and the non-relativistic description (D32), and both are fixed without free parameter by the proton quintuplet $(24, 28, 930, 10087, 11017)$. The PDL programme is self-consistent at two dynamical levels: the non-relativistic hydrogen atom and the relativistic hydrogen atom are governed by the same combinatorial axioms, with no adjustment at the interface.

8 Numerical checks

All numerical checks were performed in Python 3 (NumPy 2.x) and confirmed to machine precision. Every entry of Table 3 was computed independently.

Table 3: Numerical verification of the main algebraic results of D33. All residuals are identically zero to machine precision (float64). Python code available from the author upon request.

Quantity	Expression verified	Max. residual
$T^2 = -I_2$	$\max\lvert T^2 + I_2\rvert$	0.00×10^0
$T^4 = I_2$ (period-4)	$\max\lvert T^4 - I_2\rvert$	0.00×10^0
Unitarity of T	$\max\lvert T^\dagger T - I_2\rvert$	0.00×10^0
Anti-Hermiticity of T	$\max\lvert T^\dagger + T\rvert$	0.00×10^0
Clifford algebra	$\max_{\mu,\nu}\lvert\{\gamma^\mu, \gamma^\nu\} - 2\eta^{\mu\nu} I_4\rvert$	0.00×10^0
$(\gamma^0)^\dagger = \gamma^0$	$\max\lvert\gamma^0 - (\gamma^0)^\dagger\rvert$	0.00×10^0
$(\gamma^i)^\dagger = -\gamma^i$	$\max_i\lvert\gamma^i + (\gamma^i)^\dagger\rvert$	0.00×10^0
$(\gamma^i)^2 = -I_4$	$\max_i\lvert(\gamma^i)^2 + I_4\rvert$	0.00×10^0
$\{\gamma^0, \gamma^i\} = 0$	$\max_i\lvert\{\gamma^0, \gamma^i\}\rvert$	0.00×10^0
τ_1 excluded: $\tau_1^2 = +I_2$	$\max\lvert\tau_1^2 - I_2\rvert$	0.00×10^0

Table 4: Hydrogen energy levels from the PDL Dirac equation (13), with $\alpha_{\mathrm{PDL}}^{-1} = 137.022$ (CODATA 2022: 137.036, deviation 10^{-4}). The fine-structure splitting $\Delta E = E_{\mathrm{Dirac}}^{\mathrm{PDL}} - E_{\mathrm{Bohr}}^{\mathrm{PDL}}$ is at the sub-meV level, consistent with the 10^{-4} precision of α_{PDL}.

n	j	$E_{\mathrm{Dirac}}^{\mathrm{PDL}}$ (eV)	$E_{\mathrm{Bohr}}^{\mathrm{PDL}}$ (eV)	ΔE (eV)
1	$\frac{1}{2}$	-13.608889	-13.608707	-0.000181
2	$\frac{1}{2}$	-3.402233	-3.402177	-0.000057
2	$\frac{3}{2}$	-3.402188	-3.402177	-0.000011
3	$\frac{1}{2}$	-1.512099	-1.512079	-0.000020
3	$\frac{3}{2}$	-1.512085	-1.512079	-0.000007
3	$\frac{5}{2}$	-1.512081	-1.512079	-0.000002

9 Open problems

OP1 (Fully algebraic non-relativistic limit). Proposition 7.1 uses the standard large-component elimination, which invokes the continuum kinetic action $\hbar\omega_c = m_e^{\mathrm{PDL}} c^2$ at an intermediate step (as in D32, OP1). A fully algebraic derivation of the NR limit from the discrete PDL update rules, without the continuum action argument, would make the derivation chain purely combinatorial.

OP2 (Multi-body Dirac extension). Equation (13) covers one K_4 block in the field of one proton. Extension to helium and many-electron atoms requires specifying how two independent

Table 5: Relative spectral precision of the PDL Dirac framework with respect to $\alpha^{-1}_{\text{CODATA}} = 137.036$. The uniform value $|\Delta E|/E = 2.22 \times 10^{-4}$ across all levels confirms that the framework introduces no additional spectral error beyond the known 10^{-4} deviation of α_{PDL}.

n	j	$\lvert E_{\text{PDL}} - E_{\text{CODATA}}\rvert / \lvert E_{\text{CODATA}}\rvert$
1	$\frac{1}{2}$	2.22×10^{-4}
2	$\frac{1}{2}$	2.22×10^{-4}
2	$\frac{3}{2}$	2.22×10^{-4}
3	$\frac{1}{2}$	2.22×10^{-4}
3	$\frac{3}{2}$	2.22×10^{-4}

$(A) \wedge (B)$ criteria interact in $\mathcal{H}^{(1)}_{\text{Dirac}} \otimes \mathcal{H}^{(2)}_{\text{Dirac}}$, and whether Pauli exclusion emerges from the mixed-triangle structure.

OP3 (QED extension). The present treatment covers the electron in an external Coulomb field. A PDL–QED programme would derive the photon sector from the graph structure and yield radiative corrections (Lamb shift, anomalous magnetic moment) from first principles.

OP4 (Connection to $G_{\text{eff}}(N)$). The link between the Dirac coupling α_{PDL} and the gravitational coupling $G_{\text{eff}}(N) = \sigma(N) \cdot G_{\text{PDL}}$ of D31 [9, 8] at the level of the 18-step chain complex ε^{18}_G remains to be established algebraically (OP6 of D32, OP1 of D31).

Conclusion

This paper has derived the PDL Dirac equation from condition (B) of D29 via an algebraic chain requiring no additional postulate. The central result is that the half-cycle operator $T = -i\tau_2$ is the *unique* 2×2 operator satisfying the four PDL pulsation constraints (Theorem 5.1), and that the Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I_4$ follows from this uniqueness as a theorem (Theorem 6.3, numerical residual: 0.00×10^0).

Two results emerge beyond the primary goal. First, the PDL pulsation has period 4 in the complex amplitude representation $\mathcal{H}_{\text{cycl}}$ (Corollary 4.3): the spin-$\frac{1}{2}$ double-cover structure is a consequence of the axioms, not a postulate. The standard statement that a spinor requires a 4π rotation to return to itself corresponds here to four half-cycles of the K_4 binary pulsation. Second, the operator τ_1 is excluded by the algebraic fact $\tau_1^2 = +I_2 \neq -I_2$ (Table 2), showing that the correct Lorentz representation is uniquely selected by the PDL axioms.

The document is self-contained: Table 1 allows the reader to verify the sole external dependency (condition (B) from D29) without access to D29, and Remark 3.2 explains from first principles why the sign -1 in condition (T-b) is forced by the pulsation structure rather than being a convention.

The non-relativistic limit of the PDL Dirac equation reproduces exactly the Schrödinger–PDL equation of D32 (Proposition 7.1), with the same parameters m_e^{PDL} and α_{PDL} fixed by the proton quintuplet $(24, 28, 930, 10087, 11017)$. The PDL programme is self-consistent at two dynamical levels: the non-relativistic and relativistic descriptions of the hydrogen atom both emerge from the same combinatorial axioms, without free parameter.

References

[1] J. D. Bjorken and S. D. Drell. *Relativistic Quantum Mechanics*. McGraw-Hill, New York, 1964.

[2] P. A. M. Dirac. The quantum theory of the electron. *Proceedings of the Royal Society of London A*, 117:610–624, 1928.

[3] P. A. M. Dirac. *The Principles of Quantum Mechanics.* Oxford University Press, Oxford, 4th edition, 1958.

[4] C. Laubscher. Axiomatic derivation of the leading valence engagement amplitude in the PDL framework. *Zenodo*, 2026. PDL corpus, D29. Gate 1 resolved: 155/11017 from PDL axioms alone. Conditions (A) and (B) for mixed-triangle stability; 768 cases, 0 counter-examples.

[5] C. Laubscher. Bridge formula $\delta r_{\mathrm{val}} = \varepsilon_g \cdot r \cdot c$ and the gravitational constant in the PDL framework. *Zenodo*, 2026. PDL corpus, D20-V3. $G_{\mathrm{PDL}} = 6.67448 \times 10^{-11}\,\mathrm{m}^3\,\mathrm{kg}^{-1}\,\mathrm{s}^{-2}$, deviation 27 ppm from CODATA 2022.

[6] C. Laubscher. Coefficient 2 and the isospin mass difference from PDL axioms. *Zenodo*, 2026. PDL corpus, D30. Gate 2 resolved: $\varepsilon_G^B = \kappa C/(6+\kappa)(1+2/R_{\mathrm{tot}}) \in \mathbb{Q}(\sqrt{5})$, 17 ppm from CODATA; $\alpha_{\mathrm{PDL}} = \mu/R_{\mathrm{surf}}^2$ derived.

[7] C. Laubscher. Combinatorial derivation of the proton quintuplet and the fine-structure constant in the PDL framework. *Zenodo*, 2026. PDL corpus, D24. Quintuplet $(24, 28, 930, 10087, 11017)$ unique; $\alpha_{\mathrm{PDL}}^{-1} = \varphi^2 r_{\mathrm{val}}^2/(9\mu) = 137.022$.

[8] C. Laubscher. Effective gravitational coupling $g_{\mathrm{eff}}(n) = \sigma(n) \cdot g_{\mathrm{PDL}}$ from closure density in the PDL framework. *Zenodo*, 2026. PDL corpus, D25. $\sigma(N) = 1 - (1-\kappa)^N$, $\kappa = 310\varphi/11017 \in \mathbb{Q}(\sqrt{5})$.

[9] C. Laubscher. $g_{\mathrm{eff}}(n) = \sigma(n) \cdot g_{\mathrm{PDL}}$: Gate 3 from the linear bridge inversion hypothesis. *Zenodo*, 2026. PDL corpus, D31. Gate 3 resolved: $\Delta r_{\mathrm{val}}(N) = \sigma(N) \cdot \Delta r_{\mathrm{val}}(\infty)$, error $< 10^{-12}$; S_4 transitivity on K_4; physical obstruction.

[10] C. Laubscher. Gleason-type uniqueness of born's rule for spin-$\frac{1}{2}$ in the PDL framework. *Zenodo*, 2026. PDL corpus. Operational axiomatisation of spin-$\frac{1}{2}$ measurements; Gleason-type uniqueness theorem: Born's rule is the unique admissible probability assignment under normalisation, global phase invariance, rotational covariance, branch symmetry, and repeatability.

[11] C. Laubscher. Introduction to the projective dynamic logo framework. *Zenodo*, 2026. PDL corpus, D2.

[12] C. Laubscher. The projective dynamic logo: Physical reality from relational axioms on finite signed graphs. *Zenodo*, 2026. PDL corpus, D1. Four axioms, K_4 as unique minimal closure, $R_e = 6$, Compton pulsation $\hbar\omega_c = m_e c^2$.

[13] C. Laubscher. Relational emergence of born's rule and the golden-ratio active surface in the PDL framework. *Zenodo*, 2026. PDL corpus. Spin-$\frac{1}{2}$ doublet of the $(4, 6)$ block; mixed-triangle measurement scheme; Born probabilities $P_\pm(\theta; a, b) = |a\cos(\theta/2) \pm b\sin(\theta/2)|^2$ in the $\lambda \to \infty$ limit.

[14] C. Laubscher. Schrödinger dynamics from the $(a) \wedge (b)$ coupling criterion in the PDL framework. *Zenodo*, 2026. PDL corpus, D32. OP6 of D31 resolved: $b = -i\hbar\Delta t/(2m_e(\Delta x)^2)$ derived from condition (B) and Compton pulsation, no free parameter.

[15] C. Laubscher. Towards an Einstein–Dirac unification in the PDL framework. *Zenodo*, 2026. PDL corpus, sketch document. Spinor factorisation $\mathcal{H}_{\mathrm{Dirac}} \cong \mathcal{H}_{\mathrm{cycl}} \otimes \mathcal{H}_{\mathrm{spin}}$; emergent metric; coherence tensor; gravitational redshift of hydrogen.