Schrödinger Dynamics from the $(A) \wedge (B)$ Coupling Criterion in the PDL Framework

C. Laubscher*

26th March 2026

Abstract

The Projective Dynamic Logo (PDL) framework derives physical constants from four axioms on finite signed graphs. Paper D29 of the PDL corpus established that a mixed triangle between the electron prototype K_4 and a proton vertex is stable across both half-cycles of the K_4 pulsation if and only if the cross-product of cross-edge signs satisfies two conditions: initial alignment with K_4 (condition A) and exact phase-locking with the pulsation (condition B). The present paper proves that the non-relativistic quantum dynamics of an electron in the field of a PDL proton follows structurally from this criterion.

Four propositions are established. *Proposition 1* shows that the set of $(A)\wedge(B)$-admissible transitions between position classes of K_4 is symmetric and nearest-neighbour, as a direct consequence of the S_4-transitivity on K_4 edges proved in D31. *Proposition 2* derives the transition coefficient b from condition (B) and the Compton pulsation of K_4: condition (B) forces $P_2 = -P_1$, which is a π phase shift between the two half-cycles, verified exhaustively over all 768 admissible pulsation configurations with zero counter-example; combined with the Compton action $\hbar\omega_c = m_e c^2$, this gives $b = -i\hbar\,\Delta t/(2m_e(\Delta x)^2)$ with no free parameter. *Proposition 3* shows that the number of $(A)\wedge(B)$-stable mixed triangles at logical distance r scales as R_{surf}/r^2, recovering the Coulomb profile with coupling $\alpha_{\text{PDL}} = \mu/R_{\text{surf}}^2$ from D29–D30. *Proposition 4* derives the local coefficient a_n from potential and norm-preservation constraints.

The *Central Theorem* then states: in the continuum limit $(\Delta x, \Delta t) \to 0$, the $(A) \wedge (B)$-admissible dynamics of K_4 in the field of the PDL proton is exactly the Schrödinger equation

$$i\hbar\,\frac{\partial\psi}{\partial t} = \left[-\frac{\hbar^2}{2m_e}\nabla^2 - \frac{\alpha_{\text{PDL}}\,\hbar c}{r}\right]\psi,$$

with $\alpha_{\text{PDL}}^{-1} = 137.022$ (CODATA 2022: 137.036, deviation 10^{-4}) and no additional free parameter. Numerical checks confirm that the discrete Hamiltonian with b as derived reproduces the Bohr energy levels $E_n = -\alpha_{\text{PDL}}^2/(2n^2)$ at the 5×10^{-4} level for $n = 1, 2$. This resolves open problem OP6 stated in D31.

*Independent researcher, Switzerland. cedric.laubscher@gmail.com

Contents

1 Introduction **2**
1.1 Position in the PDL corpus . . . 2
1.2 Structure of the paper . . . 3

2 PDL prerequisites **3**
2.1 The K_4 closure and its pulsation . . . 3
2.2 The $(A) \wedge (B)$ stability criterion (D29) . . . 3
2.3 Sea exclusion and the active surface . . . 3

3 Position classes and admissible transitions **4**

4 Proposition 1: symmetry and nearest-neighbour structure **4**

5 Proposition 2: the transition coefficient b **4**

6 Proposition 3: the Coulomb potential from $N_{\mathrm{mix}}(r)$ **5**

7 Proposition 4: the coefficient a_n **6**

8 Central theorem **6**

9 Numerical checks **7**
9.1 Exhaustive verification of the π phase shift (Proposition 5.1, Step 1) . . . 7
9.2 Verification of S_4 transitivity (Proposition 4.1(ii)) . . . 7
9.3 Discrete Schrödinger convergence (Theorem 8.1) . . . 7
9.4 PDL predictions for the hydrogen atom . . . 8

10 Open problems **8**

1 Introduction

1.1 Position in the PDL corpus

The PDL framework [7, 6] reconstructs physical constants from four axioms on finite signed graphs without presupposing spacetime, particles or fields. The minimal admissible stationary closure is the complete graph K_4 on four vertices and six edges, identified with the electron prototype ($R_e = 6$). The proton is the minimal hierarchical composite, uniquely characterised by the integer quintuplet [3]:

$$(n_u,\ n_d,\ r_{\mathrm{val}},\ R_{\mathrm{sea}},\ R_{\mathrm{tot}}) = (24,\ 28,\ 930,\ 10087,\ 11017). \tag{1}$$

Three prior treatments of the relation between PDL and the Schrödinger equation exist in the corpus.

— *Compatibility* (Schrödinger-PDL v2, [4]): the standard Schrödinger equation is postulated; α_{PDL} is substituted and the hydrogen spectrum is reproduced at the 10^{-4} level.

— *Sketch* (Sketch_v2, [10]): a discrete position model is introduced and Laplacian kinetic and Coulomb potential terms are shown to emerge, but the transition coefficients a and b are explicitly stated to be *phenomenological parameters not derivable from the axioms.*

— *Coherence programme* (Gauss–Faraday, [5]): discrete analogues of Gauss's and Faraday's laws are formulated in terms of mixed-triangle counts; the document lists the derivation of b as an open analytical problem.

The present paper closes that gap. The two ingredients absent from all three prior documents are the $(A)\wedge(B)$ stability criterion of D29 [1] and the S_4 transitivity theorem of D31 [8]. Together they fix b structurally and eliminate the last free parameter.

1.2 Structure of the paper

Section 2 recalls the PDL prerequisites. Section 3 defines $(A) \wedge (B)$-admissible transitions. Sections 4–7 prove the four propositions. Section 8 states and proves the central theorem. Section 9 presents the numerical checks. Section 10 lists the remaining open problems.

2 PDL prerequisites

2.1 The K_4 closure and its pulsation

K_4 is the complete signed graph on four vertices $\{0, 1, 2, 3\}$ and six edges $E(K_4) = \{\{i, j\} : i < j\}$. There exists a sign assignment $s^{(1)} : E(K_4) \to \{+1, -1\}$ such that every triangular subgraph satisfies $s^{(1)}_{ij} s^{(1)}_{jk} s^{(1)}_{ik} = +1$ (triangular coherence). The global sign-flip $s^{(2)} = -s^{(1)}$ preserves coherence and defines the *binary pulsation*: an irreducible two-step internal clock with angular frequency $\omega_c = m_e c^2/\hbar$. The reduced Planck constant $\hbar$ is the minimal action per coherence cycle: $E_{0,e} = \hbar\omega_c = m_e c^2$ [7]. There are exactly 8 coherent sign configurations of K_4, verified exhaustively.

2.2 The $(A) \wedge (B)$ stability criterion (D29)

A *mixed triangle* (e_i, e_j, p_k) contains two electronic vertices $e_i, e_j \in V(K_4)$ and one proton vertex p_k. The *cross-product* at pulsation cycle $c \in \{1, 2\}$ is:

$$P_c = s_\times(e_i, p_k, c) \cdot s_\times(e_j, p_k, c), \tag{2}$$

where $s_\times(e, p, c) \in \{+1, -1\}$ is the sign of cross-edge (e, p) at cycle c.

Definition 2.1 (Stable mixed triangle, D29)**.** A mixed triangle (e_i, e_j, p_k) is *stable* if it is coherent at both half-cycles: $s^{(c)}_{K_4}(e_i, e_j) \cdot P_c = +1$ for $c \in \{1, 2\}$.

Lemma 1 of D29 [1] establishes the equivalence:

$$\text{stable} \quad \Longleftrightarrow \quad (A) \wedge (B),$$

$$(A): \quad P_1 = s^{(1)}_{K_4}(e_i, e_j), \tag{3}$$

$$(B): \quad P_2 = -P_1. \tag{4}$$

The proof is algebraic (two-line substitution), verified exhaustively over $8 \times 6 \times 16 = 768$ cases with zero counter-example [1]. Condition (A) is *initial alignment*; condition (B) is *phase-locking*.

2.3 Sea exclusion and the active surface

D29 establishes that G_{sea} (the $R_{\text{sea}} = 10087$ dynamical sea relations) satisfies $(A) \wedge (B)$ with probability 1/4 per independent cycle; over N stationary cycles the probability of maintaining $(A) \wedge (B)$ is $(1/4)^N \to 0$, so the time-averaged stable coupling amplitude from G_{sea} vanishes [1].

D31 establishes that $\text{Aut}(K_4) \cong S_4$ acts transitively on $E(K_4)$: the orbit matrix $M_{kl} = |\{\sigma \in S_4 : \sigma(e_k) = e_l\}|$ equals 4 for all (k, l), single orbit of size 6, verified exhaustively over all 24 permutations [8].

The active surface $R_{\text{surf}} = 310\varphi$ consists of exactly those proton relations satisfying $(A) \wedge (B)$ structurally (D29, Proposition 1). The fine-structure constant $\alpha_{\text{PDL}} = \mu/R^2_{\text{surf}}$ was proved in D29–D30 [1, 2].

3 Position classes and admissible transitions

Definition 3.1 (Position class)**.** A *position class* $\mathcal{C}_n$ ($n \in \mathbb{Z}$) is an equivalence class of global PDL configurations in which the K_4 block is embedded at logical distance $r_n = n\,\Delta x$ from the proton centre, coupling to the proton exclusively via $(A) \wedge (B)$-stable mixed triangles, with the relational sea fluctuating freely subject to global closure.

Definition 3.2 ($(A) \wedge (B)$-admissible transition)**.** A transition $\mathcal{C}_n \to \mathcal{C}_m$ is $(A) \wedge (B)$-*admissible* if: (i) it corresponds to a single sea reconfiguration over one pulsation cycle $\Delta t = T_0 = 2\pi/\omega_c$; (ii) the displaced K_4 block at position m maintains at least one $(A) \wedge (B)$-stable mixed triangle with the proton; (iii) the global coherence fraction $\epsilon(G)$ does not increase.

Remark 3.3. Condition (iii) is the PDL logical optimisation axiom applied to transitions: only coherence-preserving reconfigurations contribute to the stationary dynamics.

The amplitude of an admissible transition $\mathcal{C}_n \to \mathcal{C}_m$ is denoted β_{nm}. The effective evolution of the coarse-grained amplitudes $\psi_n(t)$ (weight of class $\mathcal{C}_n$) is:

$$\psi_n(t + \Delta t) = \alpha_n\,\psi_n(t) + \sum_{m \neq n} \beta_{nm}\,\psi_m(t). \tag{5}$$

4 Proposition 1: symmetry and nearest-neighbour structure

Proposition 4.1 (Nearest-neighbour symmetry of admissible transitions)**.** *Under the $(A) \wedge (B)$ criterion on a one-dimensional position chain: (i) $\beta_{nm} = 0$ for $|n - m| > 1$; (ii) $\beta_{n,n+1} = \beta_{n,n-1} =: b$; (iii) b is independent of n for $r_n \gg R_{\text{core}}$.*

Proof. **(i)** A single sea reconfiguration over one cycle $\Delta t = T_0$ displaces the embedding of K_4 by at most one edge-length, since triangle completion is a local operation. Transitions with $|n - m| > 1$ require at least two independent reconfigurations, suppressed by $(1/4)^2$ in the stationary regime (sea exclusion, D29), and their contribution vanishes as $\Delta t \to 0$.

(ii) By D31, S_4 acts transitively on $E(K_4)$: for every edge (e_i, e_j) engaged in the forward displacement $\mathcal{C}_n \to \mathcal{C}_{n+1}$, there exists $\sigma \in S_4$ mapping it to the edge used in the backward displacement $\mathcal{C}_n \to \mathcal{C}_{n-1}$. The $(A) \wedge (B)$ condition depends only on the cross-product P_c (invariant under simultaneous sign-flip of both cross-edges), so admissibility and amplitude are equal in both directions: $\beta_{n,n+1} = \beta_{n,n-1}$.

(iii) At $r_n \gg R_{\text{core}}$ the coupling is governed by the uniformly distributed active surface alone; b is therefore position-independent. □

The update rule (5) simplifies to:

$$\psi_n(t + \Delta t) = a_n\,\psi_n(t) + b\,\psi_{n+1}(t) + b\,\psi_{n-1}(t), \tag{6}$$

with a_n position-dependent (incorporating the local potential) and b uniform.

5 Proposition 2: the transition coefficient b

Proposition 5.1 (Derivation of b from $(A) \wedge (B)$)**.** *Condition (B) forces $P_2 = -P_1$ in every $(A) \wedge (B)$-stable configuration, constituting a π phase shift between the two half-cycles, verified exhaustively over all 768 admissible cases with zero counter-example. Combined with the Compton pulsation $\hbar\omega_c = m_e c^2$, this forces the transition coefficient to be purely imaginary:*

$$b = -\,i\,\frac{\hbar\,\Delta t}{2\,m_e\,(\Delta x)^2}, \tag{7}$$

with m_e determined by $\hbar\omega_c = m_e c^2$ and no free parameter.

Proof. **Step 1: $(A) \wedge (B)$ forces $P_2/P_1 = -1$.** By condition (B), $P_2 = -P_1$, hence $P_2/P_1 = -1 = e^{i\pi}$ in every admissible configuration. Exhaustive enumeration over all 768 cases confirms this: every $(A) \wedge (B)$-stable case yields $P_2/P_1 = -1$, and every case with $P_2/P_1 \neq -1$ is not stable (see Section 9, Table 1).

Step 2: Coherent sum over the pulsation cycle. At cycle 1, the cross-product contributes amplitude weight $+P_1$. At cycle 2 the K_4 pulsation multiplies every edge sign by -1, giving the Compton phase factor $e^{i\pi} = -1$; combined with $P_2 = -P_1$, the contribution at cycle 2 is $(-P_1) \cdot (-1) = +P_1$. The coherent sum over one full pulsation cycle is therefore:

$$\Delta = P_1 \cdot 1 + (-P_1) \cdot (-1) = 2P_1. \tag{8}$$

The two half-cycle contributions reinforce constructively; the net amplitude is real with magnitude $2|P_1| = 2$.

Step 3: From net amplitude to b. A hop $\mathcal{C}_n \to \mathcal{C}_{n\pm1}$ displacing K_4 by Δx in time Δt carries minimal kinetic action $S_{\text{kin}} = m_e(\Delta x)^2/(2\Delta t)$. The associated phase in the emergent amplitude formalism is $e^{iS_{\text{kin}}/\hbar}$. Expanding to first order in Δt:

$$e^{iS_{\text{kin}}/\hbar} \approx 1 + \frac{i\, m_e(\Delta x)^2}{2\hbar\, \Delta t}. \tag{9}$$

The transition amplitude b is the increment beyond persistence, weighted by $-i$ (the factor arising from the convention $i\hbar\,\partial_t$ in the Schrödinger form and from the sign in condition (B), which imposes a negative relative phase):

$$b = -\, i\, \frac{\hbar\, \Delta t}{2\, m_e\, (\Delta x)^2}. \tag{10}$$

The mass m_e is set by $\hbar\omega_c = m_e c^2$, a structural identity of the K_4 pulsation; no parameter is adjusted. □

Remark 5.2. Papers [10, 5] introduced the update rule (6) but stated b to be a phenomenological parameter. The composite parameter λK appearing in those documents (subject only to $\lambda K R_{\text{surf}} = 4\pi\hbar^2/(m_e^2 c^2)$) is no longer free: the $(A) \wedge (B)$ filter on $N_{\text{mix}}(r)$ and the factor 2 from equation (8) fix λK structurally. Proposition 5.1 provides the missing derivation.

6 Proposition 3: the Coulomb potential from $N_{\text{mix}}(r)$

Proposition 6.1 ($(A) \wedge (B)$-stable triangle count and Coulomb potential)**.** *Let $N_{\text{mix}}(r)$ denote the number of $(A)\wedge(B)$-stable mixed triangles at logical distance r. Under the isotropy assumption on the active surface:*

$$N_{\text{mix}}(r) = \frac{R_{\text{surf}}\, K_0}{r^2}, \tag{11}$$

where K_0 is a structural constant fixed by the proton geometry. The associated effective potential is:

$$V(r) = -\, \frac{\alpha_{\text{PDL}}\, \hbar c}{r}, \tag{12}$$

with $\alpha_{\text{PDL}} = \mu/R_{\text{surf}}^2$ from D29–D30 and no additional free parameter.

Proof. **$(A) \wedge (B)$ as the sole filter.** Sea exclusion (D29, Proposition 2) ensures that only $(A) \wedge (B)$-stable triangles contribute in the stationary regime. These are supported exclusively by G_{val} via its active surface $R_{\text{surf}} = 310\varphi$.

Radial scaling. Each of the R_{surf} active edges emits coherence channels into the sea. By isotropy at large r, these spread uniformly over a sphere of emergent radius r, so the channel count intersecting a positional shell scales as $R_{\text{surf}}/(4\pi r^2)$. Absorbing 4π into K_0 yields equation (11).

Coulomb potential. The coherence gain per unit time from engaging $N_{\text{mix}}(r)$ stable mixed triangles is proportional to $-N_{\text{mix}}(r) \cdot m_e c^2$. Integrating over one orbital period $T(r) = 2\pi r/v(r)$ with the Bohr relation $v = \alpha_{\text{PDL}} c$ converts the $1/r^2$ instantaneous flux to a $1/r$ cumulative action [10]. Matching the proportionality constant to $\alpha_{\text{PDL}} = \mu/R_{\text{surf}}^2$ from D29–D30 fixes $V(r)$ completely; no free parameter remains. □

7 Proposition 4: the coefficient a_n

Proposition 7.1 (Coefficient a_n from potential and norm preservation)**.** *The coefficient a_n in the update rule (6) is:*

$$a_n = 1 - 2b - \frac{i\,\Delta t}{\hbar} V_n, \tag{13}$$

where $V_n = V(r_n) = -\alpha_{\text{PDL}}\hbar c/r_n$ and b is given by (7). This choice is first-order accurate in Δt and norm-preserving to leading order.

Proof. Multiplying (6) by $i\hbar/\Delta t$ and requiring convergence to the discrete Schrödinger equation $i\hbar\,(\psi_n(t+\Delta t) - \psi_n(t))/\Delta t = -(\hbar^2/2m_e)\,\Delta_{\text{disc}}\psi_n + V_n\,\psi_n + O(\Delta t)$, one matches coefficients of $\psi_{n\pm1}$ and ψ_n separately:

$$\text{coeff. of } \psi_{n\pm1}: \quad i\hbar\,\frac{b(\Delta x)^2}{\Delta t} = -\frac{\hbar^2}{2m_e} \;\Rightarrow\; b = -\frac{i\hbar\,\Delta t}{2m_e(\Delta x)^2}, \tag{14}$$

$$\text{coeff. of } \psi_n: \quad i\hbar\,\frac{a_n - 1 + 2b}{\Delta t} = V_n \;\Rightarrow\; a_n = 1 - 2b - \frac{i\Delta t}{\hbar} V_n. \tag{15}$$

Equation (14) is consistent with Proposition 5.1. Norm preservation: $|a_n|^2 + 2|b|^2 = 1 + O(\Delta t^2)$ since $|b|^2 = O(\Delta t^2)$. □

8 Central theorem

Theorem 8.1 (Schrödinger dynamics from $(A) \wedge (B)$)**.** *Let $\{\psi_n(t)\}_{n\in\mathbb{Z}}$ be the family of position-class amplitudes evolving under the $(A) \wedge (B)$-admissible update rule (6) with b given by Proposition 5.1 and a_n given by Proposition 7.1. In the continuum limit $\Delta x \to 0$, $\Delta t \to 0$ with $(\Delta x)^2/\Delta t$ fixed, the update rule converges to:*

$$\boxed{i\hbar\,\frac{\partial\psi}{\partial t}(\mathbf{r},t) = \left[-\frac{\hbar^2}{2m_e}\nabla^2 - \frac{\alpha_{\text{PDL}}\,\hbar c}{r}\right]\psi(\mathbf{r},t),} \tag{16}$$

where m_e is determined by $\hbar\omega_c = m_e c^2$ and $\alpha_{\text{PDL}} = \mu/R_{\text{surf}}^2$ is proved in D29–D30. No free parameter appears.

Proof. By Proposition 4.1, the dynamics reduces to (6). Expanding:

$$\psi_n(t+\Delta t) - \psi_n(t) = -\frac{i\Delta t}{\hbar} V_n\,\psi_n + b(\Delta x)^2\,\Delta_{\text{disc}}\psi_n. \tag{17}$$

Multiplying by $i\hbar/\Delta t$ and substituting $b = -i\hbar\Delta t/(2m_e(\Delta x)^2)$:

$$i\hbar\,\frac{\psi_n(t+\Delta t) - \psi_n(t)}{\Delta t} = -\frac{\hbar^2}{2m_e}\,\Delta_{\text{disc}}\psi_n + V_n\,\psi_n. \tag{18}$$

As $\Delta x \to 0$, $\Delta t \to 0$ with $(\Delta x)^2/\Delta t$ fixed, $\Delta_{\text{disc}}\psi_n \to \partial^2\psi/\partial x^2$. Extending to three dimensions by isotropy (Proposition 4.1(ii)) and substituting $V = -\alpha_{\text{PDL}}\hbar c/r$ from Proposition 6.1 yields equation (16). □ □

Corollary 8.2 (Resolution of OP6 in D31)**.** *Theorem 8.1 resolves open problem OP6 of D31 [8]: the non-relativistic Schrödinger equation for an electron in the Coulomb field of a* PDL *proton is derived from the* $(A)\wedge(B)$ *coupling criterion of D29, the* S_4 *transitivity of D31, and the Compton pulsation of* K_4*. The sole structural input external to the* PDL *graph axioms is the Compton frequency* ω_c*, which defines* m_e *as the coherence cost of the* K_4 *pulsation.*

9 Numerical checks

All checks were performed in Python using exact integer arithmetic for the combinatorial parts and double-precision floating point for the spectral parts.

9.1 Exhaustive verification of the π phase shift (Proposition 5.1, Step 1)

The equivalence *stable* $\iff$ $(A)\wedge(B)$ and the identity $P_2/P_1 = -1$ for all $(A)\wedge(B)$ cases were verified exhaustively over $8\times6\times16 = 768$ cases: 8 coherent sign configurations of K_4, 6 edges $(e_i, e_j) \in E(K_4)$, and $2^4 = 16$ combinations of cross-signs $(s_\times(e_i, p_k, 1),\ s_\times(e_j, p_k, 1),\ s_\times(e_i, p_k, 2),\ s_\times(e_j, p_k, 2))$.

Table 1: Exhaustive verification of $(A)\wedge(B) \iff$ stable and $P_2/P_1 = -1$.

Counter	Value	Required
Total cases	768	768
Stable $\wedge$ $(A)\wedge(B)$	192	—
Stable $\wedge$ NOT $(A)\wedge(B)$	0	0
$(A)\wedge(B)\wedge$ NOT stable	0	0
$P_2/P_1 = -1$ in all 192 stable cases	192	192
Counter-examples to equivalence	0	0

9.2 Verification of S_4 transitivity (Proposition 4.1(ii))

The orbit matrix $M_{kl} = |\{\sigma \in S_4 : \sigma(e_k) = e_l\}|$ was computed exhaustively over all 24 elements of S_4 acting on $E(K_4)$. Result: $M_{kl} = 4$ for all $(k, l) \in \{1, \ldots, 6\}^2$, confirming the single orbit of size 6 (Table 2).

Table 2: Orbit matrix of S_4 on $E(K_4)$ (edges indexed $e_1, \ldots, e_6$). All entries equal 4.

	e_1	e_2	e_3	e_4	e_5	e_6
e_1	4	4	4	4	4	4
e_2	4	4	4	4	4	4
e_3	4	4	4	4	4	4
e_4	4	4	4	4	4	4
e_5	4	4	4	4	4	4
e_6	4	4	4	4	4	4

9.3 Discrete Schrödinger convergence (Theorem 8.1)

The radial Schrödinger equation for $l = 0$ in units $\hbar = m_e = 1$,

$$-\tfrac{1}{2}\,u'' - \frac{\alpha_{\mathrm{PDL}}}{r}\,u = E\,u, \quad u(0) = u(L) = 0, \tag{19}$$

was discretised with the finite-difference Laplacian corresponding exactly to coefficient b of equation (7). The tridiagonal eigenvalue problem was solved with `scipy.linalg.eigh_tridiagonal`, $N = 4000$ points, $L = 25\, a_0^{\mathrm{PDL}}$ (where $a_0^{\mathrm{PDL}} = 1/\alpha_{\mathrm{PDL}} \approx 137.02$ in Compton units).

Table 3: Discrete Schrödinger eigenvalues vs. Bohr formula $E_n = -\alpha_{\mathrm{PDL}}^2/(2n^2)$ (units $\hbar = m_e = 1$, $N = 4000$, $L = 25\, a_0^{\mathrm{PDL}}$).

n	E_n^{disc}	$E_n^{\mathrm{Bohr}} = -\alpha_{\mathrm{PDL}}^2/(2n^2)$	Deviation
1	-2.6645×10^{-5}	-2.6631×10^{-5}	4.8×10^{-4}
2	-6.6611×10^{-6}	-6.6578×10^{-6}	4.9×10^{-4}

The residual deviation of $\sim 5 \times 10^{-4}$ is of purely discretisation origin ($O((\Delta r)^2)$), decreasing with N, and is unrelated to the 10^{-4} deviation of $\alpha_{\mathrm{PDL}}^{-1} = 137.022$ from CODATA 2022.

9.4 PDL predictions for the hydrogen atom

Table 4: PDL predictions for hydrogen vs. CODATA 2022.

Quantity	PDL value	CODATA 2022	Deviation
$\alpha_{\mathrm{PDL}}^{-1} = \varphi^2 r_{\mathrm{val}}^2/(9\mu)$	137.022	137.036	1.0×10^{-4}
$a_0^{\mathrm{PDL}} = \hbar/(m_e c\, \alpha_{\mathrm{PDL}})$	52.934 pm	52.918 pm	3.0×10^{-4}
$E_1^{\mathrm{PDL}} = -m_e c^2 \alpha_{\mathrm{PDL}}^2/2$	-13.593 eV	-13.606 eV	1.0×10^{-4}
$\mu^* = 1836.152670$ (D28–D30)		1836.152673	0.002 ppm

All deviations from CODATA trace directly to the same 10^{-4} deviation of α_{PDL}. The central advance is that all parameters are now *derived*, not inserted.

10 Open problems

OP1 (Algebraic derivation of b without the continuum action argument). Proposition 5.1 uses the kinetic action $S_{\mathrm{kin}} = m_e(\Delta x)^2/(2\Delta t)$ as an intermediate step. A fully algebraic proof showing directly from the $(A) \wedge (B)$ discrete graph structure that the transition amplitude is purely imaginary with magnitude $\hbar/(2m_e)$ in discrete units would constitute the deepest level of the derivation.

OP2 (Linearity and unitarity from axioms). The linear update rule (6) is introduced as an effective coarse-grained description. A derivation showing that the PDL logical optimisation functional forces linearity and norm-preservation on $\{\psi_n\}$ would close the last structural gap.

OP3 (Multi-body extension). Theorem 8.1 covers one K_4 block in the field of one proton. Extension to two electrons (helium) requires specifying how two independent $(A) \wedge (B)$ criteria interact, and whether an effective exchange antisymmetry emerges from the mixed-triangle structure.

OP4 (Relativistic extension — Dirac equation). The K_4 pulsation is a two-state object $(s^{(1)}, s^{(2)} = -s^{(1)})$, suggesting a natural spinor representation. The passage from (16) to the Dirac equation would require restoring the full $SL(2, \mathbb{C})$ structure of the Compton pulsation.

OP5 (Born rule from Gleason-type uniqueness). The complex amplitudes ψ_n are introduced without deriving the Born rule $|\psi_n|^2$. Document [9] initiates a Gleason-type approach in the PDL setting; completing it would yield a fully internal derivation of the probability interpretation.

OP6 (Connection between α_{PDL} and $G_{\mathrm{eff}}(N) = \sigma(N) \cdot G_{\mathrm{PDL}}$). The link between the Schrödinger coupling α_{PDL} and the gravitational coupling of D31 at the level of the 18-step chain complex ε_G^{18} remains to be established algebraically (D31, OP1).

Conclusion

This paper has derived the non-relativistic Schrödinger equation for an electron in the Coulomb field of a PDL proton from the $(A) \wedge (B)$ coupling criterion of D29, the S_4 transitivity theorem of D31, and the Compton pulsation of K_4 — without free parameter.

The central advance over the prior sketch documents is the derivation of the transition coefficient b. In those documents, b was a phenomenological parameter constrained but not determined by the PDL axioms. Here, condition (B) — $P_2 = -P_1$, verified exhaustively over 768 cases with zero counter-example — forces a π phase shift between the two half-cycles of every stable mixed triangle. The coherent sum over one pulsation cycle (equation (8)) gives a constructive amplitude $\Delta = 2P_1$, and the Compton action $\hbar\omega_c = m_e c^2$ then fixes b uniquely as the purely imaginary quantity (7). The S_4 transitivity of D31 simultaneously forces the nearest-neighbour symmetric structure of admissible transitions (Proposition 1) and, through the $(A) \wedge (B)$-filtered count $N_{\mathrm{mix}}(r) \propto R_{\mathrm{surf}}/r^2$ (Proposition 3), the Coulomb form of the potential with coupling $\alpha_{\mathrm{PDL}} = \mu/R_{\mathrm{surf}}^2$ from D29–D30.

The central theorem assembles these four propositions into equation (16), resolving open problem OP6 of D31. Numerical checks confirm: 0/768 counter-examples to the $(A) \wedge (B)$-stability equivalence; orbit matrix of S_4 uniformly equal to 4; discrete tridiagonal Hamiltonian converging to the Bohr levels at the 5×10^{-4} level.

References

[1] C. Laubscher. Axiomatic Derivation of the Leading Valence Engagement Amplitude in the PDL Framework. *PDL Corpus*, 2026. D29.

[2] C. Laubscher. Axiomatic Proof of the Coefficient 2 and the Isospin Identification in the PDL Framework. *PDL Corpus*, 2026. D30.

[3] C. Laubscher. Combinatorial Proton Architecture in the PDL Framework. *Zenodo*, 2026. D24-v2.

[4] C. Laubscher. Compatibility of the Schrödinger Framework with the Projective Dynamic Logo (PDL). *PDL Corpus*, 2026. v2.

[5] C. Laubscher. From Discrete Coherence Flux to Effective Fields: A Structured Programme within the PDL Framework. *PDL Corpus*, 2026.

[6] C. Laubscher. Introduction to the Projective Dynamic Logo Framework. *PDL Corpus*, 2026. D2.

[7] C. Laubscher. Projective Dynamic Logo: Main Framework. *PDL Corpus*, 2026. D1.

[8] C. Laubscher. Proof of the Density-Dependent Gravitational Coupling $G_{\mathrm{eff}}(N) = \sigma(N) \cdot G_{\mathrm{PDL}}$: Symmetry, Obstruction, and Linearity of the Densified Bridge Map. *PDL Corpus*, 2026. D31.

[9] C. Laubscher. Toward a Gleason-Type Uniqueness Theorem for Born's Rule in the PDL Framework. *PDL Corpus*, 2026.

[10] C. Laubscher. Towards a Schrödinger-Type Dynamics from the Projective Dynamic Logo (PDL): A Relational Sketch Linking Coherence Cycles, Active Surfaces, and Effective Potentials. *PDL Corpus*, 2026. v2.