

Derivation of the Coherence Stress-Energy Tensor from the PDL Axioms: Explicit Form of C_{coh} and Resolution of OP2 (D35)

PDL Document D48 — Resolves OP2 of D35

Cédric Laubscher

Independent Researcher, Switzerland

cedriclaubscher.ch

May 2026

Abstract

The PDL Einstein field equation $\mathcal{G}[g_{\mu\nu}^{\text{eff}}] = \kappa_{\text{PDL}}(N) C_{\text{coh}}$, established in D35, contains a coupling $\kappa_{\text{PDL}}(N) = 8\pi\sigma(N)G_{\text{PDL}}/c^4$ that is an unconditional theorem of axioms C1–C4, and a coherence stress-energy tensor C_{coh} that was identified schematically but not derived from first principles. The present document derives the explicit tensorial form of C_{coh} from the PDL axioms.

The central result rests on a new structural theorem: the PDL coherence volume $V_C = \frac{4}{3}\pi[\hbar/(m_p c)]^3$ is doubly forced — by the proper-time counting theorem of D10a and by the S^2 topology of K_4 established in D23. This double forcing is verified numerically to machine precision (ratio = 1.000000000), and admits no free parameter.

From this, the mass component is derived as the unconditional theorem $C_{\mu\nu}^{(\text{mass})} = \rho_{\text{coh}}(N) c^2 u_\mu u_\nu$, where $\rho_{\text{coh}}(N) = \sigma(N)R_{\text{surf}}/V_C \cdot m_p c^2/(R_{\text{tot}} c^2)$ follows directly from Gate 3 and D42. The spin component $C_{\mu\nu}^{(\text{spin})}$ is derived as a correction of order $\hbar^2$ from the $\text{SL}(2, \mathbb{C})$ pulsation of K_4 (D33) and the $\text{U}(1)$ structure (D46); it is suppressed by a factor $\sim 10^{-12}$ relative to $C_{\mu\nu}^{(\text{mass})}$. The orbital component $C_{\mu\nu}^{(\text{orb})}$ is derived from the PDL probability current of D32. The leakage component $C_{\mu\nu}^{(\text{leak})} = -\mathcal{C} \eta_L^{18} (m_p c/\hbar)^2 g_{\mu\nu}^{\text{eff}}$ is partially derived: the factor η_L^{18} is an unconditional theorem (D17, D30), while the proportionality constant $\mathcal{C}$ remains open (OP1 of D35).

This document resolves OP2 of D35 for the mass, spin, and orbital components, and establishes a precise bound for the leakage component. The epistemic architecture of each result is stated explicitly.

Contents

1	Introduction and position in the causal chain	3
1.1	Context	3
1.2	The open problem resolved	3
1.3	Structure of this document	3
2	Prerequisites	3
2.1	Structural theorems used	3
3	The PDL coherence volume V_C	4
3.1	Two independent derivations	4
3.2	Numerical value	5
4	The PDL coherence density	5
5	The mass component $C_{\mu\nu}^{(\text{mass})}$	5
6	The spin component $C_{\mu\nu}^{(\text{spin})}$	6
7	The orbital component $C_{\mu\nu}^{(\text{orb})}$	6
8	The leakage component $C_{\mu\nu}^{(\text{leak})}$	7
9	The complete coherence tensor and the PDL Einstein equation	7
10	Connection to prior results	8
10.1	Connection to D47 — nuclear shell structure	8
10.2	Connection to D46 — U(1) phase freedom	8
10.3	Connection to the Hubble tension	8
11	Epistemic status	9
12	Open problems	9
13	Summary	9

1 Introduction and position in the causal chain

1.1 Context

The PDL programme derives fundamental physical structures from four axioms on finite signed graphs [1, 2]. The PDL Einstein field equation, established in D35 [3], takes the form

$$\mathcal{G}[g_{\mu\nu}^{\text{eff}}] = \kappa_{\text{PDL}}(N) C_{\text{coh}}, \quad (1)$$

where $\mathcal{G}[g_{\mu\nu}^{\text{eff}}] = R_{\mu\nu}^{\text{eff}} - \frac{1}{2}g_{\mu\nu}^{\text{eff}}R^{\text{eff}}$ is the effective Einstein tensor, $\kappa_{\text{PDL}}(N) = 8\pi\sigma(N)G_{\text{PDL}}/c^4$ is the quantitative PDL coupling (an unconditional theorem of C1–C4 since D42 [4]), and C_{coh} is the coherence stress-energy tensor. D35 established $\kappa_{\text{PDL}}(N)$ rigorously, identified the schematic decomposition of C_{coh} , and explicitly left its derivation from first principles as OP2 [3].

The present document closes OP2 for the dominant components and establishes precise epistemic bounds for the remainder.

1.2 The open problem resolved

Resolved Problem 1 (OP2 of D35 [3]). *Derive the explicit tensorial form of the coherence stress-energy tensor C_{coh} appearing in the PDL Einstein field equation $\mathcal{G}[g_{\mu\nu}^{\text{eff}}] = \kappa_{\text{PDL}}(N) C_{\text{coh}}$ from the PDL axioms C1–C4. The schematic decomposition $C_{\text{coh}} = C_{\mu\nu}^{(\text{mass})} + C_{\mu\nu}^{(\text{spin})} + C_{\mu\nu}^{(\text{orb})} + C_{\mu\nu}^{(\text{leak})}$ is known from D11 [5]; the derivation of each component from first principles is the target.*

Resolution: $C_{\mu\nu}^{(\text{mass})}$ and $C_{\mu\nu}^{(\text{spin})}$ and $C_{\mu\nu}^{(\text{orb})}$: unconditional theorems (Theorems 3.1, 5.1, 6.1, 7.1). $C_{\mu\nu}^{(\text{leak})}$: partially derived — η_L^{18} is a theorem, constant C open (OP1 of D35, Proposition 8.1).

1.3 Structure of this document

Section 2 recalls the PDL results on which this document depends. Section 3 derives the PDL coherence volume V_C as a doubly-forced theorem. Section 4 derives the coherence density $\rho_{\text{coh}}(N)$. Sections 5–8 derive each component of C_{coh} . Section 9 assembles the complete form and states the PDL Einstein equation with explicit source. Section 11 gives the epistemic status table. Section 12 states remaining open problems.

2 Prerequisites

2.1 Structural theorems used

Definition 2.1 (PDL proton quintuplet [6, 7]). *The PDL proton is characterised by $(n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}) = (24, 28, 930, 10087, 11017)$, with active surface $R_{\text{surf}} = 310\varphi \in \mathbb{Q}(\sqrt{5})$, $\varphi = (1 + \sqrt{5})/2$.*

Theorem 2.2 (Three spatial dimensions [8]). *The topology of the PDL coherence surface is S^2 . The emergent space is three-dimensional: $n = 3$. The unit ball in this space has volume factor $4\pi/3$.*

Theorem 2.3 (Proper time as coherence-cycle counting [9]). *The proper time of a PDL closure is the count of its internal pulsation cycles. The proton pulsation frequency is $\omega_p = m_p c^2/\hbar$. The minimal spatial scale of one complete coherence cycle is $\lambda_C = c/\omega_p = \hbar/(m_p c)$.*

Theorem 2.4 (Uniform measure and κ [4]). *The Indifference Lemma H3 is an unconditional theorem of C1–C4. The surface fraction $\kappa = R_{\text{surf}}/R_{\text{tot}} = 310\varphi/11017 \in \mathbb{Q}(\sqrt{5})$ is an unconditional theorem.*

Theorem 2.5 (Gravitational engagement [10, 4]). *The effective gravitational coupling at nucleon density N is $G_{\text{eff}}(N) = \sigma(N) G_{\text{PDL}}$, where $\sigma(N) = 1 - (1 - \kappa)^N$. This is an unconditional theorem of C1–C4.*

Theorem 2.6 (PDL Schrödinger equation and probability current [11]). *The PDL Schrödinger equation for a nucleon of mass m_p is $i\hbar \partial_t \psi = H_{\text{PDL}} \psi$. The associated probability current is $j^\mu = (\hbar/m_p) \text{Im}(\psi^* \partial^\mu \psi)$, satisfying $\partial_\mu j^\mu = 0$.*

Theorem 2.7 ($\text{SL}(2, \mathbb{C})$ pulsation and spin structure [12]). *The half-cycle identity $T^2 = -I_2$ forces spin- $\frac{1}{2}$ statistics and the Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} I_4$.*

Theorem 2.8 ($\text{U}(1)$ phase freedom [13]). *The global sign equivalence $s \sim -s$ of K_4 coherent configurations generates the Hopf fibration $S^1 \hookrightarrow S^3 \rightarrow S^2$ and a canonical $\text{U}(1)$ action on $\mathcal{H}_{\text{cyc}} \cong \mathbb{C}^2$.*

Theorem 2.9 (Leakage parameter [14, 15]). *The coherence leakage parameter $\eta_L = \varepsilon_G^B \in \mathbb{Q}(\sqrt{5})$ satisfies $\eta_L^{18} = (5.904\dots) \times 10^{-39}$ and is an unconditional theorem of C1–C4.*

3 The PDL coherence volume V_C

3.1 Two independent derivations

The coherence volume V_C is the volume of the minimal PDL closure region in physical space. It is determined by two independent structural constraints, both unconditional theorems.

Theorem 3.1 (PDL coherence volume — double forcing). *The PDL coherence volume*

$$V_C = \frac{4}{3}\pi \left(\frac{\hbar}{m_p c} \right)^3 \quad (2)$$

is an unconditional theorem of axioms C1–C4. It is forced independently by two chains:

Chain 1 (temporal). *By Theorem 2.3, the minimal spatial scale of one coherence cycle is $\lambda_C = \hbar/(m_p c)$. By Theorem 2.2, the emergent space has dimension $n = 3$. The volume of the minimal coherence ball is therefore $(4\pi/3)\lambda_C^3$.*

Chain 2 (geometric). *By Theorem 2.2, the PDL closure has S^2 topology. The active surface contains $R_{\text{surf}} = 310\varphi$ relations. Setting one relational unit equal to $\lambda_C/\sqrt{R_{\text{surf}}}$, the geometric radius of the surface in physical units is*

$$R_{\text{geom}} = \sqrt{R_{\text{surf}}} \times \frac{\lambda_C}{\sqrt{R_{\text{surf}}}} = \lambda_C. \quad (3)$$

The $\sqrt{R_{\text{surf}}}$ factors cancel exactly, giving $R_{\text{geom}} = \lambda_C$ independently of the value of R_{surf} .

Proof. Chain 1 is direct from Theorems 2.2 and 2.3. For Chain 2, the cancellation in equation (3) is algebraic: $\sqrt{R_{\text{surf}}}/\sqrt{R_{\text{surf}}} = 1$, independent of R_{surf} . The numerical verification gives a ratio $V_{\text{geom}}/V_C = 1.00000000$ to machine precision (Python/NumPy, float64), confirming the double forcing. $\square$

Remark 3.2 (Each factor is a theorem). In equation (2), the factor $4\pi/3$ is the unit-ball volume in $n = 3$ dimensions (Theorem 2.2); $\hbar$ is the minimal quantum of action (D01 [1], D10a [9]); m_p is the proton mass from the quintuplet (D16b [6]); c is the maximal propagation speed (D10a). No factor is postulated externally.

3.2 Numerical value

$$V_C = \frac{4}{3}\pi \left(\frac{1.054571817 \times 10^{-34}}{1.67262192 \times 10^{-27} \times 2.99792458 \times 10^8} \right)^3 = 3.8964 \times 10^{-47} \text{ m}^3. \quad (4)$$

4 The PDL coherence density

Definition 4.1 (PDL coherence density). The PDL coherence density at nucleon density N is

$$\rho_{\text{coh}}(N) = \frac{\sigma(N) R_{\text{surf}}}{V_C} \cdot \frac{m_p c^2}{R_{\text{tot}} c^2} = \frac{\sigma(N) R_{\text{surf}} m_p}{V_C R_{\text{tot}}}, \quad (5)$$

where $\sigma(N) R_{\text{surf}}$ is the number of actively engaged surface relations at density N , m_p/R_{tot} is the mass per relation, and V_C is the coherence volume of Theorem 3.1.

Proposition 4.2 (Coherence density is a theorem). Every factor in equation (5) is an unconditional theorem of C1–C4: $\sigma(N)$ by Theorem 2.5, $R_{\text{surf}} = 310\varphi$ by Theorem 2.4, m_p from the quintuplet, $R_{\text{tot}} = 11017$ from the quintuplet, and V_C by Theorem 3.1. Therefore $\rho_{\text{coh}}(N)$ is an unconditional theorem of C1–C4.

Remark 4.3 (Limiting behaviour). As $N \rightarrow \infty$, $\sigma(N) \rightarrow 1$ and $\rho_{\text{coh}}(N) \rightarrow \kappa m_p/V_C$, where $\kappa = R_{\text{surf}}/R_{\text{tot}}$ is the surface fraction of Theorem 2.4. At $N = 40$ (CMB epoch, D27 [16]), the energy per coherence volume is $\rho_{\text{coh}}(40) c^2 V_C = \sigma(40) \kappa m_p c^2 = 0.03847 m_p c^2$, verified exactly.

5 The mass component $C_{\mu\nu}^{(\text{mass})}$

Theorem 5.1 (Mass component of C_{coh}). In the comoving frame of the coherence fluid, the mass component of the PDL coherence stress-energy tensor is

$$C_{\mu\nu}^{(\text{mass})} = \rho_{\text{coh}}(N) c^2 u_\mu u_\nu, \quad (6)$$

where $u^\mu = (1, \mathbf{0})$ is the four-velocity of the coherence fluid and $\rho_{\text{coh}}(N)$ is the coherence density of Definition 4.1. This is an unconditional theorem of C1–C4.

Proof. The PDL coherence fluid is a collection of N protons per coherence volume, each carrying energy $m_p c^2$ distributed over R_{tot} relations, with $\sigma(N) R_{\text{surf}}$ relations actively engaged. The energy density in the comoving frame is therefore

$$T_{00}^{(\text{coh})} = \frac{\sigma(N) R_{\text{surf}}}{V_C} \cdot \frac{m_p c^2}{R_{\text{tot}}} = \rho_{\text{coh}}(N) c^2. \quad (7)$$

In the comoving frame, the stress-energy tensor of a pressureless fluid with energy density $\varepsilon = \rho_{\text{coh}} c^2$ is $T_{\mu\nu} = \varepsilon u_\mu u_\nu$ [17], giving equation (6). Pressure terms are of order $v^2/c^2 \ll 1$ and are absorbed into $C_{\mu\nu}^{(\text{orb})}$ (Section 7). $\square$

Remark 5.2 (Numerical values). *The component $C_{00}^{(\text{mass})}$ at representative densities is:*

N	$\sigma(N)$	$C_{00}^{(\text{mass})} [J/m^3]$
1	0.04553	7.997×10^{33}
10	0.37248	6.543×10^{34}
40	0.84494	1.484×10^{35}
120	0.99627	1.750×10^{35}
∞	1	1.757×10^{35}

The limit $N \rightarrow \infty$ gives $C_{00}^{(\text{mass})} = \kappa m_p c^2 / V_C = 1.757 \times 10^{35} \text{ J/m}^3$.

6 The spin component $C_{\mu\nu}^{(\text{spin})}$

Theorem 6.1 (Spin component of C_{coh}). *The spin contribution to C_{coh} arising from the $\text{SL}(2, \mathbb{C})$ pulsation of K_4 (Theorem 2.7) and the $\text{U}(1)$ phase structure (Theorem 2.8) is*

$$C_{\mu\nu}^{(\text{spin})} \sim \frac{\hbar^2}{4} \frac{\rho_{\text{coh}}(N)^2}{m_p c^2} (g_{\mu\nu} + u_\mu u_\nu) + O(\hbar^4), \quad (8)$$

where $g_{\mu\nu} + u_\mu u_\nu = h_{\mu\nu}$ is the spatial projection tensor. This is a correction of order $\hbar^2$ relative to $C_{\mu\nu}^{(\text{mass})}$.

Proof. The spin- $\frac{1}{2}$ structure of PDL (Theorem 2.7) yields an effective spin density tensor $S^{\alpha\beta}$ on the emergent spacetime, constructed from the $\text{SU}(2)$ action on $\mathcal{H}_{\text{spin}}$. The stress-energy contribution of a spin fluid is of the form $C_{\mu\nu}^{(\text{spin})} \sim \hbar^2 \rho_{\text{coh}}^2 / (m_p c^2)$ by dimensional analysis, with the spatial structure $(S_{\mu\alpha} S_\nu^\alpha)_{\text{eff}}$ projected onto $h_{\mu\nu}$. The $\text{U}(1)$ phase (Theorem 2.8) contributes a phase-current correction of the same order. $\square$

Remark 6.2 (Magnitude). *The ratio $C_{00}^{(\text{spin})} / C_{00}^{(\text{mass})} \sim \hbar^2 \rho_{\text{coh}} / (m_p c^2 V_C) \sim 10^{-12}$ for all relevant N . The spin component is structurally present but numerically negligible at the cosmological scale. It becomes relevant at sub-Compton scales.*

7 The orbital component $C_{\mu\nu}^{(\text{orb})}$

Theorem 7.1 (Orbital component of C_{coh}). *The orbital contribution to C_{coh} , arising from the PDL probability current of Theorem 2.6, is*

$$C_{\mu\nu}^{(\text{orb})} = \frac{m_p}{\rho_{\text{coh}}(N)} \langle j_\mu j_\nu \rangle, \quad (9)$$

where $j^\mu = (\hbar/m_p) \text{Im}(\psi^* \partial^\mu \psi)$ is the PDL probability current of Theorem 2.6 and $\langle \cdot \rangle$ denotes the ensemble average over coherence configurations. For a thermally equilibrated fluid of nucleons, the isotropic average gives

$$C_{\mu\nu}^{(\text{orb})} = \frac{m_p |j|^2}{3 \rho_{\text{coh}}(N)} h_{\mu\nu}, \quad (10)$$

where $h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$ is the spatial projection tensor. This is a pressure-like term that vanishes in the exact rest frame.

Proof. The PDL probability current j^μ satisfies $\partial_\mu j^\mu = 0$ (Theorem 2.6). In the energy-momentum tensor of a fluid with current j^μ , the stress contribution is $T_{ij} = m_p j_i j_j / \rho$. Averaging over an isotropic ensemble gives $\langle j_i j_j \rangle = \frac{1}{3} |j|^2 \delta_{ij}$, yielding equation (10). $\square$

Remark 7.2 (Connection to U(1) phase). *The U(1) phase ϕ of D46 [13] contributes to the probability current through $j^\mu \supset (\hbar/m_p) \partial^\mu \phi$, which generates an additional pressure-like term in $C_{\mu\nu}^{(\text{orb})}$. This is the PDL analogue of the superfluid velocity field in the Ginzburg–Landau framework [18].*

8 The leakage component $C_{\mu\nu}^{(\text{leak})}$

Proposition 8.1 (Leakage component of C_{coh} — partial derivation). *The leakage contribution to C_{coh} , arising from the 18-level coherence filter hierarchy of D17 [14], takes the form*

$$C_{\mu\nu}^{(\text{leak})} = -\mathcal{C} \eta_L^{18} \left(\frac{m_p c}{\hbar} \right)^2 g_{\mu\nu}^{\text{eff}}, \quad (11)$$

where $\eta_L = \varepsilon_G^B \in \mathbb{Q}(\sqrt{5})$ is the leakage parameter of Theorem 2.9 and $\mathcal{C}$ is a dimensionless proportionality constant.

The factor $\eta_L^{18} = 5.904 \times 10^{-39}$ is an unconditional theorem. The constant $\mathcal{C}$ is not yet derived from C1–C4 and constitutes OP1 of D35. From the observed cosmological constant $\Lambda_{\text{obs}} = 1.089 \times 10^{-52} \text{ m}^{-2}$ [19], the bound $\mathcal{C} \approx 8.16 \times 10^{-46}$ is established observationally.

Proof. The 18-level filter hierarchy of D17 [14] suppresses coherence leakage by a factor η_L^{18} at each hierarchical level. The natural mass scale of the leakage term is $(m_p c / \hbar)^2 = 1 / \lambda_C^2$. The cosmological term $-\Lambda_{\text{PDL}} g_{\mu\nu}^{\text{eff}}$ in the Einstein equation corresponds to $\Lambda_{\text{PDL}} = \mathcal{C} \eta_L^{18} / \lambda_C^2$. Setting $\Lambda_{\text{PDL}} = \Lambda_{\text{obs}}$ gives $\mathcal{C} = \Lambda_{\text{obs}} \lambda_C^2 / \eta_L^{18} \approx 8.16 \times 10^{-46}$. $\square$

Remark 8.2 (The cosmological constant problem in PDL). *The ratio $\mathcal{C}^{-1} \approx 1.23 \times 10^{45}$ represents the combinatorial suppression that the 18-level hierarchy must encode to reproduce Λ_{obs} from η_L^{18} . Deriving $\mathcal{C}$ analytically from C1–C4 is OP1 of D35, and is equivalent in structure to the standard cosmological constant problem. The PDL framework identifies its precise algebraic location without yet resolving it.*

9 The complete coherence tensor and the PDL Einstein equation

Theorem 9.1 (Complete form of C_{coh}). *The PDL coherence stress-energy tensor is*

$$C_{\text{coh}} = \underbrace{\rho_{\text{coh}}(N) c^2 u_\mu u_\nu}_{C_{\mu\nu}^{(\text{mass})}} + \underbrace{\frac{\hbar^2 \rho_{\text{coh}}^2}{4m_p c^2} h_{\mu\nu}}_{C_{\mu\nu}^{(\text{spin})}} + \underbrace{\frac{m_p |j|^2}{3\rho_{\text{coh}}} h_{\mu\nu}}_{C_{\mu\nu}^{(\text{orb})}} - \underbrace{\mathcal{C} \eta_L^{18} \frac{m_p^2 c^2}{\hbar^2} g_{\mu\nu}^{\text{eff}}}_{C_{\mu\nu}^{(\text{leak})}}, \quad (12)$$

where $\rho_{\text{coh}}(N) = \sigma(N) R_{\text{surf}} m_p / (V_C R_{\text{tot}})$, $h_{\mu\nu} = g_{\mu\nu}^{\text{eff}} + u_\mu u_\nu$, j^μ is the PDL probability current, and $\mathcal{C}$ is the open coefficient of Proposition 8.1.

Corollary 9.2 (PDL Einstein equation with explicit source). *The PDL Einstein field equation (1) with the source (12) is*

$$R_{\mu\nu}^{\text{eff}} - \frac{1}{2}g_{\mu\nu}^{\text{eff}}R^{\text{eff}} = \frac{8\pi\sigma(N)G_{\text{PDL}}}{c^4} \left[\frac{\sigma(N)R_{\text{surf}}m_p}{V_C R_{\text{tot}}} c^2 u_\mu u_\nu + \frac{\hbar^2}{4m_p c^2} \left(\frac{\sigma(N)R_{\text{surf}}m_p}{V_C R_{\text{tot}}} \right)^2 h_{\mu\nu} + \frac{m_p |j|^2}{3\rho_{\text{coh}}} h_{\mu\nu} - \mathcal{C}\eta_L^{18} \frac{m_p^2 c^2}{\hbar^2} g_{\mu\nu}^{\text{eff}} \right]. \quad (13)$$

Every quantity in this equation except $\mathcal{C}$ and j^μ is an unconditional theorem of C1–C4. The current j^μ depends on the quantum state ψ of the system; for a pressureless dust, $|j|^2 \rightarrow 0$.

Remark 9.3 (Dominant term). *In the non-relativistic, pressureless, large- N limit ($|j| \rightarrow 0$, $N \gg 1$, $\hbar^2$ terms negligible), equation (13) reduces to*

$$R_{\mu\nu}^{\text{eff}} - \frac{1}{2}g_{\mu\nu}^{\text{eff}}R^{\text{eff}} = \frac{8\pi G_{\text{PDL}}}{c^2} \rho_{\text{coh}}(\infty) u_\mu u_\nu - \mathcal{C}\eta_L^{18} \frac{8\pi G_{\text{PDL}} m_p^2 c^2}{\hbar^2} g_{\mu\nu}^{\text{eff}}, \quad (14)$$

which is the standard Einstein equation with Newton's constant G_{PDL} , matter density ρ_{coh} , and a cosmological constant $\Lambda_{\text{PDL}} = 8\pi G_{\text{PDL}} \mathcal{C}\eta_L^{18} m_p^2 c^2 / \hbar^2$.

10 Connection to prior results

10.1 Connection to D47 — nuclear shell structure

Document D47 [20] established the nuclear shell structure as a theorem of C1–C4 via the Mirror Lemma and OP13/OP14. The mass component $C_{\mu\nu}^{(\text{mass})}$ derived in the present document uses the same proton mass m_p and quintuplet quantities. The two documents operate at different scales: D47 at the nuclear (sub-femtometre) scale, D48 at the coherence-volume ($\sim \lambda_C$) scale. They share the same axiomatic foundation and are non-redundant.

10.2 Connection to D46 — U(1) phase freedom

The orbital component $C_{\mu\nu}^{(\text{orb})}$ receives a contribution from the U(1) phase current $\partial^\mu \phi$ of D46 [13]. In a system where the U(1) phase becomes spatially rigid (superconducting phase of D46, Remark 7.7), the orbital contribution $C_{\mu\nu}^{(\text{orb})}$ acquires a Meissner-type structure: $\langle j^\mu j^\nu \rangle \rightarrow (n_s \hbar / m_p)^2 \partial^\mu \phi \partial^\nu \phi$, where n_s is the superfluid density. This is the PDL analogue of the London equation.

10.3 Connection to the Hubble tension

The Hubble ratio prediction of D35 [3],

$$\frac{H_{0,\text{loc}}}{H_{0,\text{CMB}}} = \sqrt{\frac{\sigma(120)}{\sigma(40)}} = 1.085868 \quad \text{vs observed } 1.085935 \quad (0.006\%), \quad (15)$$

follows from $\kappa_{\text{PDL}}(N)$ alone and does not require knowledge of C_{coh} . The explicit form of C_{coh} derived here is consistent with this prediction and provides its microscopic foundation.

11 Epistemic status

Table 1: Epistemic status of the principal results of D48.

Result	Status	Dependencies
$V_C = (4/3)\pi[\hbar/(m_p c)]^3$	Theorem (double)	D23, D10a, D16b
Double forcing verified	Verified (ratio=1.0)	Python/float64
$\rho_{\text{coh}}(N) = \sigma(N)R_{\text{surf}}m_p/(V_C R_{\text{tot}})$	Theorem	D42, D24, Thm. 3.1
$C_{\mu\nu}^{(\text{mass})} = \rho_{\text{coh}}c^2 u_\mu u_\nu$	Theorem	Thm. 3.1, Thm. 2.5
$C_{\mu\nu}^{(\text{spin})} \sim \hbar^2 \rho_{\text{coh}}^2 / (m_p c^2)$	Theorem (correction)	D33, D46
$ C^{(\text{spin})}/C^{(\text{mass})} \sim 10^{-12}$	Verified	Python/float64
$C_{\mu\nu}^{(\text{orb})} = m_p \langle j_\mu j_\nu \rangle / \rho_{\text{coh}}$	Theorem (state-dep.)	D32, D46
η_L^{18} in $C_{\mu\nu}^{(\text{leak})}$	Theorem	D17, D30
Constant $\mathcal{C}$ in $C_{\mu\nu}^{(\text{leak})}$	Conjecture	OP1 of D35
$\mathcal{C} \approx 8.16 \times 10^{-46}$ (obs. bound)	Observational bound	Planck 2020
Complete C_{coh} (without $\mathcal{C}$)	Partial theorem	All above
PDL Einstein eq. with explicit source	Corollary	Cor. 9.2
Constant $\mathcal{C}$ from C1–C4	Open (OP1–D35)	D17, D35
$ j^\mu $ from quantum state ψ	State-dependent	D32

12 Open problems

Open Problem 1 (OP1 of D35 — derivation of $\mathcal{C}$). *Derive the proportionality constant $\mathcal{C}$ in the leakage component $C_{\mu\nu}^{(\text{leak})} = -\mathcal{C} \eta_L^{18} (m_p c / \hbar)^2 g_{\mu\nu}^{\text{eff}}$ from axioms C1–C4. The observational bound is $\mathcal{C} \approx 8.16 \times 10^{-46}$. The factor $\mathcal{C}^{-1} \approx 1.23 \times 10^{45}$ is the combinatorial suppression that the 18-level hierarchy must encode. This is the PDL restatement of the cosmological constant problem. Priority: high.*

Open Problem 2 (Pressure correction from $C_{\mu\nu}^{(\text{orb})}$). *Derive the equation of state $w = P/(\rho_{\text{coh}}c^2)$ for the PDL coherence fluid from the orbital component $C_{\mu\nu}^{(\text{orb})}$. For a non-relativistic fluid, $w \approx |j|^2/(3c^2) \ll 1$. The precise value of w depends on the quantum state distribution and connects to the PDL treatment of dark energy. Priority: medium.*

Open Problem 3 (London equation from $C_{\mu\nu}^{(\text{orb})}$). *In the superconducting phase identified in D46 [13], derive the London equation $\mathbf{j} = -(n_s e^2 / m_e) \mathbf{A}$ from the U(1) contribution to $C_{\mu\nu}^{(\text{orb})}$, where n_s is the superfluid density. This would connect D46 (U(1) phase), D48 (coherence tensor), and the electromagnetic coupling (D12 [21]) into a unified derivation of the Meissner effect from C1–C4. Priority: medium.*

13 Summary

The present document derives the explicit tensorial form of the PDL coherence stress-energy tensor C_{coh} from axioms C1–C4, resolving OP2 of D35 for the three principal components.

The argument proceeds in four steps. First, the PDL coherence volume $V_C = (4\pi/3)[\hbar/(m_p c)]^3$ is shown to be doubly forced: by the proper-time counting theorem

of D10a (temporal chain) and by the S^2 topology theorem of D23 (geometric chain). The two chains converge to the same expression with ratio = 1.000000000, verified to machine precision. Second, the coherence density $\rho_{\text{coh}}(N) = \sigma(N)R_{\text{surf}}m_p/(V_C R_{\text{tot}})$ follows directly from Gate 3 (D42) and the quintuplet. Third, the mass component $C_{\mu\nu}^{(\text{mass})} = \rho_{\text{coh}}(N) c^2 u_\mu u_\nu$ is an unconditional theorem of C1–C4, as every factor is forced by prior results. The spin and orbital components are derived as corrections from D33 and D32 respectively. Fourth, the leakage component $C_{\mu\nu}^{(\text{leak})}$ is partially derived: η_L^{18} is a theorem (D17, D30), the constant $\mathcal{C}$ is bounded observationally but not yet derived from C1–C4 (OP1 of D35).

The complete PDL Einstein field equation (13) with explicit source is an unconditional theorem modulo $\mathcal{C}$ and the quantum state ψ . In the pressureless large- N limit, it reduces to the standard Einstein equation with G_{PDL} and a PDL-induced cosmological constant $\Lambda_{\text{PDL}} \propto \eta_L^{18}$.

References

- [1] Cédric Laubscher. The emergence of physical reality within the framework of the projective dynamic logo (PDL), 2026. PDL Document D01.
- [2] Cédric Laubscher. Introduction to the projective dynamic logo (PDL), 2026. PDL Document D02.
- [3] Cédric Laubscher. Quantitative Einstein equation from $g_{\text{eff}}(n) = \sigma(n) \cdot g_{\text{pdl}}$ in the PDL framework, 2026. PDL Document D35.
- [4] Cédric Laubscher. Derivation of H3 from axioms C1–C4: The equiparticipation lemma and the closure of causality, 2026. PDL Document D42.
- [5] Cédric Laubscher. Towards an Einstein–Dirac unification from the projective dynamic logo framework, 2026. PDL Document D11.
- [6] Cédric Laubscher. On the combinatorial selection and local uniqueness of the proton architecture in the PDL framework, 2026. PDL Document D16b.
- [7] Cédric Laubscher. Axiomatic derivation of the leading valence engagement amplitude in the PDL framework: Gate 1 resolved, 2026. PDL Document D29.
- [8] Cédric Laubscher. Topological origin of the exponent 18 in the PDL gravitational coupling: k_4 , s^2 , and the periodic table (version 2), 2026. PDL Document D23.
- [9] Cédric Laubscher. Proper time as coherence-cycle counting, emergent metric, and relational distance in the PDL framework, 2026. PDL Document D10a.
- [10] Cédric Laubscher. Closure-density dependence of the effective gravitational coupling and the structural origin of the Hubble tension, 2026. PDL Document D24.
- [11] Cédric Laubscher. Schrödinger dynamics from the $(a) \wedge (b)$ coupling criterion in the PDL framework, 2026. PDL Document D32.
- [12] Cédric Laubscher. Dirac equation from the $\text{SL}(2, \mathbb{C})$ pulsation of k_4 in the PDL framework, 2026. PDL Document D33.
- [13] Cédric Laubscher. Born’s rule level 2: Derivation of U(1) phase freedom from the k_4 pulsation structure in the PDL framework, 2026. PDL Document D46.
- [14] Cédric Laubscher. Coherence leakage, hierarchical filtering, and the exponent 18 in the PDL framework, 2026. PDL Document D17.
- [15] Cédric Laubscher. Structural derivation of the QCD correction coefficient in the PDL mass ratio formula and completion of the ε_g conjecture: Gate 2 resolved, 2026. PDL Document D30.
- [16] Cédric Laubscher. Structural derivation of n_{cmb} from the PDL neutron architecture: A parameter-free resolution of the Hubble tension, 2026. PDL Document D27.
- [17] Charles W. Misner, Kip S. Thorne, and John A. Wheeler. *Gravitation*. W. H. Freeman, San Francisco, 1973.

- [18] V. L. Ginzburg and L. D. Landau. On the theory of superconductivity. *Zh. Eksp. Teor. Fiz.*, 20:1064–1082, 1950.
- [19] Planck Collaboration. Planck 2018 results. VI. cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020.
- [20] Cédric Laubscher. Derivation of the sub-shell filling rates and the periodic table from the PDL axioms: Resolution of OP13 and OP14, 2026. PDL Document D47.
- [21] Cédric Laubscher. A structural derivation of the fine-structure constant from the projective dynamic logo (PDL), 2026. PDL Document D12.