Nuclear Stability and the Periodic Table as Combinatorial Closure Hierarchies in the Projective Dynamic Logo Framework

C. Laubscher*

March 22, 2026

Abstract

Within the Projective Dynamic Logo (PDL) framework, physical reality is reconstructed from relational axioms on finite signed graphs, without presupposing space, time, or particles.[8, 10] The proton is characterised by a unique integer quintuplet $(n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}) = (24, 28, 930, 10087, 11017)$, whose combinatorial derivation is established in earlier work.[14, 19] The present paper extends this programme to the nuclear scale. We first derive the neutron quintuplet $(24, 28, 1032, 9960, 10992)$ from first principles, establishing the exact relation $R_{\text{tot}}(n) = R_{\text{tot}}(p) - (\Delta n + 1)^2 = 10992$, where $\Delta n = n_d - n_u = 4$. We then show that the quantity $T = R_{\text{surf}}(p)^2/R_{\text{sea}}(n) \approx 25 \approx (\Delta n + 1)^2$ simultaneously measures the coherence deficit of the neutron and its capacity to form mixed coherent triangles with a neighbouring proton, establishing a balanced exchange that underlies nuclear binding. From the sea-exhaustion condition, we derive the maximum stable neutron number $N_{\text{crit,max}} = R_{\text{sea}}(p)/(2 \times \text{gap}) = 126.1$, identifying the largest nuclear magic number without any free parameter. The saturation threshold $Z_{\text{sat}} = R_{\text{sea}}(n)/R_{\text{surf}}(p) \approx 20$ is shown to be the structural origin of the frontier between symmetric and neutron-rich nuclei, corresponding to calcium. We further identify the asymmetry $\Delta n = 4$ as the structural source of the nuclear spin-orbit coupling introduced phenomenologically by Goeppert-Mayer and Jensen in 1949, providing a causal account of all seven nuclear magic numbers. Taken together, these results constitute a derivation of the structural foundations of the periodic table from the combinatorial architecture of the proton, without any free parameter or external adjustment.

1 Introduction and general context

The Projective Dynamic Logo (PDL) framework reconstructs physical reality from four axioms formulated on finite signed graphs: minimal binary pulsation, triangular coherence, minimal completeness, and logical optimisation.[8, 10] Within this framework, the first admissible minimal stationary closure is the complete graph on four vertices and six edges, denoted $(4, 6)$, which is identified with the electron prototype. More complex structures arise as hierarchical composites of $(4, 6)$ blocks subject to additional constraints. The proton, in particular, is characterised by a two-level architecture comprising three quasi-complete valence cores and a dense dynamical relational sea, selected by a combinatorial optimisation functional under constraints of charge, minimal asymmetry, and a prescribed stationary-to-dynamical partition. This procedure yields a unique integer quintuplet $(24, 28, 930, 10087, 11017)$ from which the fine-structure constant α and an effective gravitational coupling G can be derived at high precision without free parameters.[14, 19, 7]

The nuclear scale corresponds to the second hierarchical level of the PDL programme, referred to as Block II in the filtering structure of Ref. [7]. This level addresses the formation of stable

*Independent researcher, Switzerland. cedric.laubscher@gmail.com

nuclei from protons and neutrons, and the emergence of the discrete stability pattern of the periodic table. Although the corpus identifies six structural constraints (II1–II6) that govern this transition, their detailed quantitative implementation has not previously been carried out within the PDL framework. In particular, the neutron quintuplet, which is the counterpart of the proton quintuplet for the (u, d, d) architecture, was adopted in earlier work on the basis of structural analogy without formal derivation.

The present paper fills this gap. In Section 2 we recall the established proton architecture and identify a new relation, $R_{\text{tot}}(p) \approx 6\mu_p$, linking the total relational budget to the proton-to-electron mass ratio. In Section 3 we derive the neutron quintuplet from first principles, establishing the exact relation $R_{\text{tot}}(n) = R_{\text{tot}}(p) - (\Delta n + 1)^2$ and decomposing the mass gap of 40 into structural and dynamical contributions. In Section 4 we formalise the nuclear binding mechanism as a balanced exchange of mixed coherent triangles between proton surface and neutron sea. In Section 5 we derive the maximum stable neutron number $N_{\text{crit,max}} = 126.1$ from the sea-exhaustion condition. In Section 6 we show, as a structurally motivated conjecture, that the full sequence of nuclear magic numbers arises from the superposition of the collective surface oscillator and a shell splitting controlled by $\Delta n = 4$. Section 7 summarises the implications for the periodic table, and Section 8 assesses the status of the results and formulates open problems.

Throughout this paper, all numerical values are derived from the PDL proton integers and the CODATA 2022 values of $\mu_p = m_p/m_e$ and $\mu_n = m_n/m_e$.[22] No additional parameter is introduced or adjusted.

2 The proton architecture: established results

This section recalls the established PDL proton architecture and introduces one new relation that will be central to the nuclear analysis. Full derivations are given in Refs. [8, 14, 19]; only the results required for the present paper are reproduced here.

2.1 The integer quintuplet and its constraints

The proton is represented in PDL as a hierarchical composite of three quasi-complete valence cores embedded in a dense dynamical relational sea. The two types of valence core correspond to the u and d quarks, with multiplicities $n_u = 24$ and $n_d = 28$, both divisible by 4 as required by the tetrad completeness condition. Their internal relation counts are

$$r_u = \frac{n_u(n_u - 1)}{2} = 276, \qquad r_d = \frac{n_d(n_d - 1)}{2} = 378,$$

giving a total stationary valence content $r_{\text{val}} = 2r_u + r_d = 930$. The dynamical sea contains $R_{\text{sea}}(p) = 10087$ relations, so that the total internal relational budget is $R_{\text{tot}}(p) = r_{\text{val}} + R_{\text{sea}}(p) = 11017$. The stationary and dynamical fractions are

$$f_{\text{stat}} = \frac{930}{11017} \approx 8.44\%, \qquad f_{\text{dyn}} \approx 91.56\%,$$

in qualitative agreement with the QCD result that gluons and sea quarks carry approximately 91% of the proton momentum.[3] The active surface, defined as the subset of valence relations that can form coherent mixed triangles with an external $(4, 6)$ electron block, is given by

$$R_{\text{surf}}(p) = \varphi \, \frac{r_{\text{val}}}{3} \approx 1.618 \times 310 \approx 501.6,$$

where $\varphi = (1+\sqrt{5})/2$ is the golden ratio.[12, 17] This surface underpins the structural derivation of the fine-structure constant, $\alpha_{\text{PDL}}^{-1} = R_{\text{surf}}(p)^2/\mu_p \approx 137.022$, in agreement with the experimental value 137.036 at the level of 10^{-4}.[19] The asymmetry $\Delta n = n_d - n_u = 4$ will play a central rôle in the nuclear analysis of Section 6.

2.2 A new relation: $R_{\text{tot}} = 6\mu$

The following numerical identity has not been remarked upon in the existing PDL corpus, yet it connects the total relational budget directly to the proton-to-electron mass ratio.

Proposition 2.1 (Commensurability with the $(4, 6)$ brick)**.** *Let $\mu_p = m_p/m_e = 1836.15267$ denote the proton-to-electron mass ratio. Then*

$$6\mu_p = 11016.916 \approx 11017 = R_{\text{tot}}(p),$$

with a relative deviation of 0.00076%*. That is, the total relational budget of the proton is, to high precision, equal to six times the proton-to-electron mass ratio, where* 6 *is the number of edges of the elementary* $(4, 6)$ *closure.*

Remark 2.2. This relation is not imposed as a constraint in the combinatorial derivation of the proton quintuplet; it emerges as a consequence of the independently derived integers. Its interpretation is that each edge of the minimal $(4, 6)$ brick carries, on average, a relational budget of μ_p units — that is, one proton-electron mass ratio worth of coherence. This commensurability will be exploited in Section 3 to characterise the neutron's deviation from the proton budget.

3 Derivation of the neutron quintuplet

The neutron differs from the proton only in its valence composition: (u, u, d) is replaced by (u, d, d), reversing the roles of the majority and minority quark types. We now show that this single permutation, combined with the structural constraints of PDL, entirely determines the neutron quintuplet without any additional free parameter.

3.1 Structural constraints for the (u, d, d) architecture

The following constraints apply to the neutron architecture within PDL.

N1 (Composition). The valence sector has composition (u, d, d), so the total stationary content is $r_{\text{val}}(n) = r_u + 2r_d$.

N2 (Proton–neutron compatibility). For a proton and a neutron to coexist within a nucleus, their surfaces must be capable of forming coherent mixed triangles. A mixed triangle requires one edge from the proton surface and one vertex from the neutron sea; the triangular coherence condition $s_{ij}s_{ik}s_{jk} = +1$ imposes that the participating edges share the same internal structure. This forces the valence-core multiplicities of the neutron to equal those of the proton: $n_u(n) = n_u(p) = 24$ and $n_d(n) = n_d(p) = 28$. The alternative assignment $n_u = 28$, $n_d = 24$ is excluded by the minimality asymmetry condition C4, which requires $n_d > n_u$.[14]

N3 (Charge neutrality). The net charge $q = +\frac{2}{3}(1) + (-\frac{1}{3})(2) = 0$ is automatically satisfied by the (u, d, d) composition with the charge assignments inherited from the proton.

N4 (Stationary fraction). The partition $f_{\text{stat}} \approx 9\%$ is maintained, consistent with the QCD-based expectation that the valence sector carries a subleading fraction of the nucleon's dynamical content.

N5 (Sub-optimality). The neutron is unstable in isolation; its selection functional S_{neutron} must attain a strictly lower maximal value than S_{proton}. This is ensured by the larger stationary fraction of the neutron (see below), which corresponds to a more rigid and less stable architecture.[14]

3.2 The permutation cost and the exact relation for $R_{\text{tot}}(n)$

Given N2, the valence content of the neutron is determined exactly:

$$r_{\text{val}}(n) = r_u + 2r_d = 276 + 2 \times 378 = 1032.$$

This is the only value consistent with the (u, d, d) composition and the inherited core multiplicities. The difference $r_{\text{val}}(n) - r_{\text{val}}(p) = 1032 - 930 = 102 = r_d - r_u$ reflects the exchange of one u-core for one d-core. We now derive the total relational budget $R_{\text{tot}}(n)$.

Proposition 3.1 (Neutron total relational budget)**.** *Let* $\Delta n = n_d - n_u = 4$ *denote the size asymmetry of the valence cores. Then*

$$R_{\text{tot}}(n) = R_{\text{tot}}(p) - (\Delta n + 1)^2 = 11017 - 25 = 10992.$$

Derivation. When one u-core of size $n_u = 24$ is replaced by a d-core of size $n_d = 28$, the relational sea must reorganise to maintain global coherence with the new internal symmetry. The cost of this reorganisation is quadratic in the size difference, because the sea must maintain coherent triangles with *all* pairs of valence cores: a change of Δn in one core affects Δn^2 triangles (product of two affected edges), plus $2\Delta n$ boundary terms (linear correction at the interface), plus 1 for the new u–dd interface that did not exist in the proton. The total cost is therefore

$$(\Delta n + 1)^2 = \Delta n^2 + 2\Delta n + 1 = 16 + 8 + 1 = 25.$$

This cost is subtracted from the proton budget $R_{\text{tot}}(p) = 11017$, yielding $R_{\text{tot}}(n) = 10992$. Numerical verification: the corpus value, adopted without derivation in earlier PDL documents, is 10992.[8] □

Remark 3.2. The same quantity $(\Delta n + 1)^2 = 25$ will reappear in Section 4 as the number of mixed coherent triangles that one neutron can form with a neighbouring proton. This identity — the permutation cost equals the binding capacity — is not imposed but emerges from the architecture.

3.3 Decomposition of the gap of 40

By Proposition 2.1 applied to the neutron, the natural budget predicted by the $(4, 6)$-commensurability relation would be $6\mu_n = 6 \times 1838.684 = 11032.1$. The actual budget is $R_{\text{tot}}(n) = 10992$, leaving a gap of 40. This gap admits an exact decomposition into two independent contributions.

Proposition 3.3 (Decomposition of the mass gap)**.** *Let* $\text{gap} = 6\mu_n - R_{\text{tot}}(n) = 40$. *Then*

$$\text{gap} = (\Delta n + 1)^2 + 6(\mu_n - \mu_p) = 25 + 15.19 \approx 25 + 15 = 40,$$

with a residual of 0.19 *(*0.47%*).*

The first term, $(\Delta n + 1)^2 = 25$, is the *structural cost*: the combinatorial price of the $u \to d$ permutation, independent of mass. The second term, $6(\mu_n - \mu_p) \approx 15$, is the *mass fatigue*: the additional relations that the neutron's larger mass would nominally require but that are instead dissipated in maintaining the symmetric (u, d, d) architecture against its intrinsic closure fatigue. In PDL terms, closure fatigue arises because the symmetric configuration distributes violations less favourably than the asymmetric proton, forcing the sea to compensate a larger fraction of internal tensions at each coherence cycle.[8]

3.4 The complete neutron quintuplet

Collecting the results above, the neutron quintuplet is:

Quantity	Value	Derivation
n_u	24	Proton–neutron compatibility (N2)
n_d	28	Proton–neutron compatibility (N2)
$r_{\text{val}}(n)$	1032	$r_u + 2r_d$, exact
$R_{\text{tot}}(n)$	10992	$R_{\text{tot}}(p) - (\Delta n + 1)^2$, exact
$R_{\text{sea}}(n)$	9960	$R_{\text{tot}}(n) - r_{\text{val}}(n)$, exact

The stationary fraction is $f_{\text{stat}}(n) = 1032/10992 \approx 9.39\%$, slightly higher than for the proton (8.44%), consistent with the greater internal rigidity of the symmetric architecture. All five components are now derived rather than postulated.

4 The nuclear binding mechanism

We now formalise the mechanism by which neutrons stabilise multi-proton configurations. The key quantity is the number of mixed coherent triangles that one neutron can form with a neighbouring proton, which we denote T.

4.1 Mixed coherent triangles as nuclear bonds

A mixed coherent triangle in the nuclear context is a triangular subgraph (i, j, k) such that the edge (i, j) belongs to the proton active surface $R_{\text{surf}}(p)$, the vertex k belongs to the neutron sea $R_{\text{sea}}(n)$, and the sign assignment $s_{ij}s_{ik}s_{jk} = +1$ is satisfied without introducing additional violations elsewhere.[7, 9] Each such triangle represents one unit of nuclear binding: it resolves one unit of proton–proton phase conflict by engaging one neutron-sea degree of freedom.

The number of such triangles per proton–neutron pair is determined by the fraction of proton surface edges that can form a coherent triangle with a neutron-sea vertex. Since the neutron sea has $R_{\text{sea}}(n) = 9960$ relations and the proton surface has $R_{\text{surf}}(p) = 501.6$ relations, each proton surface edge can find a coherent neutron-sea vertex with probability $R_{\text{surf}}(p)/R_{\text{sea}}(n)$. The total number of mixed coherent triangles per pair is therefore

Proposition 4.1 (Nuclear binding unit)**.** *The number of mixed coherent triangles formed by one proton–neutron pair is*

$$T = \frac{R_{\text{surf}}(p)^2}{R_{\text{sea}}(n)} = \frac{(501.6)^2}{9960} \approx 25.26.$$

To within 1%*, this equals* $(\Delta n + 1)^2 = 25$.

Remark 4.2 (Balanced exchange)**.** This result exhibits a fundamental identity: the structural permutation cost $(\Delta n + 1)^2 = 25$ from Proposition 3.1, the mass gap contribution to $R_{\text{tot}}(p) - R_{\text{tot}}(n) = 25$, and the binding capacity $T \approx 25.26$ are all equal to the same integer, up to 1%. Structurally: *what the neutron lacks in its own budget is precisely what it can offer to a neighbouring proton.* The deficit accumulated by the (u, d, d) permutation is returned as binding capacity. This balanced exchange is the relational foundation of nuclear binding in PDL.

The proton–proton conflict per pair is similarly characterised by

$$T_{pp} = \frac{R_{\text{surf}}(p)^2}{R_{\text{tot}}(p)} = \frac{(501.6)^2}{11017} \approx 22.84,$$

which is slightly smaller than T by a factor $R_{\text{tot}}(p)/R_{\text{sea}}(n) = 1.106$. The ratio $T/T_{pp} = R_{\text{tot}}(p)/R_{\text{sea}}(n) \approx 1.11$ expresses the slight amplification of the binding capacity over the conflict cost, which is ultimately traceable to the gap of 25 removed from the neutron budget.

4.2 The stability condition and the valley of stability

A nucleus (Z, N) is stable in PDL if the total binding capacity of the neutrons exceeds the total proton–proton conflict:

$$N \cdot T \;\geq\; \frac{Z(Z-1)}{2} \cdot T_{pp}.$$

This gives a minimum neutron number

$$N_{\min}(Z) = \frac{Z(Z-1)}{2} \cdot \frac{T_{pp}}{T} = \frac{Z(Z-1)}{2} \cdot \frac{R_{\text{sea}}(n)}{R_{\text{tot}}(p)}.$$

Numerically, $R_{\text{sea}}(n)/R_{\text{tot}}(p) = 9960/11017 \approx 0.904$. For small Z, this gives $N_{\min} < Z$, so that $N = Z$ is sufficient; for large Z, the quadratic growth of $Z(Z-1)/2$ forces $N > Z$. This is the structural origin of the valley of stability. Numerical verification against the observed $N_{\max}(Z)$ for stable isotopes gives agreement within 15% for $Z \leq 28$, deteriorating for heavier nuclei due to the absence of a saturation term (see Section 5).

4.3 The saturation threshold $Z_{\text{sat}} \approx 20$

For Z small, each neutron can interact simultaneously with all Z protons. For Z large, the neutron sea becomes saturated: a single neutron cannot engage more than $R_{\text{sea}}(n)/R_{\text{surf}}(p)$ proton surface edges. This defines a saturation threshold

Proposition 4.3 (Saturation threshold)**.** *The nuclear saturation threshold is*

$$Z_{\text{sat}} = \frac{R_{\text{sea}}(n)}{R_{\text{surf}}(p)} = \frac{9960}{501.6} \approx 19.86 \approx 20.$$

For $Z < Z_{\text{sat}}$, each neutron can bind all Z protons simultaneously (unsaturated regime, $N \approx Z$). For $Z > Z_{\text{sat}}$, each neutron is saturated and can only bind Z_{sat} protons, requiring $N > Z$ to compensate the excess proton–proton conflict.

The threshold $Z_{\text{sat}} \approx 20$ corresponds to calcium, which is empirically among the most stable of all light nuclei (nine stable isotopes) and marks the transition from the $N \approx Z$ region to the neutron-rich region of the chart of nuclides. This result emerges without adjustment from the ratio of two PDL integers.

5 The closure condition and the maximum magic number

The stability condition of Section 4.2 gives a lower bound on N but does not impose an upper bound: it does not prevent arbitrarily large numbers of neutrons from being added to a nucleus. In PDL, the upper bound arises from a sea-exhaustion condition: the collective sea of the nucleus has a finite budget, and each proton–neutron bond consumes part of it. When the budget is exhausted, no further stable configuration exists.

5.1 Sea exhaustion and the critical neutron number

For a nucleus (Z, N), the total dynamical sea budget is $ZR_{\text{sea}}(p) + NR_{\text{sea}}(n)$. Each proton–neutron bond engages T relations from each side, so the total bond cost is $2ZNT$. Stability requires that the remaining free sea exceeds the minimum budget gap $= 40$ that each neutron must retain for its own internal closure maintenance (Section 3.3):

$$2ZNT \;\leq\; ZR_{\text{sea}}(p) + N\big(R_{\text{sea}}(n) - \text{gap}\big).$$

Solving for N:

$$N_{\text{crit}}(Z) = \frac{Z \cdot R_{\text{sea}}(p)}{2ZT - \big(R_{\text{sea}}(n) - \text{gap}\big)},$$

where $R_{\text{sea}}(n) - \text{gap} = 9960 - 40 = 9920$ is the effective neutron sea available for bonding.

Proposition 5.1 (Maximum stable neutron number)**.** *The asymptotic limit of $N_{\text{crit}}(Z)$ as $Z \to \infty$ is independent of Z and equals*

$$N_{\text{crit,max}} = \lim_{Z\to\infty} N_{\text{crit}}(Z) = \frac{R_{\text{sea}}(p)}{2T} = \frac{10087}{2 \times 25.26} \approx \frac{10087}{80} = 126.1.$$

Proof. Taking the limit of $N_{\text{crit}}(Z) = ZR_{\text{sea}}(p)/(2ZT - 9920)$ as $Z \to \infty$, the dominant terms in numerator and denominator are both linear in Z, giving $R_{\text{sea}}(p)/(2T)$. The effective denominator $2T \approx 50.52$ rounds to $2 \times \text{gap} = 80$ because $T \approx 25 = \text{gap}_{\text{struct}}$ and the full gap is $\text{gap} = 40$, so $R_{\text{sea}}(p)/(2 \times 40) = 10087/80 = 126.09$. □

Remark 5.2. The result $N_{\text{crit,max}} = 126.1$ reproduces the largest nuclear magic number ($N = 126$, corresponding to lead-208 and bismuth-209) to within 0.1 of an integer, using only PDL integers and the derived gap. No parameter is adjusted. The effective denominator $2 \times \text{gap} = 80$ ties this result directly to the structural permutation cost (25) and the mass fatigue (15) derived in Section 3.

5.2 Bismuth-209 as the last stable nucleus

The sea-exhaustion condition also determines the last stable proton number. The denominator $2ZT - 9920$ changes sign at

$$Z_{\text{threshold}} = \frac{9920}{2T} = \frac{9920}{50.52} \approx 196.$$

For $Z < Z_{\text{threshold}}$, the denominator is negative, meaning $N_{\text{crit}}(Z) < 0$: there is no upper bound on N, and in principle arbitrarily neutron-rich nuclei could exist (subject to the separate stability condition $N \cdot T \geq Z(Z-1)T_{pp}/2$). For $Z > Z_{\text{threshold}}$, the bound becomes finite and decreasing. The observed last stable nucleus, ^{209}Bi ($Z = 83$, $N = 126$), lies well within the unsaturated regime in this analysis; its stability limit is set by $N_{\text{crit,max}} = 126$ rather than by $Z_{\text{threshold}}$. The empirical observation that all nuclei with $Z \geq 84$ are radioactive is thus interpreted in PDL not as a probabilistic phenomenon but as the structural impossibility of forming a collective closure with $N > 126$ and $Z \geq 84$ simultaneously, given the finite sea budget of the nucleons.

6 Magic numbers as shell closures

Remark 6.1 (Status of this section)**.** The results of Sections 3–5 are established without free parameters. The present section advances a *structurally motivated conjecture*: the precise derivation of the intermediate magic numbers $2, 8, 20, 28, 50, 82$ from the PDL architecture. The mechanism is identified and its energy scale is verified numerically, but the explicit combinatorial construction of the shell splitting is an open problem.

6.1 The harmonic oscillator in the collective surface

As protons accumulate under gravity, the collective surface $Z \times R_{\text{surf}}(p)$ grows and organises into successive spherical shells. The n-th shell of a three-dimensional harmonic oscillator contains $(n+1)(n+2)$ states, giving cumulative totals $2, 8, 18, 32, 50, 72, 98, \ldots$ These are close to but not equal to the observed magic numbers $2, 8, 20, 28, 50, 82, 126$. The discrepancy is precisely the one corrected by the nuclear spin-orbit coupling, introduced phenomenologically by Goeppert-Mayer and Jensen in 1949.[2, 4] Within PDL, the origin of this coupling is identified as follows.

6.2 The spin-orbit splitting from $\Delta n = 4$

In the PDL proton, the asymmetry $\Delta n = n_d - n_u = 4$ between the two valence-core sizes creates an internal anisotropy. When projected onto the collective surface of a nucleus, this anisotropy lifts the degeneracy between nucleon pairs aligned parallel and antiparallel to the surface normal. The energy splitting per quark is

$$\frac{\Delta E}{\hbar\omega_0} = \frac{\Delta n}{2n_u} = \frac{4}{48} = \frac{1}{12} \approx 0.083.$$

Empirically, the nuclear spin-orbit splitting is of order 0.05–0.15 $\hbar\omega_0$ for light to medium nuclei, in agreement with this estimate. The direction of the splitting is such that the sub-shell of highest angular momentum $j = l + \frac{1}{2}$ at shell n is pushed down into shell $n-1$, carrying $2(2n+1)$ states with it. This is the standard mechanism that produces the magic numbers; within PDL, the magnitude and direction of the splitting are determined by $\Delta n = 4$.

The resulting construction of all seven magic numbers is:

Contribution	Sub-shells	States	Cumul
Shell $n = 0$	$1s$	+2	2
Shell $n = 1$	$1p$	+6	8
Shell $n = 2$	$2s + 1d$	+12	20
Split: $1f_{7/2}$ descends		+8	28
Shell $n = 3$ residual	$2p + 1f_{5/2} + 1g_{9/2}$	+22	50
Shell $n = 4$ residual	$2d + 3s + 1g_{7/2} + 1h_{11/2}$	+32	82
Shell $n = 5$ residual	$2f + 3p + 1h_{9/2} + 1i_{13/2}$	+44	126

The explicit derivation of why $\Delta n = 4$ generates exactly these splittings, through a combinatorial analysis of the signed-graph structure of the collective surface, is identified as an open problem in Section 8.

7 Implications for the periodic table

The results of the preceding sections collectively provide a structural account of the major organisational features of the periodic table of elements. We summarise what is derived, what is conjectured, and what remains open.

Why nuclei exist. Two protons alone cannot coexist: their active surfaces generate $T_{pp} \approx 22.84$ units of phase conflict per pair. A neutron inserted between them provides $T \approx 25.26$ mixed coherent triangles, absorbing the conflict and forming a collective closure. The nucleus is not held together by a force; it is a relational structure whose internal coherence is maintained by the balanced exchange of Section 4.

Why $N \approx Z$ for light nuclei. For $Z < Z_{\text{sat}} = 20$, each neutron can simultaneously engage all Z protons. The stability condition $N \cdot T \geq Z(Z-1)T_{pp}/2$ is satisfied with $N \approx Z$ for small Z. This is the symmetric region of the chart of nuclides.

Why $N > Z$ for heavy nuclei. For $Z > Z_{\text{sat}} = 20$, each neutron saturates and can only engage Z_{sat} protons. More neutrons are needed to compensate the growing proton–proton conflict. The valley of stability bends away from $N = Z$ at calcium and follows the curve imposed by the sea budget.

Why ^{209}Bi is the last stable nucleus. The sea-exhaustion condition gives $N_{\text{crit,max}} = 126.1$. No nucleus can accommodate more than 126 neutrons in a stable collective closure. Above $Z = 83$, no configuration $(Z, N \leq 126)$ can simultaneously satisfy both the stability condition and

the closure condition. Radioactivity is not probabilistic decay; it is the structural impossibility of maintaining a coherent collective closure.

The parity rule and Mattauch's rule. The combinatorial preference for even–even nuclei (172 stable isotopes with Z even and N even, versus only 4 with both odd) reflects the pairing principle: two surfaces of the same type close more efficiently in the antiparallel configuration. Mattauch's rule (Z odd implies N even in 57 of 61 cases) reflects the compensation principle: an odd proton surface requires a paired neutron sea to achieve collective closure.

The magic numbers. The sequence $2, 8, 20, 28, 50, 82, 126$ arises from the harmonic oscillator structure of the growing collective surface, modified by the spin-orbit splitting generated by $\Delta n = 4$. The origin of the splitting in the valence-core asymmetry of the proton is identified, but the explicit combinatorial derivation is an open problem (Section 8).

What remains outside this framework. The chemical properties of the elements — valence, conductivity, reactivity — require the coupling between the nucleus and the electron cloud. This corresponds to the PDL Block III transition (from nuclei to atoms and molecules), which is not addressed in the present paper. Likewise, the structural impossibility of stable isotopes at $Z = 43$ (technetium) and $Z = 61$ (promethium) has been identified as a combinatorial problem but not yet derived analytically.

8 Discussion and open problems

8.1 Status of the results

The following table classifies the results of this paper according to their epistemic status within the PDL programme.

Result	Status	Precision
Neutron quintuplet $(24, 28, 1032, 9960, 10992)$	Derived	Exact
Relation $R_{\text{tot}}(p) \approx 6\mu_p$	Observed	0.001%
Permutation cost $(\Delta n + 1)^2 = 25$	Derived	Exact
Gap decomposition: $40 = 25 + 15$	Derived	0.5%
Binding unit $T \approx 25$	Derived	1%
Identity: deficit = binding capacity	Derived	1%
Saturation threshold $Z_{\text{sat}} = 20$	Derived	0.7%
Valley of stability (qualitative shape)	Derived	—
Maximum magic number $N_{\text{crit,max}} = 126.1$	Derived	0.1%
He-4 as minimum of Ω for $Z = 2$	Derived	Exact
Origin of spin-orbit coupling: $\Delta n = 4$	Conjecture	Identified
Full sequence of magic numbers $2, 8, 20, 28, 50, 82, 126$	Conjecture	Verified

The entries labelled *derived* follow from the PDL proton integers and the neutron architecture established in Section 3, without free parameters. The entries labelled *conjecture* identify a structural mechanism whose explicit combinatorial derivation is incomplete.

8.2 Open problems

The following problems are identified as natural continuations of this work within the PDL programme.

OP1 (Shell splitting from Δn). Derive analytically, within the signed-graph formalism of PDL, the precise mechanism by which the valence-core asymmetry $\Delta n = n_d - n_u = 4$ generates a spin-orbit splitting of magnitude $\Delta n/(2n_u) \approx 0.083\,\hbar\omega_0$, producing the sub-shell descent responsible for the intermediate magic numbers.

OP2 (Isotopic window width $\Delta N(Z)$). Derive the width of the stable isotope window for each element from the nuclear coordination number $c(Z, N)$, i.e. the number of nucleon neighbours with which each neutron shares direct coherent triangles. The present framework gives the correct qualitative trend but overestimates ΔN for heavy elements.

OP3 (Lacunae at $Z = 43$ and $Z = 61$). Prove that the $(u, d, d)^{43}$ and $(u, d, d)^{61}$ configurations cannot form any stable collective surface closure, regardless of N, as a theorem within the PDL coherence framework.

OP4 (Chemical properties). Extend the framework to the coupling between the nuclear collective surface and the electron cloud (Block III of the PDL hierarchy). The active surface $R_{\text{surf}}(p) \approx 502$ already mediates the proton–electron coupling in hydrogen; its extension to multi-proton, multi-electron systems should account for valence, conductivity, and reactivity patterns of the periodic table.

Conclusion

This paper has established three principal results within the PDL framework, all derived without free parameters from the proton quintuplet $(24, 28, 930, 10087, 11017)$ and the CODATA values of the proton-to-electron and neutron-to-electron mass ratios.

First, the neutron quintuplet $(24, 28, 1032, 9960, 10992)$ is derived for the first time from structural principles, via the exact relation $R_{\text{tot}}(n) = R_{\text{tot}}(p) - (\Delta n + 1)^2$. The permutation cost $(\Delta n + 1)^2 = 25$ is decomposed into a structural contribution of 25 (combinatorial cost of the $u \to d$ exchange) and a mass fatigue contribution of approximately 15 (coherence dissipated by the symmetric architecture), summing to the gap of 40 between the natural budget $6\mu_n$ and the actual $R_{\text{tot}}(n)$.

Second, the nuclear binding mechanism is formalised: the same quantity $T = R_{\text{surf}}(p)^2/R_{\text{sea}}(n) \approx 25$ simultaneously measures the neutron's deficit and its binding capacity per proton. The stability condition $N \cdot T \geq Z(Z-1)T_{pp}/2$ generates the valley of stability, with a saturation threshold $Z_{\text{sat}} \approx 20$ (calcium) that separates the symmetric from the neutron-rich regime.

Third, the sea-exhaustion condition gives $N_{\text{crit,max}} = R_{\text{sea}}(p)/(2 \times \text{gap}) = 126.1$, reproducing the largest nuclear magic number and identifying ^{209}Bi as the last stable nucleus through structural impossibility rather than probabilistic decay.

The mechanism generating the intermediate magic numbers $2, 8, 20, 28, 50, 82$ is identified as the superposition of a harmonic oscillator and a spin-orbit splitting controlled by $\Delta n = 4$, with a splitting energy $\Delta n/(2n_u) \approx 0.083\,\hbar\omega_0$ in agreement with empirical values. The derivation of this splitting from the combinatorial structure of the collective surface is the principal open problem identified for future work.

Taken together, these results constitute a structural account of the periodic table's architecture from the relational axioms of PDL: the chart of nuclides as a map of collective closures, bounded by sea exhaustion and shaped by the internal asymmetry of the proton.

References

[1] C. Amsler et al. Review of particle physics. *Physics Letters B*, 667(1–2):1–134, 2008. Particle Data Group; reference for the 9%/91% valence/sea partition of the proton momentum.

[2] Maria Goeppert-Mayer. On closed shells in nuclei. *Physical Review*, 75:1969–1970, 1949. Original proposal of nuclear magic numbers via spin-orbit coupling; the PDL provides a structural origin for this coupling via $\Delta n = n_d - n_u = 4$.

[3] H1 and ZEUS Collaborations. Combined measurement and QCD analysis of the inclusive $e^{\pm}p$ scattering cross sections at HERA. *JHEP*, 01:109, 2010. arXiv:0911.0884 [hep-ex]; confirms $\sim 91\%$ gluon/sea contribution to proton momentum.

[4] O. Haxel, J. H. D. Jensen, and H. E. Suess. On the "Magic Numbers" in nuclear structure. *Physical Review*, 75:1766, 1949. Simultaneous independent derivation of magic numbers via spin-orbit coupling.

[5] F. G. Kondev et al. The NUBASE2020 evaluation of nuclear physics properties. *Chinese Physics C*, 45:030001, 2021. Reference for stable isotope data, nuclear masses, and binding energies used in Sections 4–5.

[6] Kenneth S. Krane. *Introductory Nuclear Physics*. Wiley, 1988. Standard reference for nuclear shell model, magic numbers, and nuclear stability patterns.

[7] Cédric Laubscher. Coherence leakage, hierarchical filtering, and the exponent 18 in the projective dynamic logo framework. Zenodo, 2026. Systematic justification of the 18 coherence filters (Blocks I–III) and the gravitational scaling $G_{\text{eff}} \sim \varepsilon^{18}$.

[8] Cédric Laubscher. The emergence of physical reality within the framework of the projective dynamic logo (PDL). Zenodo, 2026. Primary reference for the PDL framework, proton architecture, and derivation of α, G, k_B.

[9] Cédric Laubscher. From discrete coherence flux to effective fields: A structured programme within the PDL framework. Zenodo, 2026. Discrete Gauss- and Faraday-type relations from mixed-triangle fluxes; $1/r^2$ Coulomb profiles; Schrödinger-type dynamics.

[10] Cédric Laubscher. Introduction to the projective dynamic logo (PDL). Zenodo, 2026. Introductory presentation of the PDL axioms and minimal closure structures.

[11] Cédric Laubscher. L'émergence de la réalité physique dans le cadre du Logo Dynamique Projectif (LDP). Zenodo, 2026. French version of the PDL main document.

[12] Cédric Laubscher. A minimal relational sketch for the emergence of the golden ratio in PDL. Zenodo, 2026. Self-similar core–surface partition; emergence of φ as relational optimisation factor; Document D18 of the corpus.

[13] Cédric Laubscher. On the combinatorial selection and local uniqueness of the PDL proton architecture. Zenodo, 2026. Refined uniqueness conjecture and selection functional S; Document D16 of the corpus.

[14] Cédric Laubscher. On the combinatorial selection of the proton architecture in PDL: Axiomatic constraints, integer structures, and a uniqueness conjecture (v2). Zenodo, 2026. Formal derivation of the proton quintuplet $(24, 28, 930, 10087, 11017)$ and the selection functional $\mathcal{S}$; uniqueness conjecture.

[15] Cédric Laubscher. PDL: Global mapping of structures, results, and open problems (v8). Zenodo, 2026. Comprehensive map of the PDL corpus; dependency diagram; list of established results and open problems; Document D_Map v8.

[16] Cédric Laubscher. Projective dynamic logo as a foundational research programme: Current status, open problems, and interface with the theory of objectivity. Zenodo, 2026. Position paper delineating structurally robust results from conjectural components; Document D3.

[17] Cédric Laubscher. Relational emergence of Born's rule and a golden-ratio active surface in PDL (v2). Zenodo, 2026. Derivation of Born's rule for spin-$\frac{1}{2}$ systems and refinement of the golden-ratio connection to the fine-structure constant.

[18] Cédric Laubscher. Schrödinger-type dynamics and coherence flux in the PDL framework. Zenodo, 2026. Formal derivation of Schrödinger-type dynamics from discrete coherence flux in PDL.

[19] Cédric Laubscher. A structural derivation of the fine-structure constant from the projective dynamic logo (PDL) (v2). Zenodo, 2026. Derivation of $\alpha_{\mathrm{PDL}}^{-1} \approx 137.022$ from R_{surf}^2/μ; agreement at $\mathcal{O}(10^{-4})$.

[20] Cédric Laubscher. A topological reformulation in the projective dynamic logo framework. Zenodo, 2026. Topological and graph-theoretic extensions of the PDL coherence framework; connection to entanglement and emergent metric.

[21] Cédric Laubscher. Universal coherence leakage and the gravitational bridge formula in PDL (v3). Zenodo, 2026. Primary result: bridge formula $\Delta r_{\mathrm{val}} = \varepsilon_G \cdot (R_{\mathrm{tot}} R_{\mathrm{surf}}/r_{\mathrm{val}}^2) \cdot (n_u^2 - 1)/n_u^2$; $G_{\mathrm{PDL}} = 6.67448 \times 10^{-11}$ (Document D20-V3).

[22] NIST. CODATA 2022 recommended values of the fundamental physical constants. `https://physics.nist.gov/cuu/Constants/`, 2022. Source for G, α, $\mu = m_p/m_e$, $\mu_n = m_n/m_e$ used throughout this work.

[23] Carlo Rovelli. Relational quantum mechanics. *International Journal of Theoretical Physics*, 35:1637–1678, 1996. Relational interpretation of quantum mechanics; conceptual background for the PDL relational framework.

[24] C. F. von Weizsäcker. Zur Theorie der Kernmassen. *Zeitschrift für Physik*, 96:431–458, 1935. Semi-empirical mass formula; the PDL derives the Coulomb-to-volume ratio $a_C/a_V \approx R_{\mathrm{surf}}/R_{\mathrm{sea}}$ structurally.