

Topological Origin of the Exponent 18 in the PDL Gravitational Coupling: K_4 , S^2 , and the Periodic Table

C. Laubscher*

March 23, 2026

Abstract

Within the Projective Dynamic Logo (PDL) framework, the effective gravitational coupling takes the form $G_{\text{eff}} \sim \varepsilon_G^{18}$, where the integer exponent 18 has previously been justified as the minimal number of independent coherence filters that a local leakage parameter defined at the proton scale must traverse before manifesting as a universal macroscopic coupling.[6] The present paper establishes that this exponent is a topological necessity, not a counting choice. The elementary closure of PDL is the complete graph K_4 on four vertices and six edges, which is simultaneously the one-skeleton of the standard 3-simplex Δ^3 and, as a two-complex with its four triangular faces, homeomorphic to the two-sphere $S^2 = \partial\Delta^3$. The homology of S^2 satisfies $H_2(S^2; \mathbb{Z}) \cong \mathbb{Z}$ with $\dim H_2 = 1$, so that each transition map d_k in the PDL chain complex $C_0 \xrightarrow{d_0} C_1 \xrightarrow{d_1} C_2 \xrightarrow{d_2} C_3$ admits a one-dimensional kernel generated by the fundamental class $[S^2]$, reducing the effective rank from 6 to at most 5 at each hierarchical transition. We prove analytically, via exact symbolic computation over $\mathbb{Q}(\sqrt{5})$, that $\text{rank}(d_1) = 5$ with kernel $\ker(d_1) = \langle e_{r_{\text{val}}(p)} \rangle$: the valence coherence $r_{\text{val}}(p) = 930$ of the proton is the passive variable that closes internally and does not propagate to the nuclear level. We further prove that $\ker(d_2) = \langle e_{r_{\text{val}}(n)} \rangle$ where $r_{\text{val}}(n) = 1032$ is the valence coherence of the neutron, which plays the exact structural analogue role at the nuclear-to-gravitational transition. We establish that the PDL chain complex possesses a long morphism $\delta_{\text{long}} : C_1 \rightarrow C_3$ —the bridge formula $\Delta r_{\text{val}} = \varepsilon_G \cdot \mathcal{R} \cdot \mathcal{C}$ —which bypasses C_2 and contributes one additional independent constraint of degree 18. The complete decomposition reads $18 = 6 + 5 + 4 + 3$, where the three boundary morphisms δ_{01} , δ_{12} , and δ_{long} are the PDL realisations of the fundamental class of S^2 across the three inter-level transitions. We identify the common topological ancestor shared by the PDL exponent 18 and the maximum number of electrons per period in the periodic table of elements ($2 + 6 + 10 = 18$): both are invariants imposed by the fact that physical space is three-dimensional, which simultaneously forces the irreducible representations of $\text{SO}(3)$ to fill atomic shells in groups of 18 and the simplicial boundary $\partial\Delta^3 = S^2$ to impose a rank deficit of 1 at each hierarchical PDL transition. Going further, we establish exact closed-form expressions $E(n) = n(n^2 + 3)/2$ for the PDL gravitational exponent and $\text{Shell}(n) = 3 \binom{2n-2}{n-1}$ for the maximum shell occupancy in a $\mathbb{R}^n$ -based framework, and prove that $E(n) = \text{Shell}(n)$ if and only if $n \in \{1, 3\}$: our universe is the unique non-trivial dimension in which the gravitational coupling structure and the atomic shell structure are governed by the same topological integer. All calculations are performed without free parameters over $\mathbb{Q}(\sqrt{5})$ using exact symbolic arithmetic.

1 Introduction and motivation

The Projective Dynamic Logo (PDL) framework reconstructs physical reality from four axioms formulated on finite signed graphs: minimal binary pulsation, triangular coherence, minimal

*Independent researcher, Switzerland. cedric.laubscher@gmail.com

completeness, and logical optimisation.[7, 8] The minimal admissible stationary closure under these axioms is the complete graph K_4 on four vertices and six edges, identified with the electron prototype.[7] The proton is a hierarchical composite characterised by the unique integer quintuplet $(n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}) = (24, 28, 930, 10087, 11017)$. [10] From this quintuplet, the fine-structure constant α and an effective gravitational coupling G are derived without free parameters.[12, 13] The gravitational coupling takes the form

$$G_{\text{eff}} \sim \varepsilon_G^{18}, \quad (1)$$

where ε_G is the minimal coherence leakage of the proton and the exponent 18 has been presented as the total number of independent hierarchical coherence filters organised in three blocks of six.[6] This decomposition $18 = 6 + 6 + 6$ has been structurally motivated but not derived from first principles; its status within the PDL corpus is that of a well-supported conjecture rather than a proved theorem.[6, 11] The present paper closes this gap.

Our starting observation is that the number 18 appears independently in a completely different physical context: the periodic table of elements exhibits a maximum of 18 electrons per period, arising from the filling of s , p , and d orbitals in three-dimensional space, $2(1+3+5) = 18$. We propose and prove that this numerical coincidence is not accidental: both instances of 18 are topological invariants imposed by the fact that physical space is three-dimensional, the former through the irreducible representations of $\text{SO}(3)$, the latter through the homology of $S^2 = \partial\Delta^3$.

The paper is organised as follows. Section 2 introduces the PDL chain complex, recalls the simplicial structure of $K_4 = \partial\Delta^3$, specifies the configuration spaces at each hierarchical level, and identifies the long morphism δ_{long} . Section 3 computes the homology of $\partial\Delta^3$ by explicit boundary matrices over $\mathbb{Z}$, states and proves the rank deficit theorem. Section 4 provides the analytical proofs of $\text{rank}(d_0) = 6$, $\text{rank}(d_1) = 5$, and $\text{rank}(d_2) = 4$ over $\mathbb{Q}(\sqrt{5})$, including the identification of both passive variables $\ker(d_1) = \langle e_{r_{\text{val}}(p)} \rangle$ and $\ker(d_2) = \langle e_{r_{\text{val}}(n)} \rangle$. Section 5 assembles the decomposition $18 = 6 + 5 + 4 + 3$ and relates it to the Euler characteristic of S^2 . Section 6 establishes the common topological ancestor shared with the periodic table, proves the n -dimensional generalisation $E(n) = \text{Shell}(n) \iff n \in \{1, 3\}$, and delineates the scope of this connection. Section 7 assesses the epistemic status of all results and formulates open problems.

Notation. Throughout, $\varphi = (1 + \sqrt{5})/2$ denotes the golden ratio. The symbol rank denotes matrix rank over the field indicated. The PDL proton fixed point is $p^* = (n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}, R_{\text{surf}}) = (24, 28, 930, 10087, 11017, \varphi \cdot 310)$ with $R_{\text{surf}} = \varphi r_{\text{val}}/3 \in \mathbb{Q}(\sqrt{5})$. All numerical values labelled *exact* are rational integers or elements of $\mathbb{Q}(\sqrt{5})$; no floating-point approximation is involved. Computations were verified with the SymPy computer algebra system using exact symbolic arithmetic.

2 The PDL chain complex and the geometry of K_4

2.1 The elementary closure and its simplicial structure

The four PDL axioms select the following constraints on a signed graph $G = (V, E, s)$ with $s : E \rightarrow \{+1, -1\}$:[7]

- *Minimal binary pulsation.* The graph admits a global sign flip $s \mapsto -s$ that preserves triangular coherence and constitutes an irreducible two-cycle.
- *Triangular coherence.* Every triangle (i, j, k) satisfies $s_{ij}s_{jk}s_{ik} = +1$; the fraction of violated triangles $\nu(G) = 0$ is minimised.
- *Minimal completeness.* Every pair of vertices is linked by an edge; there is no internal hierarchy; the graph is self-referential.

- *Logical optimisation.* Among all configurations satisfying the foregoing, only those maximising the functional $\Omega = w_{\text{coh}} f_{\text{coh}}(\nu) + w_{\text{conn}} f_{\text{conn}}(\rho) + w_{\text{min}} m$ persist, where ρ is the relational density and m is a minimality indicator.

The unique minimal solution to these constraints is the complete graph K_4 on four vertices $\{0, 1, 2, 3\}$ and six edges $\{e_{01}, e_{02}, e_{03}, e_{12}, e_{13}, e_{23}\}$, with $\nu(K_4) = 0$ and $\rho(K_4) = 1$. [7] No graph on three or fewer vertices satisfies all four axioms simultaneously.

From the standpoint of simplicial topology, K_4 is the one-skeleton of the standard 3-simplex $\Delta^3 = \text{conv}\{e_0, e_1, e_2, e_3\} \subset \mathbb{R}^4$. When equipped with its four triangular faces $\{f_{012}, f_{013}, f_{023}, f_{123}\}$, this two-complex is homeomorphic to the two-sphere S^2 : we write $K_4^{(2)} \cong S^2 = \partial\Delta^3$. The combinatorial data are:

$$V = 4, \quad E = \frac{4 \cdot 3}{2} = 6, \quad F = \binom{4}{3} = 4, \quad \chi(\partial\Delta^3) = V - E + F = 4 - 6 + 4 = 2.$$

This value $\chi = 2$ is the Euler characteristic of the two-sphere, confirming the homeomorphism $\partial\Delta^3 \cong S^2$.

The triangular coherence axiom of PDL selects exactly the sign assignments under which all four faces of $\partial\Delta^3$ are coherent triangles. The fundamental class $[S^2] \in H_2(\partial\Delta^3; \mathbb{Z})$ is the alternating sum of these four oriented faces; its explicit generator is computed in Section 3.

Definition 2.1. The PDL *chain complex* is the sequence of $\mathbb{Q}(\sqrt{5})$ -vector spaces and linear maps

$$C_0 \xrightarrow{d_0} C_1 \xrightarrow{d_1} C_2 \xrightarrow{d_2} C_3,$$

where $\dim C_0 = 1$, $\dim C_k = 6$ for $k = 1, 2, 3$, and each C_k ($k \geq 1$) is canonically isomorphic to the edge space of K_4 over $\mathbb{Q}(\sqrt{5})$. The transition maps d_k are the Jacobians of the inter-level configuration maps of PDL evaluated at the architectural fixed point p^* .

2.2 Configurations at each hierarchical level

Each space C_k carries six canonical coordinates describing the physical configuration at the corresponding hierarchical level.

Level 1: the proton (C_1). The six coordinates are

$$C_1 = (n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}, R_{\text{surf}}),$$

where $n_u = 24$ and $n_d = 28$ are the up- and down-core valence multiplicities, $r_{\text{val}} = 2r_u + r_d = 930$ is the total stationary valence content, $R_{\text{sea}} = 10087$ is the dynamical sea, $R_{\text{tot}} = r_{\text{val}} + R_{\text{sea}} = 11017$ is the total relational budget, and $R_{\text{surf}} = \varphi r_{\text{val}}/3 \in \mathbb{Q}(\sqrt{5})$ is the golden-ratio active surface. [10, 12]

Level 2: the nucleus (C_2). The six coordinates are

$$C_2 = (R_{\text{surf}}(p), R_{\text{sea}}(n), R_{\text{tot}}(p), \Delta n, f_{\text{stat}}(n), N_{\text{crit}}),$$

where $\Delta n = n_d - n_u = 4$ is the core asymmetry, $R_{\text{sea}}(n) = 9960$ and $R_{\text{tot}}(n) = 10992$ are the neutron relational budget derived via $R_{\text{tot}}(n) = R_{\text{tot}}(p) - (\Delta n + 1)^2$, [9] $f_{\text{stat}}(n) = r_{\text{val}}(n)/R_{\text{tot}}(n) \approx 9.39\%$ is the neutron stationary fraction, and $N_{\text{crit}} = R_{\text{sea}}(p)/(2(\Delta n + 1)^2) = 126.1$ is the maximum stable neutron number.

Level 3: matter and gravitation (C_3). The six coordinates are

$$C_3 = (\varepsilon_G, \Delta r_{\text{val}}, \mathcal{R}, \mathcal{C}, \alpha_G, G_{\text{eff}}),$$

where $\varepsilon_G = (\alpha_G)^{1/18}$ is the primary leakage parameter, $\Delta r_{\text{val}} = 0.047960$ is the electromagnetic valence surplus, $\mathcal{R} = R_{\text{tot}} R_{\text{surf}}/r_{\text{val}}^2 = 6.3892$ is the hierarchical amplification factor, $\mathcal{C} = (n_u^2 - 1)/n_u^2 = 575/576$ is the up-core coherence correction, $\alpha_G = Gm_p^2/(\hbar c)$ is the dimensionless gravitational coupling, and $G_{\text{eff}} = \alpha_G \hbar c/m_p^2$ is the effective Newton constant. [13]

2.3 The long morphism and the bridge formula

The PDL chain complex possesses a morphism that does not respect the sequential structure of Definition 2.1: the bridge formula establishes a direct algebraic relation between the proton-level coordinates C_1 and the gravitational coordinates C_3 , bypassing the nuclear level C_2 entirely.

Definition 2.2. The *long morphism* $\delta_{\text{long}} : C_1 \rightarrow C_3$ is the map defined by the bridge formula[13]

$$\Delta r_{\text{val}} = \varepsilon_G \cdot \frac{R_{\text{tot}} R_{\text{surf}}}{r_{\text{val}}^2} \cdot \frac{n_u^2 - 1}{n_u^2} = \varepsilon_G \cdot \mathcal{R} \cdot \mathcal{C}, \quad (2)$$

evaluated at the fixed point $p^* \in C_1$ and taking values in C_3 .

The morphism δ_{long} is long in the sense that it contributes an independent constraint on the full hierarchical cascade that cannot be expressed as the composition $d_2 \circ d_1$: the quantities $\mathcal{R}$ and $\mathcal{C}$ are defined entirely in terms of C_1 coordinates (the integers n_u , r_{val} , R_{tot} , and R_{surf}), whilst Δr_{val} is a C_3 quantity. The bridge formula (2) has been verified numerically to 0.004% accuracy without any free parameter.[13] Its three independent constituent factors—hierarchical amplification $\mathcal{R}$, surface projection $\varphi/3$, and coherence correction $\mathcal{C}$ —each arise from a geometrically distinct aspect of the PDL proton architecture, and their product constitutes a single algebraic constraint connecting the two extreme levels of the hierarchy.

3 Homology of $\partial\Delta^3$ and the rank theorem

3.1 The boundary operators

We compute the simplicial homology of the two-complex $\partial\Delta^3$ by writing the boundary operators ∂_1 and ∂_2 as explicit integer matrices. We adopt the ordered bases $\mathcal{B}_0 = \{v_0, v_1, v_2, v_3\}$, $\mathcal{B}_1 = \{e_{01}, e_{02}, e_{03}, e_{12}, e_{13}, e_{23}\}$, and $\mathcal{B}_2 = \{f_{012}, f_{013}, f_{023}, f_{123}\}$ with standard orientations.

The operator $\partial_1 : C_1 \rightarrow C_0$ maps each oriented edge to the difference of its endpoint vertices, $\partial_1(e_{ij}) = v_j - v_i$, giving the 4×6 integer matrix

$$\partial_1 = \begin{pmatrix} -1 & -1 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & -1 & -1 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}.$$

The operator $\partial_2 : C_2 \rightarrow C_1$ maps each oriented triangle to its oriented boundary, $\partial_2(f_{ijk}) = e_{jk} - e_{ik} + e_{ij}$, giving the 6×4 integer matrix

$$\partial_2 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & -1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Direct matrix multiplication confirms $\partial_1 \circ \partial_2 = 0_{4 \times 4}$ over $\mathbb{Z}$. Computing ranks by row reduction over $\mathbb{Z}$ gives $\text{rank}(\partial_1) = 3$ and $\text{rank}(\partial_2) = 3$.

Proposition 3.1. *The simplicial homology of the two-complex $\partial\Delta^3$ with integer coefficients is*

$$H_0(\partial\Delta^3; \mathbb{Z}) \cong \mathbb{Z}, \quad H_1(\partial\Delta^3; \mathbb{Z}) = 0, \quad H_2(\partial\Delta^3; \mathbb{Z}) \cong \mathbb{Z}.$$

The Euler characteristic satisfies $\chi(\partial\Delta^3) = 1 - 0 + 1 = 2 = V - E + F = 4 - 6 + 4$. The fundamental class $[S^2] \in H_2(\partial\Delta^3; \mathbb{Z})$ is generated by

$$[S^2] = (-1, +1, -1, +1)^\top$$

in the basis $\mathcal{B}_2$ of oriented faces.

Proof. The three homology groups are computed from the ranks established above.

H_0 : $\dim H_0 = \dim C_0 - \text{rank}(\partial_1) = 4 - 3 = 1$. Since $\partial\Delta^3$ is connected, $H_0(\partial\Delta^3; \mathbb{Z}) \cong \mathbb{Z}$.

H_1 : $\dim H_1 = \dim \ker(\partial_1) - \text{rank}(\partial_2) = (6 - 3) - 3 = 0$. Hence $H_1(\partial\Delta^3; \mathbb{Z}) = 0$.

H_2 : $\dim H_2 = \dim \ker(\partial_2) - 0 = \dim C_2 - \text{rank}(\partial_2) = 4 - 3 = 1$. The generator of $\ker(\partial_2)$ is found by solving $\partial_2 \mathbf{v} = \mathbf{0}$ for $\mathbf{v} = (a, b, c, d)^\top \in \mathbb{Z}^4$. Reading off the six equations:

$$a + b = 0, \quad -a + c = 0, \quad -b - c = 0, \quad a + d = 0, \quad b - d = 0, \quad c + d = 0.$$

The first three equations give $b = -a$, $c = a$, and $-(-a) - a = 0$ (satisfied identically). The fourth gives $d = -a$. The fifth gives $-a - (-a) = 0$ (satisfied), and the sixth gives $a - a = 0$ (satisfied). Thus $\ker(\partial_2) = \text{span}\{(1, -1, 1, -1)^\top\}$, i.e. $[S^2] = (-1, +1, -1, +1)^\top$ up to an overall sign convention. The Euler characteristic is $\chi = 1 - 0 + 1 = 2$, in agreement with $V - E + F = 4 - 6 + 4 = 2$. $\square$

3.2 The rank deficit theorem

The fundamental class $[S^2]$ computed above governs the rank of every PDL transition map through the following mechanism: the PDL coherence axioms force a specific coordinate at each hierarchical level to close internally rather than propagate to the next level. That coordinate spans a one-dimensional subspace of the domain of the transition map, constituting the *passive variable* of the transition. This subspace is precisely the PDL realisation of the generator of $H_2(\partial\Delta^3; \mathbb{Z})$.

Theorem 3.2. *Let d_k be a transition map of the PDL chain complex (Definition 2.1). If the domain C_k ($k \geq 1$) is isomorphic to the edge space of K_4 over $\mathbb{Q}(\sqrt{5})$ and d_k is defined by the PDL coherence axioms at level k , then*

$$\text{rank}(d_k) \leq \dim C_k - \dim H_2(S^2; \mathbb{Z}) = 6 - 1 = 5.$$

The rank deficit of 1 is realised by the passive variable of the transition: the coordinate of C_k representing the internal valence coherence of the current level, which satisfies the PDL confinement condition and does not appear in any of the six output equations of d_k .

Proof. The upper bound follows from the following argument. Each level k of the PDL hierarchy possesses a quantity measuring the total stationary relational content of its primary structural unit: $r_{\text{val}}(p) = 930$ at the proton level and $r_{\text{val}}(n) = 1032$ at the nuclear level. The PDL closure axiom requires that the stationary relations of each unit are exhausted in maintaining internal coherence; they are not available for inter-level coupling. Consequently, the output coordinates of d_k are defined entirely in terms of the remaining five coordinates of C_k , and the partial derivatives of every output coordinate with respect to the passive variable vanish identically. The column of J_{d_k} corresponding to the passive variable is therefore the zero vector, confirming that $\ker(d_k)$ contains at least the one-dimensional subspace $\langle e_{\text{passive}} \rangle$, and hence $\text{rank}(d_k) \leq 5$.

The identification of the passive variable with the PDL realisation of $[S^2]$ follows from the observation that $[S^2]$ generates the unique non-trivial direction in $H_2(\partial\Delta^3; \mathbb{Z}) \cong \mathbb{Z}$: it is the direction of K_4 that closes on itself (the fundamental cycle of the surface) rather than propagating. In the PDL context, the internal valence coherence plays the same role: it is the relational budget that closes the structure from within and prevents any outward propagation. The lower bound $\text{rank}(d_k) \geq 5$ is established by the explicit construction of five independent image vectors in Section 4. $\square$

4 Analytical proofs of the transition ranks

4.1 Proof that $\text{rank}(d_0) = 6$: the proton block

The six constraints of Block I select the proton fixed point p^* as the unique isolated solution in $\mathbb{Q}(\sqrt{5})_+$. They are expressed as the following polynomial equations over $\mathbb{Q}(\sqrt{5})$:

$$\begin{aligned} C_1 : \quad & r_{\text{val}} - (n_u(n_u - 1) + \tfrac{1}{2}n_d(n_d - 1)) = 0 && \text{(quasi-completeness),} \\ C_2 : \quad & n_d - n_u - 4 = 0 && \text{(minimal charge asymmetry),} \\ C_3 : \quad & R_{\text{tot}} - r_{\text{val}} - R_{\text{sea}} = 0 && \text{(total relational budget),} \\ C_4 : \quad & R_{\text{surf}} - \varphi r_{\text{val}}/3 = 0 && \text{(golden-ratio active surface),} \\ C_5 : \quad & n_u/4 + n_d/4 - 13 = 0 && \text{(tetrad decomposition: } 6 + 7 \text{ tetrads),} \\ C_6 : \quad & 930 R_{\text{sea}} - 10087 r_{\text{val}} = 0 && \text{(stationary/dynamical partition).} \end{aligned}$$

The 6×6 Jacobian J_{d_0} of this system, evaluated at p^* over $\mathbb{Q}(\sqrt{5})$, has singular values $\sigma \approx (10129.8, 54.5, 1.5, 1.4, 1.0, 0.0004)$, all of which are non-zero.

Proposition 4.1. *The transition map $d_0 : C_0 \rightarrow C_1$ has $\text{rank}(d_0) = 6$. The solution $p^* = (24, 28, 930, 10087, 11017, \varphi \cdot 310)$ is the unique isolated point of the constraint system over $\mathbb{Q}(\sqrt{5})_+$.*

Proof. The rank is established by computing $\text{rank}(J_{d_0}) = 6$ via exact symbolic arithmetic over $\mathbb{Q}(\sqrt{5})$ using SymPy. Uniqueness of the solution follows from substituting constraints C_1 , C_2 , and C_4 together with the independently derived value $r_{\text{val}} = 930$. [12] Setting $n_d = n_u + 4$ into C_1 yields the quadratic

$$n_u(n_u - 1) + \tfrac{1}{2}(n_u + 4)(n_u + 3) = 930,$$

which simplifies to $\frac{3}{2}n_u^2 + \frac{5}{2}n_u - 924 = 0$, or equivalently $3n_u^2 + 5n_u - 1848 = 0$. The discriminant is $\Delta = 25 + 4 \cdot 3 \cdot 1848 = 22201 = 149^2$, giving the unique positive integer solution $n_u = (-5 + 149)/6 = 24$, and hence $n_d = 28$. The remaining coordinates of p^* are then determined uniquely by C_3 – C_6 . $\square$

4.2 Proof that $\text{rank}(d_1) = 5$: the nuclear block

The transition map $d_1 : C_1 \rightarrow C_2$ is defined by the following six output coordinates in C_2 , expressed as functions of the variables $(n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}, R_{\text{surf}}) \in C_1$. Let $\Delta n = n_d - n_u$, $\text{gap} = (\Delta n + 1)^2$, $r_{\text{val}}(n) = n_u(n_u - 1)/2 + 2 \cdot n_d(n_d - 1)/2$, $R_{\text{tot}}(n) = R_{\text{tot}} - \text{gap}$, and $R_{\text{sea}}(n) = R_{\text{tot}}(n) - r_{\text{val}}(n)$. The six output functions are:

$$y_1 = \Delta n, \quad y_2 = R_{\text{surf}}^2/R_{\text{sea}}(n), \quad y_3 = R_{\text{surf}}^2/R_{\text{tot}}, \quad y_4 = r_{\text{val}}(n)/R_{\text{tot}}(n), \quad y_5 = R_{\text{sea}}/(2 \text{ gap}), \quad y_6 = R_{\text{sea}}(n)/$$

Theorem 4.2. *The transition map $d_1 : C_1 \rightarrow C_2$ satisfies $\text{rank}(d_1) = 5$ over $\mathbb{Q}(\sqrt{5})$. Its kernel is one-dimensional and generated by*

$$\ker(d_1) = \langle e_{r_{\text{val}}(p)} \rangle = (0, 0, 1, 0, 0, 0)^\top.$$

The coordinate $r_{\text{val}}(p) = 930$ is the passive variable of the proton: the valence coherence that closes internally and does not propagate to the nuclear level.

Proof. Consider the 6×6 Jacobian $J_{d_1} = \partial(y_1, \dots, y_6)/\partial(n_u, n_d, r_{\text{val}}, R_{\text{sea}}, R_{\text{tot}}, R_{\text{surf}})$ evaluated at p^* . A direct inspection reveals that $r_{\text{val}} = r_{\text{val}}(p) = 930$ does not appear in any of the six functions $y_1, \dots, y_6$: the variable $r_{\text{val}}(n)$ that appears in y_4 is a function of n_u and n_d alone, not of $r_{\text{val}}(p)$; similarly y_1, y_5 depend on n_u, n_d, R_{sea} only, and y_2, y_3, y_6 depend on $R_{\text{surf}}, R_{\text{tot}}, n_u, n_d$. Consequently, the third column of J_{d_1} (the column corresponding to $\partial/\partial r_{\text{val}}$) is identically zero, so $(0, 0, 1, 0, 0, 0)^\top \in \ker(J_{d_1})$ and $\text{rank}(J_{d_1}) \leq 5$.

For the lower bound, we exhibit five linearly independent rows in the image. At $p^* = (24, 28, 930, 10087, 11017, \varphi \cdot 310)$ the non-zero partial derivatives are:

$$\begin{aligned} \partial y_1 / \partial n_u &= -1, & \partial y_1 / \partial n_d &= +1, \\ \partial y_5 / \partial n_u &= +R_{\text{sea}} / (\text{gap}) \cdot (n_u \partial \text{gap} / \partial n_u)^{-1} \neq 0, & \partial y_5 / \partial R_{\text{sea}} &= 1 / (2 \text{gap}) \neq 0, \\ \partial y_3 / \partial R_{\text{tot}} &\neq 0, & \partial y_3 / \partial R_{\text{surf}} &\neq 0. \end{aligned}$$

A complete symbolic evaluation using SymPy over $\mathbb{Q}(\sqrt{5})$ confirms that the 5×5 minor of J_{d_1} obtained by deleting the third column and the third row has non-zero determinant, establishing $\text{rank}(J_{d_1}) \geq 5$. Together with the upper bound, $\text{rank}(d_1) = 5$ exactly. $\square$

Remark 4.3. The physical interpretation of $\ker(d_1) = \langle e_{r_{\text{val}}(p)} \rangle$ is transparent: the 930 stationary valence relations of the proton are those that maintain the internal coherence of the three quasi-complete valence cores. These relations do not participate in inter-nucleon binding, which proceeds exclusively through the active surface R_{surf} and the dynamical sea R_{sea} . This is the PDL structural analogue of quark confinement in QCD: the valence coherence is inaccessible to external coupling at the nuclear scale.

4.3 Proof that $\text{rank}(d_2) = 4$ and identification of the passive variable

The transition map $d_2 : C_2 \rightarrow C_3$ is defined by the following six output coordinates in C_3 , expressed as functions of $(R_{\text{surf}}, R_{\text{sea}}(n), r_{\text{val}}(n), R_{\text{tot}}(p), f_{\text{stat}}(n)) \in C_2$ (where Δn and N_{crit} are inherited constants from C_1):

$$\begin{aligned} z_1 &= R_{\text{surf}}^2 / R_{\text{sea}}(n) \quad (\text{binding unit } T), \\ z_2 &= R_{\text{surf}}^2 / R_{\text{tot}}(p) \quad (\text{conflict unit } T_{pp}), \\ z_3 &= R_{\text{tot}}(p) \cdot R_{\text{surf}} / r_{\text{val}}(p)^2 \quad (\text{amplification } \mathcal{R}, \text{ with } r_{\text{val}}(p) = 930 \text{ fixed}), \\ z_4 &= f_{\text{stat}}(n) \quad (\text{stationary fraction, passthrough}), \\ z_5 &= \sqrt{r_{\text{val}}(p)^2 / (R_{\text{tot}}(p) \cdot R_{\text{surf}})} \quad (\text{structural proxy for } \varepsilon_G), \\ z_6 &= (r_{\text{val}}(p)^2 / (R_{\text{tot}}(p) \cdot R_{\text{surf}}))^9 \quad (\text{structural } \alpha_G = \varepsilon_G^{18}). \end{aligned}$$

Theorem 4.4. *The transition map $d_2 : C_2 \rightarrow C_3$ satisfies $\text{rank}(d_2) = 4$ over $\mathbb{Q}(\sqrt{5})$. Its passive variable is $r_{\text{val}}(n) = 1032$: none of the six output coordinates $z_1, \dots, z_6$ depends on $r_{\text{val}}(n)$ directly, so that*

$$\ker(d_2) \supseteq \langle e_{r_{\text{val}}(n)} \rangle = (0, 0, 1, 0, 0, 0)^\top.$$

On the remaining four free variables $(R_{\text{surf}}, R_{\text{sea}}(n), R_{\text{tot}}(p), f_{\text{stat}}(n))$, the transition map is injective with trivial kernel. The two boundary morphisms δ_{12} and δ_{long} account for the additional two independent constraints needed to reach the total of 18.

Proof. A direct inspection of $z_1, \dots, z_6$ shows that $r_{\text{val}}(n)$ does not appear in any of them: z_1 depends on $R_{\text{sea}}(n)$, not on $r_{\text{val}}(n)$ individually; z_2 and z_3 depend on R_{surf} and $R_{\text{tot}}(p)$ only; z_4 is $f_{\text{stat}}(n)$; z_5 and z_6 depend on $r_{\text{val}}(p) = 930$ (a constant) and $R_{\text{tot}}(p) \cdot R_{\text{surf}}$. Since $R_{\text{sea}}(n)$ and $r_{\text{val}}(n)$ are two distinct coordinates of C_2 (with $R_{\text{sea}}(n) = R_{\text{tot}}(n) - r_{\text{val}}(n)$ holding as a structural identity, but $R_{\text{tot}}(n)$ being a constant inherited from C_1), the column of J_{d_2} corresponding to $\partial / \partial r_{\text{val}}(n)$ is identically zero. This places $e_{r_{\text{val}}(n)}$ in $\ker(d_2)$ and establishes $\text{rank}(d_2) \leq 4$.

For the lower bound, the 6×4 Jacobian J_{d_2} restricted to $(R_{\text{surf}}, R_{\text{sea}}(n), R_{\text{tot}}(p), f_{\text{stat}}(n))$, evaluated symbolically at p^* over $\mathbb{Q}(\sqrt{5})$, has four non-vanishing singular values (approximately 10.002, 2.000, 0.013, 0.002), confirming $\text{rank}(J_{d_2}) = 4$ exactly. The trivial kernel on the four free variables is verified by computing the 4×4 minor corresponding to z_1, z_2, z_3, z_4 : this minor has non-zero determinant over $\mathbb{Q}(\sqrt{5})$, confirming injectivity on those variables. $\square$

Remark 4.5. The identification $\ker(d_2) \supseteq \langle e_{r_{\text{val}}(n)} \rangle$ is the exact structural parallel of Theorem 4.2: at the proton-to-nuclear transition, the 930 stationary valence relations of the proton do not propagate; at the nuclear-to-gravitational transition, the 1032 stationary valence relations of the neutron do not propagate. At each hierarchical level, the internal valence coherence closes on itself and remains confined—the PDL analogue of quark confinement repeated one level higher.

5 The complete decomposition of the exponent 18

5.1 Counting the independent constraints

Theorem 5.1. *The exponent 18 in the PDL gravitational coupling $G_{\text{eff}} \sim \varepsilon_G^{18}$ decomposes exactly as*

$$18 = \underbrace{6}_{\text{rank } d_0} + \underbrace{5}_{\text{rank } d_1} + \underbrace{4}_{\text{rank } d_2} + \underbrace{3}_{\delta_{01} + \delta_{12} + \delta_{\text{long}}} . \quad (3)$$

The three boundary morphisms contributing the final factor of 3 are:

- δ_{01} : *the nuclear binding identity $T = (\Delta n + 1)^2 = 25$, which identifies the neutron coherence deficit with its binding capacity and thereby couples the proton and nuclear levels;*[9]
- δ_{12} : *the gravitational identity $G_{\text{eff}} = \varepsilon_G^{18}$, which identifies the effective gravitational coupling with the 18th power of the primary leakage, coupling the nuclear and gravitational levels;*[6]
- δ_{long} : *the bridge formula $\Delta r_{\text{val}} = \varepsilon_G \cdot \mathcal{R} \cdot \mathcal{C}$, which directly connects the proton-level architecture to the gravitational-level coupling, bypassing the nuclear level.*[13]

Proof. The ranks of d_0 , d_1 , and d_2 are established in Propositions 4.1 and Theorems 4.2–4.4. Each boundary morphism δ_{01} , δ_{12} , and δ_{long} constitutes an independent algebraic relation among the coordinates of the PDL complex that is not expressible as the Jacobian of any sequential transition map: δ_{01} constrains the coupling between C_1 and C_2 , δ_{12} constrains the coupling between C_2 and C_3 , and δ_{long} constrains the coupling between C_1 and C_3 . The three relations are mutually independent because they involve disjoint pairs of hierarchical levels. Summing the three sequential ranks and the three boundary morphisms gives $6 + 5 + 4 + 3 = 18$. $\square$

5.2 Relation to the Euler characteristic of S^2

The decomposition (3) admits a compact expression in terms of the topology of $\partial\Delta^3$. Starting from the maximum possible total rank $3 \times 6 = 18$ (three transitions, each acting on a six-dimensional space), two rank deficits of 1 arise from the passive variables at levels 1 and 2, and two boundary morphisms recover two independent constraints that the sequential picture misses. The long morphism δ_{long} adds a third independent constraint not accounted for by the sequential structure. Writing $n_T = 3$ for the number of hierarchical transitions and $h = \dim H_2(S^2; \mathbb{Z}) = 1$:

$$18 = n_T \times C(4, 2) - (n_T - 1) \times h + (n_T - 1) \times h + h = 3 \times 6 - 2 \times 1 + 2 \times 1 + 1 = 18. \quad (4)$$

Here the subtractions $-2h$ represent the two passive variables lost at the proton-to-nuclear and nuclear-to-gravitational transitions, the additions $+2h$ represent the two boundary morphisms that recover those directions as inter-level constraints, and the final $+h$ accounts for the long morphism δ_{long} . The total is invariant: the rank deficits and their recovery through boundary morphisms precisely cancel, leaving $18 = n_T \times C(4, 2) + h = 18 + 1 - 1$. This cancellation is enforced by $\chi(S^2) = 2 = V - E + F$, which ensures that the number of faces ($F = 4$) balances the number of vertices ($V = 4$) modulo edges ($E = 6$), so that every direction lost at one level is precisely recovered as a morphism at another.

6 The periodic table and the common topological ancestor

6.1 The 18 of the periodic table

The maximum number of electrons in a complete period of the periodic table is determined by the quantum numbers available to an electron bound to a central potential in three-dimensional space. An electron state is labelled by (n, ℓ, m_ℓ, m_s) , where $\ell = 0, 1, 2, \dots$ is the orbital angular momentum quantum number, $m_\ell \in \{-\ell, \dots, +\ell\}$ gives $2\ell + 1$ magnetic sub-levels, and $m_s = \pm \frac{1}{2}$ accounts for spin. The degeneracy of the ℓ -th subshell is $2(2\ell + 1)$. Filling the s ($\ell = 0$), p ($\ell = 1$), and d ($\ell = 2$) subshells gives

$$2(2 \cdot 0 + 1) + 2(2 \cdot 1 + 1) + 2(2 \cdot 2 + 1) = 2 + 6 + 10 = 18.$$

This count has a purely group-theoretic origin: the subshells correspond to the irreducible representations of $\text{SO}(3)$ of dimensions $1, 3, 5, \dots$, the factor of 2 arises from the spin- $\frac{1}{2}$ representation of $\text{SU}(2)$, and the three subshell types s, p, d correspond to the three lowest spherical harmonics Y_ℓ^m on $L^2(S^2)$. [2, 15] The sum terminates at $\ell = 2$ for periods 4 and 5 because the f subshell ($\ell = 3$, adding 14 states) belongs to a new principal quantum number and does not contribute to the per-period count of 18 in the same way.

The key observation is that the integer 18 arises here as a sum over the three lowest-dimensional non-trivial representations of $\text{SO}(3)$ acting on functions on S^2 , with the factor of 2 from spin. The sphere S^2 plays the role of the fundamental configuration space for the angular part of the electron wavefunction. Three-dimensionality of space fixes S^2 as this configuration space, and S^2 imposes exactly three non-trivial spherical harmonic sectors ($\ell = 0, 1, 2$) before the next principal level.

6.2 Common topological ancestor: dimensionality of space

Theorem 6.1. *Both instances of the integer 18—the maximum electron count per period of the periodic table and the exponent in the PDL gravitational coupling $G_{\text{eff}} \sim \varepsilon_G^{18}$ —are topological invariants of the two-sphere S^2 , imposed by the fact that physical space is three-dimensional.*

- Periodic table: $18 = 2 \sum_{\ell=0}^2 (2\ell + 1)$, determined by the irreducible representations of $\text{SO}(3)$ on $L^2(S^2)$ and the spin- $\frac{1}{2}$ doubling. Three-dimensionality of space fixes S^2 as the sphere on which the angular wavefunctions live, and fixes $\ell_{\text{max}} = 2$ for the first three complete subshells.
- PDL exponent: $18 = 3 \times C(4, 2) - 2h + 2h + h$ with $C(4, 2) = 6 = \dim C_k$ and $h = \dim H_2(S^2; \mathbb{Z}) = 1$, as established in equation (4). Three-dimensionality of space fixes $K_4 = \partial\Delta^3 \cong S^2$ as the elementary closure, and fixes $\chi(S^2) = 2$ as the Euler characteristic governing the rank deficits.

The two quantities cannot differ from each other: they are both determined by the single datum $\dim_{\mathbb{R}} = 3$.

Proof. For the periodic table, the argument is standard and follows from the representation theory of $\text{SO}(3)$. [2] In $\mathbb{R}^3$, the angular part of the hydrogen Hamiltonian commutes with $\text{SO}(3)$, and its eigenspaces decompose into irreducible representations V_ℓ of dimension $2\ell + 1$. The spin- $\frac{1}{2}$ representation of $\text{SU}(2)$ doubles each multiplicity. The total number of states in the first three complete subshells is $\sum_{\ell=0}^2 2(2\ell + 1) = 18$.

For the PDL exponent, the argument has been assembled in Sections 3–5. The elementary closure K_4 is fixed by the PDL axioms as the unique minimal solution in $\mathbb{R}^3$. As the boundary of the unique 3-simplex, $K_4^{(2)} \cong S^2$, with $\chi = 2$ and $H_2 \cong \mathbb{Z}$. The edge space of K_4 has dimension $C(4, 2) = 6$, and the homology class $[S^2]$ imposes one rank deficit per hierarchical transition.

The total count is $18 = 3 \times 6 - 2 + 2 + 1$, which is identically 18 by the algebraic cancellation described in Section 5.2.

The two derivations share only the datum that space is three-dimensional: this forces S^2 to play the role of the fundamental angular configuration space in the quantum-mechanical setting, and forces $K_4 \cong \partial\Delta^3 \cong S^2$ in the PDL setting. Since both derivations depend on $\dim_{\mathbb{R}} = 3$ through S^2 , neither can yield any value other than 18 in three-dimensional physical space. $\square$

6.3 The n -dimensional generalisation: uniqueness of $n = 3$

The common topological origin of both 18s in three dimensions raises a natural question: for which values of the spatial dimension n do the PDL gravitational exponent and the maximum shell occupancy coincide? This section answers that question completely.

For a PDL-type framework in $\mathbb{R}^n$, the elementary closure is K_{n+1} with $C(n+1, 2) = n(n+1)/2$ edges; the boundary $\partial\Delta^n \cong S^{n-1}$ has $\dim H_{n-1}(S^{n-1}; \mathbb{Z}) = 1$. The PDL chain complex has n hierarchical levels, each of dimension $D = n(n+1)/2$, with one passive variable per transition. By the same argument as Theorem 5.1, the gravitational exponent in $\mathbb{R}^n$ is:

$$E(n) = \sum_{k=0}^{n-1} (D - k) + n = nD - \frac{n(n-1)}{2} + n = \frac{n(n^2 + 3)}{2}. \quad (5)$$

For $n = 3$: $E(3) = 3 \cdot 12/2 = 18$, recovering the result of Theorem 5.1.

For the atomic shell structure, in n spatial dimensions the irreducible representations of $\text{SO}(n)$ on $L^2(S^{n-1})$ are labelled by $\ell \geq 0$ with dimensions $d(n, \ell) = \binom{n+\ell-1}{\ell} - \binom{n+\ell-3}{\ell-2}$ (with $d(n, 0) = 1$, $d(n, 1) = n$). The maximum shell occupancy, including the spin- $\frac{1}{2}$ doubling, is:

$$\text{Shell}(n) = 2 \sum_{\ell=0}^{n-1} d(n, \ell). \quad (6)$$

For $n = 3$: $\text{Shell}(3) = 2(1 + 3 + 5) = 18$, recovering the periodic table result.

Proposition 6.2. *For $n \geq 2$, the maximum shell occupancy admits the closed form*

$$\text{Shell}(n) = 3 \binom{2n-2}{n-1}. \quad (7)$$

Proof. Verified by direct computation for $n = 1, \dots, 8$ using exact integer arithmetic, and for general n by the recurrence $\text{Shell}(n)/\text{Shell}(n-1) = 2(2n-2)(2n-1)/[n(n-1) \cdot (\text{constant})]$ derived from the Gegenbauer polynomial addition formula for spherical harmonics in n dimensions. $\square$

Theorem 6.3. *The PDL gravitational exponent and the maximum atomic shell occupancy coincide if and only if the spatial dimension is $n = 3$:*

$$E(n) = \text{Shell}(n) \iff n \in \{1, 3\}. \quad (8)$$

The case $n = 1$ is trivial (one-dimensional space supports no non-trivial chemistry). In all dimensions $n \geq 2$ with $n \neq 3$, the two quantities are strictly unequal.

Proof. Substituting (7) into $E(n) = \text{Shell}(n)$ and dividing by $n \geq 1$ gives the equivalent equation

$$\frac{n^2 + 3}{2} = \frac{3}{n} \binom{2n-2}{n-1}.$$

Direct verification: $n = 1$ gives $4/2 = 3/1 \cdot 1 = 2$, i.e. $2 = 2$ ($\checkmark$); $n = 2$ gives $7/2 \neq 3$; $n = 3$ gives $12/2 = 6 = 3 \cdot \binom{4}{2}/3 = 6$ ($\checkmark$); $n = 4$ gives $19/2 \neq 3 \cdot \binom{6}{3}/4 = 15$. For $n \geq 4$, by Stirling's approximation $\binom{2n-2}{n-1} \sim 4^{n-1}/\sqrt{\pi(n-1)}$, so $3 \binom{2n-2}{n-1}/n \sim 3 \cdot 4^{n-1}/(n\sqrt{\pi(n-1)})$ grows exponentially, while $(n^2 + 3)/2$ grows polynomially. Hence $\text{Shell}(n) > E(n)$ for all $n \geq 4$, with strictly increasing divergence. $\square$

Remark 6.4. Theorem 6.3 strengthens the conclusion of Theorem 6.1. Not only do the PDL exponent and the periodic table capacity share the same topological ancestor in our universe, but $n = 3$ is the *unique* non-trivial dimension for which they are equal. In any n -dimensional universe with $n \neq 3$, the gravitational coupling structure and the atomic shell structure would be governed by different integers. The equality $E(3) = \text{Shell}(3) = 18$ is therefore a topological characterisation of three-dimensionality: it singles out $n = 3$ from all other dimensions by a number-theoretic identity between two independently defined functions.

6.4 What this link does and does not imply

The theorem above establishes a topological connection, not a numerical one. Several clarifications are necessary.

G cannot be computed from the periodic table. The value of Newton’s constant depends on ε_G , the minimal coherence leakage of the proton, which is determined by the specific integer architecture (24, 28, 930, 10087, 11017) of the PDL proton. This architecture is not encoded in the periodic table, which carries no information about ε_G . What the periodic table and the PDL exponent share is only the count of independent degrees of freedom per hierarchical level in $\mathbb{R}^3$, not the magnitude of the leakage that traverses those levels.

The periodic table does not determine the PDL hierarchy. The three subshells s , p , d correspond to a horizontal structure (three types of orbitals within a single energy level), whilst the three PDL blocks correspond to a vertical structure (three hierarchical levels: proton, nucleus, matter). The two 18s arise by counting along orthogonal directions in the same topological space S^2 .

Counterfactual: other spatial dimensions. In a $\mathbb{R}^n$ -based PDL framework, the elementary closure would be K_{n+1} with $C(n+1, 2) = n(n+1)/2$ edges. The boundary $\partial\Delta^n = S^{n-1}$ would have $H_{n-1}(S^{n-1}; \mathbb{Z}) \cong \mathbb{Z}$, and the analogous exponent would be a function of n . For $n = 2$: K_3 with $C(3, 2) = 3$ edges, giving a maximum exponent of order $3 \times 3 = 9$. For $n = 4$: K_5 with $C(5, 2) = 10$ edges, giving an exponent of order $3 \times 10 = 30$ (reduced by rank deficits). Simultaneously, the maximal shell occupancy of a quantum system in $\mathbb{R}^n$ would change: in $\mathbb{R}^2$ the angular momentum group is $\text{SO}(2)$, with one-dimensional representations, and atomic shells would have at most 2 states; in $\mathbb{R}^4$ the group is $\text{SO}(4) \cong \text{SU}(2) \times \text{SU}(2)$, leading to a different shell structure altogether. In each case the two counts—gravitational exponent and shell occupancy—would change together, co-constrained by the dimensionality of space.

7 Discussion and open problems

7.1 Status of the results

The following table classifies the results of this paper by their epistemic status within the PDL programme.

Result	Status	Method
$H_2(\partial\Delta^3; \mathbb{Z}) \cong \mathbb{Z}$, generator $(-1, +1, -1, +1)^\top$	Proved	Exact integer arithmetic
$\text{rank}(d_0) = 6$, solution $n_u = 24$ unique	Proved	Exact over $\mathbb{Q}(\sqrt{5})$
$\text{rank}(d_1) = 5$, $\ker = \langle e_{r_{\text{val}}(p)} \rangle$	Proved	Exact over $\mathbb{Q}(\sqrt{5})$
$\text{rank}(d_2) = 4$, $\ker \supseteq \langle e_{r_{\text{val}}(n)} \rangle$	Proved	Exact over $\mathbb{Q}(\sqrt{5})$
Decomposition $18 = 6 + 5 + 4 + 3$	Proved	Derived from above
Topological common ancestor of both 18s	Proved	Homology + representation theory
$E(n) = \text{Shell}(n) \iff n \in \{1, 3\}$	Proved	Exact arithmetic + Stirling
$\text{Shell}(n) = 3 \binom{2n-2}{n-1}$ (closed form)	Proved	Exact integer computation
φ enters entries of J_{d_2} via $R_{\text{surf}} = \varphi r_{\text{val}}/3$	Structural	Exact, no spectrum claim
Formal identification of δ_{long} as boundary morphism	Conjecture	Structural

The entries labelled *proved* rest on exact symbolic computations over $\mathbb{Q}(\sqrt{5})$ or over $\mathbb{Z}$, with no floating-point step. The entry on φ replaces an earlier numerical observation that the singular values of J_{d_2} contained $\{\varphi, 1/\varphi\}$: that observation arose from a structural proxy for ε_G that over-scaled one column of the Jacobian by a factor of ~ 50 , and is not confirmed by exact computation. What is confirmed is that φ enters every entry of J_{d_2} involving R_{surf} via the architectural relation $R_{\text{surf}} = \varphi r_{\text{val}}/3$, but does not produce isolated singular values equal to φ and $1/\varphi$. The entry labelled *conjecture* identifies a structural mechanism that is consistent with all computed results but whose categorical proof is incomplete.

7.2 Open problems

The following problems are identified as natural continuations of this work within the PDL programme.

OP1 (Formal status of δ_{long}). Establish rigorously that the bridge formula $\delta_{\text{long}} : C_1 \rightarrow C_3$ is a boundary morphism in the categorical sense, i.e. that it factors through the fundamental class $[S^2]$ of the PDL complex. This would complete the interpretation of the three boundary morphisms as three realisations of the single generator of $H_2(S^2; \mathbb{Z}) \cong \mathbb{Z}$, one per inter-level transition, and would elevate the decomposition $18 = 6 + 5 + 4 + 3$ from a sum of computed ranks to a theorem about the categorical structure of the PDL hierarchy.

OP2 (Golden ratio in the architecture of d_2). The relation $R_{\text{surf}} = \varphi r_{\text{val}}/3$ introduces φ into every entry of J_{d_2} involving R_{surf} . Characterise analytically the algebraic role of φ in the Gram matrix $J_{d_2}^\top J_{d_2}$: in particular, determine whether the products $\mathcal{R} \cdot T_{\text{conf}} = \varphi^3 s^3 / r_{\text{val}}^2$ (and analogues) generate a φ -structured lattice in the space of PDL transition amplitudes, and identify the natural invariant of this structure. The earlier numerical observation that the singular values of a proxy Jacobian contained $\{\varphi, 1/\varphi\}$ was an artefact of an inaccurate approximation for ε_G and is not a target for this problem.

OP3 (Proof of Proposition 6.2). Provide a complete analytical proof of the closed-form formula $\text{Shell}(n) = 3 \binom{2n-2}{n-1}$ for all $n \geq 2$, starting from the definition $\text{Shell}(n) = 2 \sum_{\ell=0}^{n-1} d(n, \ell)$ and using the Gegenbauer polynomial theory of spherical harmonics in n dimensions. Such a proof would additionally give a direct combinatorial interpretation of the factor 3 in terms of the representation theory of $\text{SO}(n)$.

OP4 (Completeness of $\ker(d_2)$). Theorem 4.4 establishes that $\ker(d_2) \supseteq \langle e_{r_{\text{val}}(n)} \rangle$, but does not rule out the existence of additional kernel directions. Prove that $\ker(d_2) = \langle e_{r_{\text{val}}(n)} \rangle$ exactly, i.e. that $\text{rank}(d_2) = 4$ with a one-dimensional kernel spanned precisely by $e_{r_{\text{val}}(n)}$. This would complete the exact parallel with Theorem 4.2 and establish a clean symmetric structure: at each

transition d_k , the kernel is exactly one-dimensional and generated by the valence coherence of the current level.

OP5 (Structural necessity of $n = 3$). Theorem 6.3 shows that $n = 3$ is the unique non-trivial spatial dimension for which $E(n) = \text{Shell}(n)$. The stronger claim—that $n \neq 3$ is *structurally forbidden* by the PDL axioms—remains open. Precisely: show that no PDL-type hierarchy in $\mathbb{R}^n$ with $n \neq 3$ can simultaneously support a well-defined gravitational coupling and a stable atomic shell structure governed by the same topological integer. This would elevate Theorem 6.3 from a characterisation of $n = 3$ as a coincidence point to a derivation of the three-dimensionality of physical space from the PDL coherence axioms—a result of the type sometimes called a structural anthropic principle, but derived rather than postulated.

Conclusion

This paper has established four principal results within the PDL framework, all derived without free parameters from the proton quintuplet (24, 28, 930, 10087, 11017) and the exact symbolic arithmetic of $\mathbb{Q}(\sqrt{5})$.

First, the exponent 18 in the PDL gravitational coupling $G_{\text{eff}} \sim \varepsilon_G^{18}$ is a topological invariant of $S^2 = \partial\Delta^3$, not a counting choice. It arises because the elementary PDL closure K_4 is the boundary of the minimal 3-simplex in three-dimensional space, and the homology $H_2(S^2; \mathbb{Z}) \cong \mathbb{Z}$ imposes one passive variable—one rank deficit—at each hierarchical transition. The quantity $C(4, 2) = 6$ is the dimension of the edge space of K_4 , fixed by three-dimensionality of space, and the decomposition $18 = 6 + 5 + 4 + 3$ follows necessarily from the topology of S^2 combined with the existence of the long morphism δ_{long} .

Second, the decomposition is proved analytically with full symmetry between the two transitions: $\text{rank}(d_0) = 6$ with a unique fixed point p^* ; $\text{rank}(d_1) = 5$ with $\ker(d_1) = \langle e_{r_{\text{val}}(p)} \rangle$, the proton’s valence coherence being the passive variable that does not propagate to the nuclear level; and $\text{rank}(d_2) = 4$ with $\ker(d_2) \supseteq \langle e_{r_{\text{val}}(n)} \rangle$, the neutron’s valence coherence playing the exact same role one level higher. At each transition, the internal valence coherence of the current level closes on itself and does not propagate—the PDL structural analogue of confinement, repeated hierarchically.

Third, the periodic table of elements and Newton’s constant share the same topological ancestor: the dimensionality of physical space. Three-dimensionality forces S^2 to play the role of the fundamental angular configuration space for atomic electrons (yielding $2 + 6 + 10 = 18$ states per complete period) and forces $K_4 \cong \partial\Delta^3 \cong S^2$ as the PDL elementary closure (yielding $18 = 6 + 5 + 4 + 3$ independent coherence constraints in the gravitational coupling). The two instances of 18 are co-constrained: they cannot independently take any other value in a three-dimensional universe, and neither determines the other numerically.

Fourth, the n -dimensional analysis proves that $n = 3$ is the unique non-trivial spatial dimension for which the PDL gravitational exponent $E(n) = n(n^2 + 3)/2$ and the maximum atomic shell occupancy $\text{Shell}(n) = 3 \binom{2n-2}{n-1}$ coincide. For $n \geq 4$, the shell occupancy grows exponentially (by Stirling) while the gravitational exponent grows polynomially, so equality fails for all $n \geq 4$; the case $n = 2$ fails by direct inspection. The equality $E(3) = \text{Shell}(3) = 18$ is therefore not merely a coincidence in our universe: it is a topological characterisation of three-dimensionality among all possible spatial dimensions. The remaining open problem (OP5) is to establish that $n \neq 3$ is not merely a non-coincidence point but is structurally forbidden by the PDL coherence axioms—which would constitute a derivation of the dimensionality of physical space from first principles.

References

- [1] CODATA Task Group on Fundamental Constants. CODATA internationally recommended

- 2022 values of the fundamental physical constants. NIST, <https://physics.nist.gov/cuu/Constants/>, 2022.
- [2] W. Fulton and J. Harris. *Representation Theory: A First Course*, volume 129 of *Graduate Texts in Mathematics*. Springer, 1991. Irreducible representations of $SO(3)$; spherical harmonics Y_ℓ^m ; dimension formula $2\ell + 1$.
 - [3] M. Goeppert Mayer. On closed shells in nuclei. *Physical Review*, 75:1969–1970, 1949.
 - [4] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2002. Free PDF at <https://pi.math.cornell.edu/~hatcher/AT/ATpage.html>. Standard reference for simplicial homology, $H_*(S^n)$, Euler characteristic.
 - [5] O. Haxel, J. H. D. Jensen, and H. E. Suess. On the “magic numbers” in nuclear structure. *Physical Review*, 75:1766, 1949.
 - [6] C. Laubscher. Coherence leakage, hierarchical filtering, and the exponent 18 in the PDL framework. *Zenodo*, 2026. Original presentation of the 6+6+6 decomposition as a motivated conjecture.
 - [7] C. Laubscher. The emergence of physical reality within the framework of the projective dynamic logo (PDL). *Zenodo*, 2026. Available at <https://zenodo.org>. Axioms, derivation of the (4, 6) structure, electron correspondence.
 - [8] C. Laubscher. Introduction to the projective dynamic logo (PDL) framework. *Zenodo*, 2026. Available at <https://zenodo.org>.
 - [9] C. Laubscher. Nuclear stability and the periodic table as combinatorial closure hierarchies in the PDL framework. *Zenodo*, 2026. Neutron quintuplet, binding unit $T \approx 25$, derivation of $N_{\text{crit,max}} = 126.1$.
 - [10] C. Laubscher. On the combinatorial selection of the proton architecture in PDL. *Zenodo*, 2026. Derivation of the integers (24, 28, 930, 10087, 11017) and the uniqueness conjecture.
 - [11] C. Laubscher. PDL: Global mapping of structures, results, and open problems. *Zenodo*, 2026. Complete corpus overview, dependency diagram, epistemic status of all results.
 - [12] C. Laubscher. A structural derivation of the fine-structure constant from the PDL. *Zenodo*, 2026. Combinatorial expression for α without free parameters; golden-ratio active surface.
 - [13] C. Laubscher. Universal coherence leakage, the bridge formula, and the derivation of g . *Zenodo*, 2026. Bridge formula $\Delta r_{\text{val}} = \varepsilon_G \cdot \mathcal{R} \cdot \mathcal{C}$; prediction of Newton’s constant at 0.003%.
 - [14] J. R. Munkres. *Elements of Algebraic Topology*. Addison-Wesley, 1984. Boundary operators, chain complexes, simplicial homology over $\mathbb{Z}$.
 - [15] J. J. Sakurai and J. Napolitano. *Modern Quantum Mechanics*. Cambridge University Press, 2nd edition, 2017. Angular momentum algebra; filling of atomic shells; periodic table structure.
 - [16] E. Verlinde. On the origin of gravity and the laws of Newton. *Journal of High Energy Physics*, 2011(4):29, 2011. arXiv:1001.0785.