

Model specification / goodness of fit

Consider a generic multiple Gaussian linear model

$$Y = X\beta + \varepsilon \quad \varepsilon \sim N_n(0, \sigma^2 I_n)$$

Y vector of response variables

X $n \times p$ matrix of covariates

$\beta = [\beta_1 \beta_2 \dots \beta_p]^T$ vector of regression parameters

We have seen the statistical tests to evaluate the model's adequacy:

- Test about an individual coefficient

$$\begin{cases} H_0: \beta_j = 0 \\ H_1: \beta_j \neq 0 \end{cases}$$

If I do not reject $\beta_j = 0$ for some j , I can remove that covariate from the model specification and estimate a new one with one less covariate but the same accuracy at predicting y .

- Test about a subset of coefficients

$$\begin{cases} H_0: \beta_{k+1} = \beta_{k+2} = \dots = \beta_p = 0 \\ H_1: \text{at least one is } \neq 0 \end{cases}$$

similarly, if I do not reject H_0 , I can remove that subset of covariates without losing accuracy

- Test about the overall significance

$$\begin{cases} H_0: \beta_1 = \dots = \beta_p = 0 \\ H_1: \bar{H}_0 \end{cases}$$

in this case, the model is useless.

R^2 and R^2_{adj} (adjusted R^2)

We have seen how the coefficient R^2 describes the proportion of variability explained by the model. Hence, we could think of using R^2 to choose between different model specifications.

However, if I use R^2 to compare nested models (i.e. one can be obtained starting from the other through a set of constraints), R^2 is not a valid measure.

consider: (a) $\hat{y}_i = \hat{\beta}_1 + \hat{\beta}_2 x_i$

(b) $\tilde{y}_i = \tilde{\beta}_1 + \tilde{\beta}_2 x_i + \tilde{\beta}_3 w_i$ | add one covariate

$$R^2_{(a)} \leq R^2_{(b)} \quad \text{BY CONSTRUCTION!}$$

Recall that $R^2 = \frac{SSR}{SST}$

$$SST = \sum_{i=1}^n (y_i - \bar{y})^2 \quad \text{does not depend on the model} \Rightarrow SST_{(a)} = SST_{(b)}$$

$$\text{However, } SSR_{(b)} \geq SSR_{(a)}$$

the SSR of model (b) can not be smaller than $SSR_{(a)}$.

If w_i is useful to predict y , $SSR_{(b)} > SSR_{(a)}$

In the worst case (if w_i is really useless), I set $\tilde{\beta}_3 = 0$ and I obtain $SSR_{(b)} = SSR_{(a)}$.

The more variables I include in the model, the larger R^2 will be.

So we can not use it to compare, for example, models (a) and (b) — or, in general, NESTED MODELS.

In general:

MORE COVARIATES $\begin{cases} R^2 \text{ increases} \\ \text{less interpretable} \\ \text{overfit} \end{cases}$

FEWER COVARIATES $\begin{cases} \text{parsimony} \\ \text{interpretable} \end{cases}$

of course, we want few covariates, but not too few!

• ADJUSTED R^2 $R^2_{adj} = 1 - (1 - R^2) \cdot \frac{n-1}{n-p}$

it is "adjusted" for the model dim. p

penalizes models with many covariates.

when I introduce a new covariate:

— R^2 can remain the same or increase

R^2_{adj} can increase, remain the same, or decrease

\Downarrow
 R^2_{adj} can be < 0 !

\Rightarrow To compare nested models we can use R^2_{adj} .