



Investigating the effects of NO_x emission and chemistry on the diurnal variation of NO₂ over East Asia using ground-based and geostationary satellite observations

Laura Hyesung Yang

Daniel Jacob, Ruijun Dang, Yujin Oak, Haipeng Lin, Shixian Zhai, Drew Pendergrass, Ellie Beaudry, Nadia Colombi, Viral Shah, Xu Feng, Robert Yantosca, Heesung Chong, Junsung Park, Hanlim Lee, Won-Jin Lee, Soontae Kim, Eunhye Kim, Katherine Travis, James Crawford, Jhoon Kim

The 14th GEMS Workshop, September 7th, 2023



Differences in NO_x ($\equiv \text{NO} + \text{NO}_2$) lifetime between different seasons can result in distinct diurnal variations in NO_2 column

NO_x lifetimes and loss pathways




Figure from Shah et al. (2020)

** In the absence of transport **

$$\frac{dN_v}{dt} = \alpha(t)(E(t) - k(t)\frac{\text{NO}_2}{\text{NO}_x}\text{column ratio})$$


Trop. NO_2 column $\frac{\text{NO}_2}{\text{NO}_x}$ column ratio NO_x emission NO_x chemical loss

Equation from Boersma et al. (2008)

Laura Yang · laurayang@g.harvard.edu

A chemical transport model (CTM) can elucidate the factors driving diurnal variation in retrieved and measured NO_2

GEMS
 NO_2 column



Pandora
 NO_2 column



Surface
 NO_2 concentration



Drivers of
diurnal variation in NO_2


Emission? Chemistry?

GEOS
Chem

Diurnal variation in
simulated NO_2


The main difference in diurnal variation of NO_x emissions occurs between day and night, which impacts the morning behavior of NO_2

Diurnal Variation of NO_x Emissions in BTH



MEIC inventory diurnal scaling factor
(Miao et al. 2020)




Diurnal Variation of NO_x Emissions in the SMA







Developed KORUSv5 scaling factor based on TOPIS traffic count and construction activity time

NO_2 diurnal variation from surface measurements in summer (JJA) is affected by monsoon

Mixed layer growth complicates interpreting diurnal variation of NO_2 from surface measurements



Pandora NO₂ column measurements are sensitive to advection





Too few Pandora
measurements available
for JJA

GEMS and GEOS-Chem from one $0.25^\circ \times 0.3125^\circ$ grid box

Dot shows the location of the Pandora station

Extended NO_x lifetime and stronger wind in the winter increases effect of transport, complicating urban-scale diurnal analysis using GEMS



Trop. NO₂ column


$$\frac{dN_v}{dt} = \alpha(t)(E(t) - k(t)N_x(t) - \nabla \cdot N_x(t)U)$$

$\frac{\text{NO}_2}{\text{NO}_x}$ column ratio

NO_x emission NO_x chemical loss Transport





- Strong Northwesterly winds blow over the Seoul Metropolitan Area (SMA) during the winter (Seo et al. 2021)
- Over a small horizontal domain, the transport term becomes important
- Spatial averaging over large domain is needed to minimize the effect of the transport term

A wider spatial domain is required to reduce the impact of transport when interpreting the diurnal variation of NO₂





** DJF and JJA have different y-axis scale **

In the summer, emission regions exhibit an accumulation of emissions in the early morning, followed by a decrease driven by chemical processes



** BGI has a different y-axis scale

In the winter, the accumulation of high emissions lead to a net increase in the NO_2 column over emission regions



** Remote site has a different y-axis scale

Takeaways

During summer, the diurnal variation of the NO₂ column is driven by active photochemistry while in winter, it is driven by emission

Urban-scale analysis can be complicated by the transport term and must be done carefully especially during the winter over East Asia

GEOS-Chem agrees with GEMS observations and can provide an explanation for the observed diurnal variation in GEMS