

Assessment of Hydrogen's Climate Impact Is Affected by Model OH Biases



Laura Yang

D. Jacob, H. Lin, R. Dang, K. Bates, J. East, K. Travis,
D. Pendergrass, L. Murray




[Under review by GRL]

AGU 2024
December 9, 2024





Hydrogen (H₂) is an indirect climate forcer due to its atmospheric oxidation by OH



Future source: H₂ leakage
in a hydrogen economy


Deposition
to soils



H₂ total lifetime: 2.4 ± 0.3 years

H ₂ GWP-100	Model	References
8 ± 2	2-D model	Derwent et al. (2023)
10^{+7}_{-4}	Box model	Chen et al. (2024)
11.6 ± 2.8	Ensemble of five 3-D models	Sand et al. (2023)

Global 3-D models overestimate tropospheric OH by 10-30% as inferred by the methyl chloroform (MCF) proxy






The OH biases in current models may lead to biases in the GWP estimates

¹: Prather et al. (2012), ²: Sand et al. (2023), ³: Hauglustaine et al. (2022),

⁴: Warwick et al. (2023), ⁵: Naik et al. (2013)

Underestimated OH reactivity (OHR) in the models may be driving an overestimation of OH and H₂ GWP



OH reactivity (OHR): OH loss frequency

$$\text{OHR} = k_{\text{H}_2+\text{OH}}[\text{H}_2] + \sum k_{\text{VOC}+\text{OH}}[\text{VOC}] + \dots$$

Missing/
under-accounted
species






OHR
underestimate

Overestimate of
H₂ GWP

↑ importance of H₂
for affecting
 τ_{CH_4} & P_{O_3}





τ_{CH_4} : Lifetime of methane
 P_{O_3} : O₃ production

Modified GEOS-Chem with increased OHR makes H_2 less of a driver for OH loss and relaxes CH_4 - H_2 coupling



- 1) Added volatile chemical product^{1,2} & oceanic³ emissions of NMVOCs
 - 2) Added terminal OH sink over continents
- * *The change in O_3 concentration is minor*

Modified GEOS-chem shows improved agreement with observationally constrained OH values



¹: Prather et al. (2012), ²: Sand et al. (2023), ³: Hauglustaine et al. (2022), ⁶

⁴: Warwick et al. (2023), ⁵: Naik et al. (2013)

The roles that H₂ and CH₄ play in ozone production decrease in the modified GEOS-Chem

	Standard GEOS-Chem	Modified GEOS-Chem
Chemical source (Tg O₃ equivalent a⁻¹)		
NO + HO ₂ → NO ₂ + OH	3970	3590
NO + CH ₃ O ₂ → NO ₂ + CH ₃ O	1420	1320
NO + RO ₂ → NO ₂ + RO	330	900
Total	5720	5810
Chemical loss (Tg O₃ equivalent a⁻¹)		
O(¹ D) + H ₂ O → 2OH	2170	2290
O ₃ + HO ₂ → OH + 2O ₂	1340	1360
O ₃ + OH → HO ₂ + O ₂	660	500
Others	1050	1140
Total	5220	5290
Lifetime against chemical loss (days)		
	24.2	24.8
Tropospheric O₃ burden (Tg)		
	346	359

Modified GEOS-Chem

Increased contribution of RO₂ to P_{O₃}
due to added NMVOCs




Decreased contributions of H₂ and CH₄
to P_{O₃}

O₃ concentrations in the modified model
show little change from the standard model

RO₂: Organic peroxy radicals
NMVOCs: Non-methane VOCs
P_{O₃}: O₃ production

Correcting OHR and OH biases in modified GEOS-Chem leads to 20% smaller hydrogen GWP-100



- GWP-100 calculation method, soil sink, and stratospheric H₂O follow Sand et al. (2023)
- Increasing OHR to match observations decreases the GWP regardless of how this OHR increase is implemented

Takeaways

Correcting the OHR underestimate in global 3-D models decreases the coupling between CH_4 and H_2 and reduces the effect of H_2 on O_3

OH and OHR biases may lead to a 20% overestimation of H_2 GWP-100 in current 3-D atmospheric chemistry models

Better understanding of the factors controlling global OH concentrations and OHR is needed for GWP estimates



Acknowledgements. The authors acknowledge funding support from the ExxonMobil Technology & Engineering Company, the NASA ACCDAM Program, and the NSF GRFP.

We thank B. Mignone, E. Reidy, O. Clifton, R. Skeie, L. Mickley, and T. Mooring for helpful discussions. The views expressed in this work are solely those of the authors.