


# Interpreting GEMS geostationary satellite observations of the diurnal variation of nitrogen dioxide (NO<sub>2</sub>) over East Asia

Laura H. Yang<sup>1</sup>, Daniel J. Jacob<sup>1,2</sup>, Ruijun Dang<sup>1</sup>, Yujin J. Oak<sup>1</sup>, Haipeng Lin<sup>1</sup>, Jhoon Kim<sup>3</sup>, Shixian Zhai<sup>4</sup>, Nadia K. Colombi<sup>2</sup>, Drew C. Pendergrass<sup>1</sup>, Ellie Beaudry<sup>1</sup>, Viral Shah<sup>5,6</sup>, Xu Feng<sup>1</sup>, Robert M. Yantosca<sup>1</sup>, Heesung Chong<sup>7</sup>, Junsung Park<sup>7</sup>, Hanlim Lee<sup>8</sup>, Won-Jin Lee<sup>9</sup>, Soontae Kim<sup>10</sup>, Eunhye Kim<sup>10</sup>, Katherine R. Travis<sup>11</sup>, James H. Crawford<sup>11</sup>, Hong Liao<sup>12</sup>


<sup>1</sup> Harvard University, John A. Paulson School of Engineering and Applied Sciences <sup>2</sup> Harvard University, Department of Earth and Planetary Sciences <sup>3</sup> Yonsei University, Department of Atmospheric Sciences <sup>4</sup> The Chinese University of Hong Kong, Earth and Environmental Sciences Programme and Graduation Division of Earth and Atmospheric Sciences <sup>5</sup> Global Modeling and Assimilation Office, NASA Goddard Space Flight Center <sup>6</sup> Science Systems and Applications, Inc. <sup>7</sup> Center for Astrophysics | Harvard & Smithsonian <sup>8</sup> Pukyong National University, Division of Earth Environmental System Science <sup>9</sup> Environmental Satellite Center, National Institute of Environmental Research <sup>10</sup> Ajou University, Department of Environmental and Safety Engineering <sup>11</sup> NASA Langley Research Center <sup>12</sup> School of Environmental Science and Engineering, Nanjing University of Information Science and Technology




## Introduction

- Nitrogen oxide radicals (NO<sub>x</sub> ≡ NO + NO<sub>2</sub>) emitted by fuel combustion are important precursors of ozone and particulate matter pollution.
- The GEMS, launched in space in 2020, now provides hourly daytime observations of NO<sub>2</sub> columns over East Asia.
- Here we investigate the drivers of the diurnal variation of NO<sub>2</sub> observed by GEMS during winter and summer over Beijing and Seoul.

## Model & Observations




## Diurnal variation of NO<sub>x</sub> emissions




- Emissions are dominated by industrial and transport sources in Beijing, and by mobile sources in Seoul.
- Both sectors show a broad maximum between 7 and 18 LT.
- There is no significant rush hour peaks in transport emissions, suggesting that surface NO<sub>2</sub> maxima observed in rush hours are driven more by shallow mixing depth.

## Diurnal variation of NO<sub>2</sub> over urban core




- NO<sub>2</sub> in winter increases throughout the day due to high daytime emissions and increasing NO<sub>2</sub>/NO<sub>x</sub> ratio from entrainment of ozone, partly balanced by loss from transport.
- NO<sub>2</sub> in summer, chemical loss combined with transport drives a minimum in the NO<sub>2</sub> column at 13-14 local time.

## Segregation by wind speed in winter



- At high wind speed, emission is balanced by transport leading to weak diurnal variation.
- At low wind speed, NO<sub>2</sub> accumulates as transport cannot keep up with emission.

## Diurnal variation of NO<sub>2</sub> at a regional scale



- In winter, NO<sub>2</sub> shows a steady increase over the daytime hours due to emissions, but the transport term remains the major sink of NO<sub>x</sub>.
- In summer, NO<sub>2</sub> shows negligible loss from transport in daytime because chemical loss term is much faster, but the diurnal amplitude is weak because of diluted emission and long residence time.

## Conclusions

- We can independently infer the emission in the winter and chemistry in the summer by quantifying the transport term using an inversion with a chemical transport model.
- Our conclusions regarding the interpretation of the diurnal variation of NO<sub>2</sub> columns can be extended to other instruments of geostationary air quality constellation like TEMPO and Sentinel-4.

## Acknowledgement

- This work was funded by the Samsung Advanced Institute of Technology and the Nanjing University of Information Science and Technology. LHY was funded by a US National Science Foundation Graduate Research Fellowship Program (NSF GRFP).

- Contact LHY at [aurayang@g.harvard.edu](mailto:aurayang@g.harvard.edu)
- Read the preprint at EGUsphere: <https://doi.org/10.5194/egusphere-2023-2979>

