

Materials 31 - Parameterized expectations against value iteration

Laura Gáti

May 23, 2020

The value function problem takes the form

$$V(\mathbf{x}_t, t) = \max_{i_t} - \left\{ \Omega_6 i_t^2 + \Omega_7 \bar{\pi}_{t-1}^2 + \Omega_8 i_t \bar{\pi}_{t-1} + \Omega_9 i_t s_t + \Omega_{10} \bar{\pi}_{t-1} s_t + \Omega_{11} s_t + \beta \mathbb{E}_t V(\mathbf{x}_{t+1}, t+1) \right\} \quad (1)$$


s.t.

$$\bar{\pi}_t = \bar{\pi}_{t-1} + k_t^{-1} (-\kappa \sigma i_t + \Omega_{12} \bar{\pi}_{t-1} + \Omega_5 s_t - b_1 s_{t-1}) \quad (2)$$

$$k_t^{-1} = \rho_k k_{t-1}^{-1} + \gamma_k (-\kappa \sigma i_t + \Omega_{12} \bar{\pi}_{t-1} + \Omega_5 s_t - b_1 s_{t-1})^2 \quad (3)$$

where the parameters Ω_i are defined in Materials 25, as is $\mathbf{x}_t \equiv (k_t^{-1}, \bar{\pi}_{t-1}, r_t^n, u_t)$.

Figure 1: Parameterized expectations solution



A Model summary

$$x_t = -\sigma i_t + \hat{\mathbb{E}}_t \sum_{T=t}^{\infty} \beta^{T-t} ((1-\beta)x_{T+1} - \sigma(\beta i_{T+1} - \pi_{T+1}) + \sigma r_T^n) \quad (\text{A.1})$$

$$\pi_t = \kappa x_t + \hat{\mathbb{E}}_t \sum_{T=t}^{\infty} (\alpha\beta)^{T-t} (\kappa\alpha\beta x_{T+1} + (1-\alpha)\beta\pi_{T+1} + u_T) \quad (\text{A.2})$$

$$i_t = \psi_\pi \pi_t + \psi_x x_t + \bar{i}_t \quad (\text{if imposed}) \quad (\text{A.3})$$

$$\text{PLM:} \quad \hat{\mathbb{E}}_t z_{t+h} = a_{t-1} + b h_x^{h-1} s_t \quad \forall h \geq 1 \quad b = g_x h_x \quad (\text{A.4})$$

$$\text{Updating:} \quad a_t = a_{t-1} + k_t^{-1} (z_t - (a_{t-1} + b s_{t-1})) \quad (\text{A.5})$$

$$\text{Anchoring function:} \quad k_t = k_{t-1} + \mathbf{g}(f e_{t-1}^2) \quad (\text{A.6})$$

$$\text{Forecast error:} \quad f e_{t-1} = z_t - (a_{t-1} + b s_{t-1}) \quad (\text{A.7})$$

$$\text{LH expectations:} \quad f_a(t) = \frac{1}{1-\alpha\beta} a_{t-1} + b(\mathbb{I}_{nx} - \alpha\beta h)^{-1} s_t \quad f_b(t) = \frac{1}{1-\beta} a_{t-1} + b(\mathbb{I}_{nx} - \beta h)^{-1} s_t \quad (\text{A.8})$$

This notation captures vector learning (z learned) for intercept only. For scalar learning, $a_t = (\bar{\pi}_t \ 0 \ 0)'$ and b_1 designates the first row of b . The observables (π, x) are determined as:

$$x_t = -\sigma i_t + \begin{bmatrix} \sigma & 1-\beta & -\sigma\beta \end{bmatrix} f_b + \sigma \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} (\mathbb{I}_{nx} - \beta h_x)^{-1} s_t \quad (\text{A.9})$$

$$\pi_t = \kappa x_t + \begin{bmatrix} (1-\alpha)\beta & \kappa\alpha\beta & 0 \end{bmatrix} f_a + \begin{bmatrix} 0 & 0 & 1 \end{bmatrix} (\mathbb{I}_{nx} - \alpha\beta h_x)^{-1} s_t \quad (\text{A.10})$$

B Target criterion

The target criterion in the simplified model (scalar learning of inflation intercept only, $k_t^{-1} = \mathbf{g}(f e_{t-1})$):

$$\pi_t = -\frac{\lambda_x}{\kappa} \left\{ x_t - \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_t^{-1} + ((\pi_t - \bar{\pi}_{t-1} - b_1 s_{t-1})) \mathbf{g}_\pi(t) \right) \right. \\ \left. \left(\mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1} (1 - k_{t+1+j}^{-1} - (\pi_{t+1+j} - \bar{\pi}_{t+j} - b_1 s_{t+j}) \mathbf{g}_{\bar{\pi}}(t+j)) \right) \right\} \quad (\text{B.1})$$

where I'm using the notation that $\prod_{j=0}^0 \equiv 1$. For interpretation purposes, let me rewrite this as follows:

$$\pi_t = -\frac{\lambda_x}{\kappa} x_t + \frac{\lambda_x}{\kappa} \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_t^{-1} + f e_{t|t-1}^{eve} \mathbf{g}_\pi(t) \right) \mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i} \\ - \frac{\lambda_x}{\kappa} \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_t^{-1} + f e_{t|t-1}^{eve} \mathbf{g}_\pi(t) \right) \left(\mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1} (k_{t+1+j}^{-1} + f e_{t+1+j|t+j}^{eve} \mathbf{g}_{\bar{\pi}}(t+j)) \right) \quad (\text{B.2})$$

Interpretation: **tradeoffs from discretion in RE** + **effect of current level and change of the gain on future tradeoffs** + **effect of future expected levels and changes of the gain on future tradeoffs**