Materials 37 - Cross-section, neighborhood of zero forecast errors

Laura Gáti

July 14, 2020

Overview

1 Simulated "true" data

1.1 Odd number of breakpoints (5)

Figure 1: Mean estimates for increasing N, imposing convexity with weight 10M, truth with $nfe = 5, fe \in (-2, 2)$



(a) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=1000



(b) Autocovariogram



(c) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=2000



(d) Autocovariogram



(e) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=10000



(f) Autocovariogram

1.2 Even number of breakpoints (6)

Figure 2: Mean estimates for increasing N, imposing convexity with weight 100K, truth with $nfe = 6, fe \in (-3.5, 3.5)$



(a) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=100



(b) Autocovariogram



(c) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=1000



(d) Autocovariogram



(e) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, N=10000



(f) Autocovariogram

1.3 Finer grid in the zero neighborhood

Figure 3: Mean estimates for $N = 100$, imposing convexity with weight 100K, truth with $nfe = 5, fe \in (-2, 2)$



(a) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, 5 breakpoints, uniformly spaced



(b) Autocovariogram



(c) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, 7 breakpoints, uniformly spaced



(d) Autocovariogram



(e) $\alpha^{true}, \alpha_0, mean(\hat{\alpha})$, 5 breakpoints, concentrated around 0



(f) Autocovariogram

2 Autocovariogram for real data

Figure 4: Mean estimated parameters over the cross-section of size N, convexity imposed, mean moment not imposed



(a) $mean(\hat{\alpha}) = (0.0979; 0.0076; 0.0058; 0.006; 0.0864)$, N=1000, convexity weight 10M



(b) Autocovariogram



(c) $mean(\hat{\alpha}) = (0.0773; 0.0103; 0.0097; 0.0104; 0.0771)$, N=1000, convexity weight 100K



(d) Autocovariogram

3 Impulse responses to iid monpol shocks across a wide range of learning models

$T = 400, N = 100, n_{drop} = 5$, shock imposed at $t = 25$, calibration as above, Taylor rule assumed to be known, PLM = learn constant only, of inflation only.

Figure 5: IRFs and gain history (sample means)



(a) Decreasing gain learning



(b) Mean gain



(c) Constant gain learning



(d) Mean gain



(e) CEMP criterion (vector)



(f) Mean gain

Figure 6: IRFs and gain history (sample means), continued



(a) CUSUM criterion (vector)



(b) Mean gain



(c) Smooth criterion, approximated, using $\alpha^{true} = (0.05; 0.025; 0; 0.025; 0.05)$, on $fe \in (-2, 2)$.



(d) Mean gain

A Model summary

$$x_t = -\sigma i_t + \hat{\mathbb{E}}_t \sum_{T=t}^{\infty} \beta^{T-t}\big((1-\beta)x_{T+1} - \sigma(\beta i_{T+1} - \pi_{T+1}) + \sigma r_T^n\big) \tag{A.1}$$

$$\pi_t = \kappa x_t + \hat{\mathbb{E}}_t \sum_{T=t}^{\infty} (\alpha\beta)^{T-t}\big(\kappa\alpha\beta x_{T+1} + (1-\alpha)\beta\pi_{T+1} + u_T\big) \tag{A.2}$$

$$i_t = \psi_\pi \pi_t + \psi_x x_t + \bar{i}_t \qquad \text{(if imposed)} \tag{A.3}$$

$$\text{PLM:} \qquad \hat{\mathbb{E}}_t z_{t+h} = a_{t-1} + b h_x^{h-1} s_t \quad \forall h \geq 1 \qquad b = g_x\, h_x \tag{A.4}$$

$$\text{Updating:} \qquad a_t = a_{t-1} + k_t^{-1}\big(z_t - (a_{t-1} + b s_{t-1})\big) \tag{A.5}$$

$$\text{Anchoring function:} \qquad k_t^{-1} = \rho_k k_{t-1}^{-1} + \gamma_k f e_{t-1}^2 \tag{A.6}$$

$$\text{Forecast error:} \qquad f e_{t-1} = z_t - (a_{t-1} + b s_{t-1}) \tag{A.7}$$

$$\text{LH expectations:} \qquad f_a(t) = \frac{1}{1-\alpha\beta} a_{t-1} + b(\mathbb{I}_{nx} - \alpha\beta h)^{-1} s_t \qquad f_b(t) = \frac{1}{1-\beta} a_{t-1} + b(\mathbb{I}_{nx} - \beta h)^{-1} s_t \tag{A.8}$$

This notation captures vector learning (z learned) for intercept only. For scalar learning, $a_t = \begin{pmatrix}\bar{\pi}_t & 0 & 0\end{pmatrix}'$ and b_1 designates the first row of b. The observables (π, x) are determined as:

$$x_t = -\sigma i_t + \begin{bmatrix}\sigma & 1-\beta & -\sigma\beta\end{bmatrix} f_b + \sigma \begin{bmatrix}1 & 0 & 0\end{bmatrix} (\mathbb{I}_{nx} - \beta h_x)^{-1} s_t \tag{A.9}$$

$$\pi_t = \kappa x_t + \begin{bmatrix}(1-\alpha)\beta & \kappa\alpha\beta & 0\end{bmatrix} f_a + \begin{bmatrix}0 & 0 & 1\end{bmatrix} (\mathbb{I}_{nx} - \alpha\beta h_x)^{-1} s_t \tag{A.10}$$

B Target criterion

The target criterion in the simplified model (scalar learning of inflation intercept only, $k_t^{-1} = \mathbf{g}(f e_{t-1})$):

$$\pi_t = -\frac{\lambda_x}{\kappa}\Bigg\{x_t - \frac{(1-\alpha)\beta}{1-\alpha\beta}\Big(k_t^{-1} + ((\pi_t - \bar{\pi}_{t-1} - b_1 s_{t-1}))\mathbf{g}_\pi(t)\Big)$$
$$\Big(\mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1}(1 - k_{t+1+j}^{-1} - (\pi_{t+1+j} - \bar{\pi}_{t+j} - b_1 s_{t+j})\mathbf{g}_{\bar{\pi}}(t+j))\Big)\Bigg\} \tag{B.1}$$

where I'm using the notation that $\prod_{j=0}^{0} \equiv 1$. For interpretation purposes, let me rewrite this as follows:

$$\pi_t = -\frac{\lambda_x}{\kappa} x_t + \frac{\lambda_x}{\kappa}\frac{(1-\alpha)\beta}{1-\alpha\beta}\Big(k_t^{-1} + f e_{t|t-1}^{eve}\mathbf{g}_\pi(t)\Big)\mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i}$$
$$-\frac{\lambda_x}{\kappa}\frac{(1-\alpha)\beta}{1-\alpha\beta}\Big(k_t^{-1} + f e_{t|t-1}^{eve}\mathbf{g}_\pi(t)\Big)\Big(\mathbb{E}_t \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1}(k_{t+1+j}^{-1} + f e_{t+1+j|t+j}^{eve})\mathbf{g}_{\bar{\pi}}(t+j)\Big) \tag{B.2}$$

Interpretation: tradeoffs from discretion in RE + effect of current level and change of the gain on future tradeoffs + effect of future expected levels and changes of the gain on future tradeoffs