Materials 44 - I think I've got the estimation

Laura Gáti

September 9, 2020

Overview


1	Inconsistency was: didn't annualize expectations in true data		
	1.1 Check with same seed as true data and $N=1$	4	
2	Real data	5	
3	Policy isn't a function of k_t^{-1}	7	
A	Model summary	8	
В	Target criterion	8	
\mathbf{C}	Impulse responses to iid monpol shocks across a wide range of learning models	g	

1 Inconsistency was: didn't annualize expectations in true data

0.8 0.70.6 0.50.40.30.2 0.10 -4 -3 -2 -1 0 1 2 3 $--{\rm true}$ —initial —estimated (a) π_t vs. x_{t-k} π_t vs. i_{t-k} 0.1 1 0.40.5 0.20.50 0 0 0 -0.1 $\overline{x_t}$ vs. π_{t-k} $\underset{x_{t} \text{ vs. } x_{t-k}}{2}$ $x_t \text{ vs. } i_{t-k}$ $0 \quad \underset{x_t \text{ vs. } E_{t-k}\pi_{t-k+1}}{\underbrace{2}}$ 0 0 1 0 0 0.5-0.2 0.5 -0.1 -0.4 0 -0.5 -0.6 -0.2 0 $\sum_{i_t \text{ vs.}}^2 \pi_{t-k}$ $\underset{i_t \text{ vs.}}{2} x_{t-k}$ $i_t \text{ vs. } i_{t-k}$ 4 4 0 0 $i_t \text{ vs. } E_{t-k}^2 \pi_{t-k}$ 0 1 -0.1 0 0.50 -0.2 0 -1 $E_t \pi_{t+1}$ vs. π_{t-k} $E_t \pi_{t+1}$ vs. x_{t-k} $E_t \pi_{t+1}$ vs. i_{t-k} 0 $0_{E_t\pi_{t+1}} \text{ vs. } 2_{E_{t-k}\pi_{t-k+1}} 4$ 0.10 $0.03 \\ 0.02 \\ 0.01$ 0.1-0.1 0 0.05-0.2 -0.1 Data $\begin{array}{c}
4 & 0 \\
\text{Initial} & 0 \\
\end{array} \text{Optimal}$ 0 2 2 4 0 (b)

Figure 1: Calibration C, use expectations, ridge tuning = 0.01, initialize at truth


Figure 2: Calibration C, use expectations, ridge tuning = 0.01, initialize above truth, annualize expectations in true data



I think that's success!

1.1 Check with same seed as true data and N=1


Figure 3: Calibration C, use expectations, ridge tuning = 0.01, initialize at truth, annualize expectations in true data




Yes!


2 Real data

Figure 4: Adopt settings from Fig 2



Problem: sensitive to where you initialize.





3 Policy isn't a function of k_t^{-1}

The anchoring function is (A.6): $k_t^{-1} = \sum_i \alpha_i b_i (f e_{t|t-1})$. This essentially eliminates k as a state variable.

A Model summary

$$x_{t} = -\sigma i_{t} + \hat{\mathbb{E}}_{t} \sum_{T=t}^{\infty} \beta^{T-t} \left((1-\beta) x_{T+1} - \sigma(\beta i_{T+1} - \pi_{T+1}) + \sigma r_{T}^{n} \right)$$
(A.1)

$$\pi_t = \kappa x_t + \hat{\mathbb{E}}_t \sum_{T=t}^{\infty} (\alpha \beta)^{T-t} \left(\kappa \alpha \beta x_{T+1} + (1-\alpha) \beta \pi_{T+1} + u_T \right)$$
(A.2)

$$i_t = \psi_\pi \pi_t + \psi_x x_t + \bar{i}_t$$
 (if imposed) (A.3)

PLM:
$$\hat{\mathbb{E}}_t z_{t+h} = a_{t-1} + b h_x^{h-1} s_t \quad \forall h \ge 1 \qquad b = g_x h_x$$
 (A.4)

Updating:
$$a_t = a_{t-1} + k_t^{-1} (z_t - (a_{t-1} + bs_{t-1}))$$
 (A.5)

Anchoring function:
$$k_t^{-1} = \sum_i \alpha_i b_i (f e_{t|t-1})$$
 (A.6)

Forecast error:
$$fe_{t-1} = z_t - (a_{t-1} + bs_{t-1})$$
 (A.7)

LH expectations:
$$f_a(t) = \frac{1}{1 - \alpha \beta} a_{t-1} + b(\mathbb{I}_{nx} - \alpha \beta h)^{-1} s_t$$
 $f_b(t) = \frac{1}{1 - \beta} a_{t-1} + b(\mathbb{I}_{nx} - \beta h)^{-1} s_t$ (A.8)

This notation captures vector learning (z learned) for intercept only. For scalar learning, $a_t = \begin{pmatrix} \bar{\pi}_t & 0 & 0 \end{pmatrix}'$ and b_1 designates the first row of b. The observables (π, x) are determined as:

$$x_t = -\sigma i_t + \begin{bmatrix} \sigma & 1 - \beta & -\sigma \beta \end{bmatrix} f_b + \sigma \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} (\mathbb{I}_{nx} - \beta h_x)^{-1} s_t$$
(A.9)

$$\pi_t = \kappa x_t + \begin{bmatrix} (1 - \alpha)\beta & \kappa \alpha \beta & 0 \end{bmatrix} f_a + \begin{bmatrix} 0 & 0 & 1 \end{bmatrix} (\mathbb{I}_{nx} - \alpha \beta h_x)^{-1} s_t$$
 (A.10)

B Target criterion

The target criterion in the simplified model (scalar learning of inflation intercept only, $k_t^{-1} = \mathbf{g}(fe_{t-1})$):

$$\pi_{t} = -\frac{\lambda_{x}}{\kappa} \left\{ x_{t} - \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_{t}^{-1} + ((\pi_{t} - \bar{\pi}_{t-1} - b_{1}s_{t-1})) \mathbf{g}_{\pi}(t) \right) \right\}$$

$$\left(\mathbb{E}_{t} \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1} (1 - k_{t+1+j}^{-1} - (\pi_{t+1+j} - \bar{\pi}_{t+j} - b_{1}s_{t+j}) \mathbf{g}_{\bar{\pi}}(t+j)) \right) \right\}$$
(B.1)

where I'm using the notation that $\prod_{j=0}^{0} \equiv 1$. For interpretation purposes, let me rewrite this as follows:

$$\pi_{t} = -\frac{\lambda_{x}}{\kappa} x_{t} + \frac{\lambda_{x}}{\kappa} \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_{t}^{-1} + f e_{t|t-1}^{eve} \mathbf{g}_{\pi}(t) \right) \mathbb{E}_{t} \sum_{i=1}^{\infty} x_{t+i}$$

$$-\frac{\lambda_{x}}{\kappa} \frac{(1-\alpha)\beta}{1-\alpha\beta} \left(k_{t}^{-1} + f e_{t|t-1}^{eve} \mathbf{g}_{\pi}(t) \right) \left(\mathbb{E}_{t} \sum_{i=1}^{\infty} x_{t+i} \prod_{j=0}^{i-1} (k_{t+1+j}^{-1} + f e_{t+1+j|t+j}^{eve}) \mathbf{g}_{\pi}(t+j) \right)$$
(B.2)

Interpretation: tradeoffs from discretion in RE + effect of current level and change of the gain on future tradeoffs + effect of future expected levels and changes of the gain on future tradeoffs

C Impulse responses to iid monpol shocks across a wide range of learning models

 $T = 400, N = 100, n_{drop} = 5$, shock imposed at t = 25, calibration as above, Taylor rule assumed to be known, PLM = learn constant only, of inflation only.

Figure 6: IRFs and gain history (sample means)




Figure 7: IRFs and gain history (sample means), continued

