



The risks of shell-boring polychaetes to shellfish aquaculture in Washington, USA: A mini-review to inform mitigation actions

Journal:	<i>Aquaculture Research</i>
Manuscript ID:	ARE-RA-20-Mar-280
Manuscript Type:	Review Article
Date Submitted by the Author:	23-Mar-2020
Complete List of Authors:	Spencer, Laura; University of Washington, School of Aquatic and Fishery Sciences Martinelli, Julieta C.; University of Washington, School of Aquatic and Fishery Sciences King, Teri; University of Washington Washington Sea Grant, NA Crim, Ryan; Puget Sound Restoration Fund, NA Blake, Brady; Washington Department of Fish and Wildlife, NA Lopes, Heather; University of Washington, School of Aquatic and Fishery Sciences Wood, Chelsea L.; University of Washington, School of Aquatic and Fishery Sciences
Keywords:	Polydora, mudworm, invasive species, oyster

SCHOLARONE™
Manuscripts

1 *The risks of shell-boring polychaetes to shellfish aquaculture in Washington, USA:*
2 *A mini-review to inform mitigation actions*

3
4 Short running title: *Minimizing impacts of shell-boring polychaetes*
5

6 Laura H Spencer¹, Julieta C Martinelli¹, Teri L King², Ryan Crim³,
7 Brady Blake⁴, Heather M Lopes¹, Chelsea L Wood¹

8
9 ¹School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98105

10 ²Washington Sea Grant, University of Washington, Shelton, WA 98584

11 ³Puget Sound Restoration Fund, Bainbridge Island, WA 98110

12 ⁴Washington State Department of Fish and Wildlife, Olympia, WA 98501

13
14 Corresponding author: Laura H Spencer, lhs3@uw.edu

15 **ABSTRACT**

16 In 2017, *Polydora websteri*, a shell-boring spionid polychaete worm and cosmopolitan
17 invader, was identified for the first time in Washington State. *Polydora websteri* and some of its
18 congeners bore into the shells of calcareous marine invertebrates, reducing the host's shell
19 integrity, growth, survivorship, and market value. Shell-boring *Polydora* spp. have a history of
20 harming shellfish aquaculture industries worldwide by devaluing products destined for the half-
21 shell market, and requiring burdensome treatments and interventions to manage against
22 infestation. Here, we explore the risks of *Polydora* spp. to the historically unaffected aquaculture
23 industry in Washington State. This mini-review is intended to inform shellfish stakeholders by
24 synthesizing the information needed for immediate action in Washington State. We discuss
25 *Polydora* life history and pathology, summarize the recent documentation of *Polydora* spp. in
26 Washington State, and discuss its history as a pest species globally, including farm management
27 strategies developed in other infested regions. Finally, we review existing regulations that may
28 be leveraged by stakeholders to avoid introduction of *Polydora* spp. into uninfested regions.

29

30 Keywords: Polydora, mudworm, invasive species, oyster

31 INTRODUCTION

32 In 2017, shell-boring *Polydora* spp. polychaete worms were positively identified in
33 Washington State (Figure 1), including the cosmopolitan invader *Polydora websteri* (Martinelli
34 *et al.*, 2019). These parasitic marine polychaetes in the family Spionidae bore into the shells of
35 calcareous marine invertebrates, and may pose an economic and ecological risk to cultured and
36 native shellfish species (Lunz 1941; Simon and Sato-Okoshi 2015). Prior to positive
37 identification in 2017, no native or introduced shell-boring *Polydora* species had been described
38 from Washington State (Martinelli *et al.* 2019; Lie 1968).

39 *P. websteri* is common to many other shellfish aquaculture regions (Simon and Sato-
40 Okoshi 2015), with a broad host range, including seven oyster, one mussel, and three scallop
41 species (Simon and Sato-Okoshi 2015). *Polydora* spp. are colloquially known as mud worms, or
42 mud blister worms, and have a long history of reducing shellfish aquaculture production and
43 value in regions such as Australia, New Zealand, South Africa, Chile, Mexico, the East and Gulf
44 coasts of the United States, Hawaii, New Brunswick, and British Columbia (Table 1). Despite
45 previous observations of *P. websteri* in nearby regions such as British Columbia (Bower *et al.*
46 1992) and California (Hartman 1961), neither benthic surveys nor shellfish growers have
47 historically identified shell-boring mud worms in Washington State. The worm's local history,
48 whether as an invader or a species that was not previously identified, and its state-wide
49 infestation rates are unknown. The 2017 study reports that *Polydora* prevalence in Pacific oysters
50 sampled from public beaches was as high as 53% in one embayment of South Puget Sound
51 (Martinelli *et al.* 2019) and suggests that infestation rates may have recently increased to levels
52 at which observers (e.g., growers, agency personnel) take notice. Ongoing work will determine
53 infestation rates for the Salish Sea and Willapa Bay regions.

54 Given the negative impacts of *Polydora* spp. on shellfish aquaculture in other regions, its
55 presence in Washington State warrants a region-focused review to inform further investigation
56 and stakeholder awareness. Here, we explore *Polydora* spp. as a potential risk to Washington
57 State aquaculture. We summarize *Polydora* pathology and life history, review the recent
58 documentation of this pest in Washington State, discuss its history as a pest species, and finally
59 outline measures that stakeholders can take to mitigate the risks and impacts of *Polydora* spp. to
60 Washington State shellfish aquaculture given existing regulations.

61

62 HOST PATHOLOGY

63 Shellfish infected with boring *Polydora* spp. can have reduced shell integrity, growth,
64 survivorship, and marketability (Morse *et al.* 2015; Simon and Sato-Okoshi 2015). *Polydora* spp.
65 worms bore into calcareous shells and line their tunnel with shell fragments, mucus, and detritus
66 (Figure 2) (Wilson 1928; Zottoli and Carriker 1974). If the tunnel breaches the inner shell
67 surface, the host responds by laying down a layer of nacre to protect itself from the burrow and
68 the worm (Whitelegge 1890; Lunz 1941). This can produce a blister, where a thin layer of shell
69 lies over a mass of anoxic detritus. In oysters, the blister is unsightly, its contents malodorous,
70 and if the blister is breached during shucking the detritus can contaminate oyster meat and brine,
71 detracting from flavor and presentation (Morse *et al.* 2015). Burrows can also decrease shell
72 strength, causing cracks during shipping and handling, and making shucking difficult (Bergman,
73 Elner and Risk 1982; Bishop and Hooper 2005; Calvo, Luckenbach and Burreson 1999; Kent
74 1981). Since half-shell oysters are the most lucrative option for oyster farmers, and *Polydora*-
75 infested oysters are often not salable on the half-shell market, infestation significantly
76 depreciates oyster products.

77 *Polydora* infestation can also devalue other oyster products by compromising growth and
78 survival. *Polydora* worm burden is negatively correlated with growth rate, and while the
79 mechanisms are not fully understood, this may be due to the energetic drain of nacre production
80 (Ambariyanto and Seed 1991; Boonzaaier *et al.* 2014; Handley 1998; Kojima and Imajima 1982;
81 Leonart *et al.* 2003a; Royer *et al.* 2006; Simon 2011; Wargo and Ford 1993). For instance,
82 Pacific oysters (*C. gigas*) infested with *P. websteri* grow more slowly, exhibit more frequent but
83 shorter valve gaping, and have higher blood oxygenation, a sign of metabolic changes (Chambon
84 *et al.* 2007). Infested *C. gigas* also demonstrate a three-fold increase in abundance of
85 Cytochrome P450, a protein involved in the oyster's stress response, which could increase
86 susceptibility to secondary stressors (Chambon *et al.* 2007). Shell strength is negatively
87 correlated with *Polydora ciliata* burden in the mussel *Mytilus edulis*, which increases
88 vulnerability to predation (Kent 1981). Reproductive capacity can be altered by *Polydora*, for
89 instance oocyte size was significantly reduced in infested *C. gigas* (Handley 1998). Interestingly,
90 fecundity in the rock oyster *Striostrea margaritacea* increases with *P. websteri* infestation
91 (Schleyer 1991). The rock oyster could be exhibiting a response to stress from infestation
92 reproducing while resources allow it. Similar phenomena have been documented in nematode-
93 parasitized mice, which produce larger litters than uninfected mice (Kristan 2004; Schleyer
94 1991) and plants that prematurely reproduce ("bolt") during periods of drought (Barnabás *et al*
95 2008). While mortality directly associated with *Polydora* infestation is not common, these
96 studies indicate that shellfish harboring *Polydora* may be more susceptible to secondary
97 stressors, including predation, disease, and environmental stress (Wargo & Ford, 1993).

98

99

100 **POLYDORA LIFE HISTORY**

101 The impact of *Polydora* on shellfish aquaculture arises from its life history as a shell-borer. After
102 a planktonic larval stage, a burrowing *Polydora* worm settles onto the prospective host's shell
103 and begins building a tunnel (Wilson 1928; Loosanoff and Engle 1943; Blake 1969a; Blake and
104 Arnofsky 1999). The worm enters along the margin of the shell and excavates its burrow toward
105 the shell center, then often turns back toward the margin to create a characteristic U-shaped
106 borrow (Figure 2). The worm secretes a viscous fluid to dissolve the calcium carbonate shell
107 material, and uses its specialized segment, the 5th setiger, to stabilize its tunnel during burrowing
108 (Haigler 1969; Zottoli and Carricker 1974). The *Polydora* adult dwells within the tunnel, but can
109 emerge from openings on the outer surface of the host's shell to feed on particles in the water
110 column and materials on the shell surface (Figures 2, 3) (Loosanoff and Engle 1943).

111 *Polydora* spp. reproduction has been thoroughly reviewed by Blake and Arnofsky (1999).
112 Briefly, reproduction occurs when the male deposits sperm in a female's burrow, and the female
113 deposits egg capsules along the burrow wall, with each capsule containing dozens of eggs. Many
114 species are capable of reproducing more than once during a season, and while species vary, one
115 fecund female can produce hundreds of larval progeny (Blake 1969a; Blake and Arnofsky 1999).
116 For instance, *P. websteri* females lay strings of approximately 10 capsules, each containing 50-
117 55 eggs (Blake 1969a; Blake and Arnofsky 1999). Larvae hatch from eggs and emerge from their
118 maternal burrow at the 3-chaetiger stage and are free-swimming until they settle onto a substrate
119 (Orth 1971; Blake 1969a). Growth rate in the larval stage depends on ambient water temperature;
120 thus, the time spent in the water column differs among species and across environmental
121 conditions, and may last as long as 85 days (Blake and Woodwick 1971; Blake and Arnofsky
122 1999). This potential for a long pelagic larval duration, particularly in colder climates, may allow

123 for long dispersal distances (Simon and Sato-Okoshi 2015). Additionally, in some instances,
124 early hatched larvae can feed on underdeveloped eggs (“nurse eggs”), and complete development
125 in the burrow (Haigler 1969). This could result in an individual host’s parasitic burden
126 compounding over time due to high rates of autoinfection. 

127 Understanding when planktonic *Polydora* larvae are most abundant in Washington State
128 will be important for shellfish growers managing infestations, as *Polydora* colonize hosts during
129 the larval phase. Generally, planktonic larval abundance tends to correlate with temperature and
130 phytoplankton abundance, but temporal patterns vary geographically (Blake and Arnofsky 1999;
131 Dorsett 1961). In Maine and New Zealand, *Polydora* larvae are reportedly only in the water
132 column during spring and summer months (March to September) and in Maine peak abundance
133 occurs in May and June (Blake 1969a; Blake 1969b; Handley and Bergquist 1997). In the Sea of
134 Japan off the coast of Russia, *Polydora* spp. larvae are present year round, but abundance peaks
135 in May, then persists at moderate levels through October (Omel’yanenko, Kulikova and Pogodin
136 2004). In the Gulf of Mexico, *Polydora* larvae are found in the water column year-round (Cole
137 2018; Hopkins 1958), and larval abundance peaks in May and/or November, depending on the
138 location (Cole 2018). The breeding season can also vary within a region. For instance, in
139 northern Japan (Hokkaido), *P. variegata* breeding occurs during the warmest months, from
140 August to October (Sato-Okoshi, Sugawara, Nomura 1990). In contrast, in northeastern Japan,
141 *Polydora* larvae (species not reported) are most abundant during winter and spring months, from
142 December through June, and loosely coincide with phytoplankton blooms (Abe, Sato-Okoshi and
143 Endo 2011). Although it has not been confirmed in the field, laboratory experiments indicate that
144 diatoms may be an important larval food source for some *Polydora* species, as opposed to

145 flagellates, and thus larval abundances or recruitment could coincide with diatom blooms
146 (Anger, Anger and Hagmeier 1986).

147 How *Polydora* larvae select settlement locations is not understood. *Polydora* larvae are
148 attracted to light (positively phototactic) during early stages, which is commonly leveraged to
149 isolate polydorid larvae from plankton samples (Ye *et al.* 2017). *Polydora* readily recruit to dead
150 oyster shells, so larvae probably do not respond to chemical cues from live hosts, but may
151 respond to chemical or tactile signatures from shells (Clements *et al.* 2018). Some studies
152 indicate that *Polydora* spp. may prefer to colonize certain mollusc species over others, possibly
153 due to shell traits such as texture and size (Ambariyanto and Seed 1991; Lemasson and Knights
154 2019). Higher infestation rates were reported in *Ostrea edulis* compared to *C. gigas* (Lemasson
155 and Knights 2019). Compared to *C. virginica*, however, *C. gigas* was more susceptible to
156 *Polydora* infestation, which the authors attributed to the thinness of *C. gigas* shells (Calvo *et al.*
157 1999). Larger hosts are commonly infested with more worms. In the surf clam, *Mesodesma*
158 *donacium*, infestation rates increase with size and juveniles smaller than 34 mm do not harbor
159 any *Polydora* spp., suggesting a shell size threshold for settlement (Riascos *et al.* 2008). Stressed
160 or unhealthy hosts may be more prone to *Polydora* spp. infestation. When exposed to petroleum
161 pollutants from the Providence River system, the hard clam *Mercenaria mercenaria* is more
162 likely to be infested with *Polydora*; the authors suggest that the pollutants alter clam burrowing
163 behavior, increasing the chances of *Polydora* colonization (Jeffries 1972). Finally, *Polydora*
164 infestation may differ among locations due to environmental conditions, particularly salinity. A
165 recent survey of wild *C. virginica* in two Gulf of Mexico estuaries found that *P. websteri*
166 prevalence and abundance decrease with increasing salinity, with a marked drop in infestation at
167 salinities exceeding 28 ppt (Hanley *et al.* 2019). High infestation rates were reported for *C. gigas*

168 and *C. virginica* grown in low- and moderate salinity locations across Virginia, but infestation
169 rates were much lower in areas with high salinity (Calvo *et al.* 1999). *Polydora* infestation has
170 also been associated with low-salinity environments in the Indian backwater oyster *C.*
171 *madrasensis* (Stephen 1978). In Gulf of Mexico farms, *P. websteri* was reportedly least abundant
172 in *C. virginica* where salinity was most variable (Cole 2018). Whether salinity influences the
173 current *Polydora* spp. distribution and abundance in Washington State is not yet clear.

174

175 RECENT *POLYDORA* IDENTIFICATION IN WASHINGTON STATE

176 Historically, Washington shellfish farmers have not reported losses from shell-boring *Polydora*
177 on their farms, and until recently no shell-boring *Polydora* species had been formally
178 documented from the state. Related spionid polychaetes have been present, such as *Polydora*
179 *cornuta* (Fermer & Jumars 1999), *Pseudopolydora* spp. (e.g. Woodin 1984), and *Boccardia*
180 *proboscidea* (Hartman 1940, Oyarzun *et al.* 2011). These are primarily benthic species, and
181 while they can occupy mud deposits within oyster shell crevices, they do not burrow and
182 therefore do not create blisters.

183 In 2017, mud worm blisters were noticed in increasing abundance in cultured Pacific
184 oysters from southern Puget Sound, which triggered a preliminary survey. Martinelli *et al.*
185 (2019) sampled Pacific oysters from public beaches in Totten Inlet and Oakland Bay (Figure 1).
186 Across the two sites, 41% of oysters were infested with a shell-boring worm (53% of Oakland
187 Bay oysters, 34% of Totten Inlet oysters) (Martinelli *et al.* 2019). The worm species was
188 identified using morphology (from scanning electron microscope images), and phylogenetics
189 (comparing 18s rRNA & mtCOI sequences against published *Polydora* sequences). Some of the
190 worms collected from Oakland Bay were positively identified as *P. websteri*, while others did

191 not group with any of the available sequences and their identity remains unresolved
192 (phylogenetic trees from Martinelli *et al.* 2019 are reproduced in Figures 4 & 5).

193 It is unknown whether *P. websteri* was historically present in Washington State at low
194 abundance or recently introduced. If the species was recently introduced, eradication might be
195 possible (see Williams & Grosholz, 2008 for examples of successful programs). But if
196 eradication of *P. websteri* is not possible, it could still be contained to a few Puget Sound basins
197 through education, mitigation, and regulation (Çinar 2013; Paladini *et al.* 2017). If *P. websteri*
198 has been present but dormant, the high infestation intensity reported by Martinelli *et al.* (2019)
199 may be the result of a recent outbreak, caused by factors such as genetic changes, relaxation of
200 biotic pressures (e.g. predators), or environmental changes (e.g., ocean warming, siltation)
201 (Crooks 2005; Clements *et al.* 2017a).

202 Washington State aquaculture produces 45% of the molluscs cultured in the U.S. (2013,
203 USDA) and is an iconic industry that supports rural communities, protects water quality, and
204 collaborates closely with research and restoration programs. Within Washington, Puget Sound
205 growers produce 70% of the state's shellfish (80% by value, over \$92 million annually),
206 concentrated mostly in South Puget Sound, where the *Polydora*-infested oysters were sourced
207 (Figure 1). Economic losses associated with *Polydora* outbreaks in this highly productive
208 shellfish region could have nation-wide repercussions for the aquaculture industry.

209

210 IMPACTS TO AQUACULTURE PRODUCTION

211 *Polydora* has caused economic losses for shellfish aquaculture operations worldwide. Of the
212 shell borers, *P. websteri*, *P. ciliata*, and *P. hoplura* are the most widely distributed and notorious
213 for infesting shellfish farms (Radashevsky *et al.* 2006) (Table 1). The primary impact is product



214 devaluation due to negative consumer responses to blisters and anoxic material within the inner
215 shell, particularly in freshly shucked oysters (Shinn *et al.* 2015). In rare instances, large mortality
216 events have been attributed to *Polydora* infestation. For instance, in British Columbia, *P.*
217 *websteri* caused up to 84% mortality in scallop grow-out sites from 1989 to 1990, resulting in up
218 to US \$449,660 in lost revenue that year (Shinn *et al.* 2015, Bower *et al.* 1992). In Tasmania and
219 South Australia, *P. hoplura* killed over 50% of abalone stocks between 1995 and 2000, causing
220 an estimated US \$550,000 to \$1.16 million in losses per year (Shinn *et al.* 2015). In the summer
221 of 1997, one million juvenile scallops were culled in a Norwegian nursery due to a *Polydora* spp.
222 infestation; as a result, one-third of Norway's 1997 scallop cohort was lost (Mortensen *et al.*
223 2000). In 1998, intense infestations (up to 100 worms per oyster) of *P. ciliata* in *C. gigas* oysters
224 in Normandy, France correlated with considerable reduction in growth and meat weight, which
225 may have contributed to unusually high summer mortality rates of up to 51% (Royer *et al.* 2006).

226 In other regions, *Polydora* infestations have made certain growing practices impractical
227 or unprofitable. In New Zealand, fattening intertidally-grown oysters in longlines for a few
228 weeks prior to sales improves oyster condition, but this practice is not recommended due to the
229 risk it entails of *Polydora* spp. infestation (Curtin 1982). Following the collapse of native *C.*
230 *virginica* in North Carolina, triploid *Crassostrea ariakensis* were assessed for culture. Feasibility
231 was contingent on harvesting oysters prior to summer months to avoid *Polydora* colonization, as
232 revenue would be lost if infestation rate exceeded 54% (Bishop & Peterson 2005; Grabowski *et*
233 *al.* 2007). Many regions have experienced chronic *Polydora* infestation for decades (*e.g.*, South
234 Africa and New South Wales, Australia). Growers incur costs associated with cleaning or
235 treating stocks to control *Polydora*, and having grow-out methods restricted to specific high tidal
236 heights or locations, but these economic impacts have not been quantified.

237

238 **MANAGEMENT STRATEGIES DEVELOPED IN OTHER REGIONS**

239 In regions with noxious *Polydora* spp., producers control infestation by modifying gear and grow
240 methods, and treating shellfish stocks regularly. Farm management approaches focus on keeping
241 oysters free of mud and air drying oysters by growing them at high tidal elevations (Morse *et al.*
242 2015; Handley & Bergquist, 1997). Since the early 20th century, Australian oyster farmers in
243 New South Wales have used off-bottom growing methods with long tidal exposures to reduce
244 mud worm infestation rates (Smith 1981; Diggles 2013; Ogburn 2011). Oysters are grown at
245 approximately the mean low water neap height using rack and rail, long-line, and elevated tray
246 systems, such that stocks are exposed for 30 percent of each daily tidal cycle (Ogburn 2011). On
247 the U.S. Atlantic Coast, researchers report that exposing *C. virginica* for 40 percent of a tidal
248 cycle is an effective method of avoiding substantial *Polydora* infestation (Littlewood *et al.*
249 1992). Growing oysters in bags that are easily raised above the water line for aerial exposures
250 can also reduce infestation rates, particularly during the *Polydora* breeding season. For instance,
251 some growers on the U.S. Gulf Coast use floating cages and rack-and-rail systems to easily
252 expose bags weekly for up to 24 hours (Gamble 2016; Cole 2018). These off-bottom methods
253 have proven effective for avoiding high rates of infestation, but do slow oyster growth rates
254 (Ogburn *et al.* 2007; Nell 2007; Nell 2001), and do not always prevent infestation (Cole 2018;
255 Clements *et al.* 2017a). For instance, recent *Polydora* outbreaks were reported in oysters
256 suspended off-bottom in New Brunswick, Canada and may have been related to high siltation
257 levels, which can increase *Polydora* infestation rates (Clements *et al.* 2017a). Increasing cleaning
258 frequency to reduce siltation may therefore help to control *Polydora*, particularly in areas with
259 heavy siltation. Frequent cleaning can also reduce impacts of non-boring *Polydora* species, such

260 as *P. nuchalis* and *P. cornuta*, which foul culture equipment with large masses of sediment and
261 tubes (Bailey-Brock 1990). 

262 A variety of treatments have been developed to kill worms in oysters infested with
263 *Polydora* spp. Methods include freshwater soaks (up to 72 hours), salt brine soaks (up to 5
264 hours), extended cool air storage (up to 3-4 weeks at 3°C), heat treatments (e.g., 40 seconds at
265 70°C), chemical treatments (e.g., chlorine, iodine), and various combinations thereof. Treatment
266 efficacy can differ among species, season, and exposure duration, but generally the most
267 commonly used treatments are hyper-saline dips followed by air drying, and extended cold-air
268 storage. For Washington State growers, hyper-saline dips followed by air drying may be a
269 feasible treatment regime, but precise methods will need to be developed for local conditions and
270 species. For *C. virginica* and *C. ariakensis* grown in North Carolina, weekly treatments using a
271 20-minute hypersaline dip followed by air drying for 2 hours reduced *Polydora* spp. infestation
272 to only 5% from up to 47.5% in untreated oysters (Bishop and Hooper 2005). Currently, the most
273 effective treatment appears to be the “Super Salty Slush Puppy” (SSSP), first developed by Cox
274 *et al.* (2012). The protocol involves a 2-minute full submersion of oysters in brine (250 g/L)
275 between -10°C and -30°C (i.e., ice-water), followed by air drying for 3 hours. The SSSP also
276 effectively kills other fouling epibionts, such as barnacles. Petersen (2016) recently compared the
277 SSSP method against other saltwater, freshwater, and chemical dips followed by air exposure for
278 infested *C. gigas*, and confirmed SSSP as the best method, killing 95% of *P. websteri* while
279 causing only minimal oyster mortality.

280 Freshwater immersion is another treatment option for Washington growers, and for some
281 host or polychaete species, may be more effective than hypersaline dips. For Chilean flat oysters
282 (*Tiostrea chilensis*), freshwater immersion for 180-300 minutes was more effective than

283 hypersaline immersion (64 ppt) at killing *Boccardia acus*, another shell-boring polychaete
284 species (Dunphy, Wells and Jeffs 2005). In heavily infested *C. virginica*, nearly 98% *Polydora*
285 mortality was achieved with a 3-day freshwater immersion followed by four days of cold-air
286 storage (Brown 2012). Without the cold-air storage, the freshwater immersion only killed 25-
287 60% of *Polydora*, and worms occupying deep burrows were unaffected (Brown 2012).

288 Interestingly, worms that were removed from burrows and placed in freshwater were killed
289 within three days, which highlights the protection that shell burrows provide for *Polydora* worms
290 (Brown 2012). In other regions, chemical treatments have effectively controlled *Polydora*
291 infestation (Gallo-Garcia *et al.* 2004). However, environmental and health and safety regulations
292 will probably preclude chemicals from being used in Washington State (Morse *et al.* 2015).

293 Treating infested oysters mitigates the effects of severe infestation, but costs may be
294 prohibitive. Growers incur expenses associated with handling and specialized equipment (Nell
295 2007). Modifying grow methods to accommodate frequent *Polydora* treatments, or to minimize
296 secondary stressors following treatments, may also be necessary. Treatment costs also depend on
297 reinfection rates, which occur more readily on farms that harbor *Polydora* reservoirs, such as
298 dead oyster shell or wild shellfish growing nearby (Clements *et al.* 2018; Lemasson and Knights
299 2019). Many of the existing treatments have been developed for species not commonly grown in
300 Washington State. A common treatment for *C. virginica* is long-term cold-air storage. Maine
301 growers have found that after 3-4 weeks (~3°C), 100% of adult *Polydora* worms are killed, with
302 minimal *C. virginica* mortality (Morse *et al.* 2015). Prolonged air exposure is also commonly
303 used for the Australian oyster *Saccostrea glomerata* (7-10 days, in the shade; Nell 2007). These
304 oyster species have different physiological tolerances than *C. gigas*, the dominant aquaculture
305 species in Washington, and therefore the same treatments may not be feasible for many of the

306 state's oyster growers (Morse *et al.* 2015; Nell 2007). For instance, while *C. virginica* can
307 survive cold-air storage for six months with ~80% survival, no *C. gigas* seed or adults survived
308 similar cold-air conditions after 20 weeks of storage (Hidu, Chapman and Mook 1998). Irrigating
309 stored *C. gigas* continuously with seawater can increase survival in cold air storage (52% adults
310 and 80% juveniles at 7°C), but whether irrigation also increases *Polydora* survival is not known
311 (Seaman 1991).

312 Oyster mortality can be an issue following treatments for *Polydora* (Nell 2007). Growers
313 are highly encouraged to test treatments on a small number of oysters before applying it to large
314 batches (Morse *et al.* 2015). Making adjustments to grow methods might be necessary to
315 improve oyster survival following treatments. For instance, increasing flow rates in a nursery
316 upweller system can increase *C. ariakensis* and *C. virginica* survival following hypersaline and
317 drying treatments (Bishop and Hooper 2005). More details and recommendations for treatment
318 options are available in Morse *et al.* (2015) and Nell (2007).

319 It is important to recognize that the majority of treatments to kill *Polydora* have been
320 developed for oysters (but see Bilbao *et al.* 2017 and Lleonart, Handliger & Powell 2003b for
321 abalone treatments). Shellfish species that are sensitive to exposures cannot be treated using
322 these extreme methods, and therefore are vulnerable to infestation and may provide refuge to
323 *Polydora*. Finally, no method to date has assessed whether these interventions render *Polydora*
324 eggs inviable, which is an important question that needs to be answered.

325

326 **POLYDORA INTRODUCTION VIA SHELLFISH TRANSLOCATION**

327 *Polydora* spp. have a long history of accompanying shellfish during translocation and becoming
328 invasive pests. In the early 1880's, oysters believed to have been infected with *P. ciliata* were

329 imported from New Zealand into the George's River in Southeast Australia. Before being sold in
330 Australian markets, they were routinely refreshed or fattened in bays adjacent to native shellfish
331 beds (Roughley 1922; Edgar 2001; Ogburn 2007). By 1889, mud worm outbreaks had infected
332 thirteen separate estuaries in the region, and oyster growers abandoned leases that were below
333 the low-water mark (Roughley 1922). The introduction and translocation of mud worm species
334 to Australia may have contributed to the disappearance of native subtidal oyster beds (*Saccostrea*
335 *glomerata*, *Ostrea angasi*), some of which never recovered (Diggle 2013; Ogburn 2011). More
336 recently, *Polydora* spp. were introduced into Hawaii, probably from stock shipped from 
337 mainland United States or Mexico (Eldredge 1994). In one notable case, *P. websteri* brought to
338 Oahu via California oyster seed resulted in a severe infestation, and caused farmers to abandon
339 their land-locked oyster pond (Bailey-Brock and Ringwood 1982).

340 When invasive *Polydora* spp. are introduced to new regions, they can disperse during
341 their planktonic larval stage to infect other shellfish within a basin (Simon and Sato-Okoshi
342 2015; Blake and Arnofsky 1999; David *et al.* 2014; Hansen *et al.* 2010). As shellfish farmers
343 grow oysters in high-density bags, racks, or lines, a *Polydora* infestation can spread readily
344 within a farm, and the subsequent movement of stock is considered the primary pathway for
345 *Polydora* introduction into new regions (Simon and Sato-Okoshi 2015; Moreno *et al.* 2005). 
346 *Polydora* worms do not usually kill the host, nor do they inhabit living host tissue, so infections
347 can go undetected via traditional disease screening and may not be recognized until an area is
348 fully infested (Korringa 1976). The infection mechanism might explain why *Polydora* spp. were
349 found to be very prevalent in the year in which the infections were first reported from Puget
350 Sound (up to 53% of *C. gigas* infected in Oakland Bay) (Martinelli *et al.* 2019). Many *Polydora*
351 species have broad host ranges, making it possible for all cultured shellfish species in

352 Washington State to be infested, including the native Olympia oyster (*Ostrea lurida*) and
353 introduced *C. gigas*, *C. virginica*, and *C. sikamea*. Furthermore, *Polydora* species can persist in
354 non-cultured reservoir hosts, regardless of growers' control treatments, making it difficult to
355 eradicate from a farm (Moreno *et al.* 2006).

356

357  **EXAMPLES OF *POLYDORA* MONITORING AND REGULATIONS GLOBALLY**

358 In Australia, *Polydora* spp. have been common since they were introduced in the late 1800's, and
359 are not identified as invasive species but are considered pests to abalone and oyster growers. In
360 New South Wales, the Department of Primary Industries continues to develop and test control
361 measures for shellfish farmers (Nell 2007). In 2005, Tasmania developed a comprehensive
362 management program for mud worm control in cultured abalone in response to outbreaks
363 (Handlinger *et al.* 2004). In Victoria, Australia, the Abalone Aquaculture Translocation Protocol
364 categorizes mud worms as a "significant risk", and now regulates the movement of infected
365 stock to uninfected areas (Victorian Fisheries Authority 2015). In New Brunswick, Canada the
366 Canadian Aquaculture Collaborative Research and Development Program (ACRDP) recently
367 funded a project to identify potential causes of increasing, sporadic *P. websteri* outbreaks in off-
368 bottom oyster sites. Despite Canada characterizing *Polydora* spp. as a Category 4 species of
369 "negligible regulatory significance in Canada," the recent outbreaks raise questions about the
370 potential for *Polydora* spp. intensity to shift geographically and over time, particularly in
371 response to changing climate conditions (Government of Canada and Services 2017).

372

373 **STATUS OF *POLYDORA* MONITORING AND REGULATIONS IN THE USA**

374 Marine polychaete species, including shell-boring *Polydora* spp., are not monitored or regulated
375 in the United States. According to a 2013 review, 292 polychaete species (15% of all described
376 polychaetes) have been relocated to new marine regions via human transport. Of these, 180 are
377 now established and 16 are in the genus *Polydora* (Çinar 2013). Despite this, there is no
378 international or national governing body regulating this transport, and aquatic parasites are not
379 recognized as invasive or injurious species in the United States. For example, the U.S.
380 Geological Services list of Nonindigenous Aquatic Species includes only two annelids, both
381 freshwater species (USDI n.d.). While the United States Department of Agriculture's 2017
382 reportable disease list does include seven molluscan parasites, it does not include shell-boring
383 polychaetes (USDA 2017).

384 The ubiquity of *Polydora* species and their long history as pests in the Atlantic and Gulf
385 Coasts may be the reason for this lack of federal regulation (Lunz 1941; Lafferty and Kuris
386 1996). Nevertheless, researchers and government agencies continue to help Atlantic and Gulf
387 farmers control infection. In the past five years, the Maine Sea Grant (Morse *et al.* 2015),
388 Alabama Cooperative Extension System (Walton *et al.* 2012; Gamble 2016), New Jersey Sea
389 Grant (Calvo *et al.* 2014), and the USDA Sustainable Agriculture Research & Education (USDA
390 Grant no. FNE13-780) invested in communication tools and methods for farmers to mitigate the
391 effects of mud worm on their shellfish products. These investments highlight that *Polydora* is an
392 ongoing, real issue for farmers in infected regions, and that Washington growers may need to
393 respond if *Polydora* prevalence continues to increase in the state.

394

395 **LIVE SHELLFISH REGULATIONS IN WASHINGTON STATE**

396 In Washington State, regulations are in place to avoid introducing diseases and invasive species,
397 which are identified in the Washington Administrative Code (WAC). Here, we review existing
398 Washington State code to highlight regulations that control the spread of invasive species
399 throughout the state, which may be leveraged to limit movement of shellfish heavily infested
400 with *Polydora* spp. to uninfected regions, if warranted.

401 Under WAC 220-340-050 and WAC 220-370-200, import permits are mandatory for any
402 entity importing live shellfish from outside Washington State for any purpose, such as
403 aquaculture, research, or display, but excluding animals that are market-ready and not expected
404 to contact Washington waters. Import permits require a “clean bill of health” certifying that the
405 origin is disease-free, and free of the invasive green crab (*Carcinus maenas*) and oyster drills
406 (*Urosalpinx cinerea* and *Ocinebrellus inornatus*). The WDFW import permits can require that
407 clam, oyster, and mussel seed or stock intended to touch Washington waters be treated for the
408 invasive green crab using a dilute chlorine dip (WDFW, n.d.); this treatment may be effective
409 against shell-boring species such as *Polydora* spp., but has yet to be tested. In instances where
410 the chlorine dip is lethal (e.g., mussels and geoduck), imports are only allowed from locations
411 isolated from European green crab-infested waters, and thus the treatment is not required. The
412 chlorine dip has not been evaluated for use against *Polydora*. If effective, it could be adopted as
413 a treatment required by WDFW when translocating stocks from areas with heavy *Polydora*
414 infections. Transfer permits are also required under WAC 220-340-150 when moving adult
415 shellfish and seed between and within Washington State basins. These permits are regulated by
416 the Washington State Department of Fish and Wildlife (WDFW). Oyster shell (cultch), which is
417 moved throughout the state for oyster bed enrichment and hatchery seeding for farming and
418 restoration purposes, is required to be "aged" out of the water for a minimum of 90 days and is

419 inspected by WDFW prior to placement into state waters, so it is unlikely to translocate viable
420 *Polydora* worms or eggs (WDFW, personal communication). Permits do not certify that
421 translocated organisms are free of *Polydora* spp., as they are not currently designated as invasive
422 or pest species.

423 Under WAC 220-370-200 and WAC 220-370-180, aquaculture groups must report any
424 disease outbreak to the WDFW. Consequently, hatchery staff and farmers monitor for large
425 mortality events that indicate disease. Widespread mortalities due to infectious pathogens are
426 common to shellfish aquaculture. However, aided by diligent stakeholders, Washington has so
427 far avoided several of the most notorious diseases infecting other regions, such as oyster herpes
428 virus variants (e.g. OsHV-1 found in Tomales Bay, CA), the highly lethal OsHV-1 microvariant
429 (OsHV-1 μVar, recently found in San Diego, CA, likely transferred from Europe or Oceania),
430 abalone withering syndrome (present in California), dermo (*Perkinsus marinus*, Gulf and
431 Atlantic Coasts of USA), Pacific oyster nocardiosis (Atlantic and Gulf Coast), MSX disease
432 (*Haplosporidium nelsoni*, detected in British Columbia), and bonamiasis (although bonamiasis
433 was once identified in WA in oyster stock sourced from California) (Elston *et al.* 1986;
434 Alfjorden, *et al.* 2017; Meyer 1991). These regulations do not currently require *Polydora*
435 infestation to be reported, as it is not a designated disease.

436

437 **STAKEHOLDER COMMUNICATION AND RESEARCH NEEDS IN WASHINGTON STATE**

438 To minimize the impact of *Polydora* spp. on Washington State shellfish aquaculture,
439 stakeholders need to be informed of the risks of *Polydora* infestation and treatment options.
440 Shellfish growers should be equipped to recognize *Polydora*-infected product, and to understand
441 the impact *Polydora* could have on their businesses. Growers in uninfected regions may wish to

442 inspect for *Polydora* before translocating shellfish to their properties. The best method to screen
443 for *Polydora* in oysters is to shuck and inspect the inside of the valves for evidence of burrowing
444 and blisters (Figure 2) (Bower *et al.* 1994). If *Polydora* is found on their properties, shellfish
445 growers and aquaculture facilities will probably need to implement treatment measures to control
446 *Polydora* spp. in their products, and to avoid further spread. While prior work in other regions
447 provides some hints as to which treatments might work for eliminating *Polydora*, growers
448 require information on the relative efficacy and practicality of these treatments in local
449 conditions, on locally cultured species, and on whether existing handling practices can be
450 effective against the worm. For example, air drying during long tidal exposures, or
451 environmental conditions such as high salinity, may mitigate or inhibit *Polydora* infestation in
452 some areas (e.g., coastal estuaries such as Willapa Bay).

453 Hatcheries and nurseries produce shellfish seed that is sold to growers in Washington
454 State. These facilities are particularly important in pest management, since they are the nodes
455 from which a significant portion of shellfish move about the region. Oyster larvae are reared in
456 the hatchery, sent to nurseries to grow to seeding size, and then are distributed to shellfish
457 growers. Broodstock are frequently held in one location, brought to the hatchery for spawning 
458 and returned. As a result, hatchery production involves moving oysters multiple times throughout
459 their lifespans. Shellfish seed are also imported into Washington from hatcheries in Canada,
460 Hawaii, California, and Oregon. Hatcheries and nurseries may need to update biosecurity
461 protocols to inspect and treat translocated stock for *Polydora*. How infestation rate and
462 abundance change as a function of shellfish seed size and age, and whether viable *Polydora* spp.
463 eggs can be transferred alongside translocated shellfish larvae, will be important considerations
464 and require additional research.

465 To better inform Washington State stakeholders and to control further human-aided
466 spread into uninfected areas, *Polydora* presence and baseline infestation rates need to be fully
467 established with a quantitative survey of live oysters. To understand why *Polydora* infestation
468 rates are higher in certain areas, sampling site details should be collected alongside the
469 distribution survey, including sediment type, culture gear type and tidal elevation, and
470 environmental data such as salinity, temperature, and pH (Calvo *et al.* 1999; Clements *et al.*
471 2017b; Cole 2018). Species distributions will inform potential regulatory and control actions. It
472 is possible that *Polydora* spp. have been present in Washington State at low levels of abundance
473 for many years, perhaps controlled by environmental conditions, local ecology, or culture
474 techniques. Environmental data will also help to characterize *Polydora* spp. potential impact on
475 shellfish aquaculture under projected climate conditions. Finally, phytoplankton abundance and
476 community composition should be monitored in areas where *Polydora* has been positively
477 identified to understand factors predicting *Polydora* larval abundance. Predicting when and
478 where larvae are most likely to colonize shellfish may allow growers to relocate products
479 temporarily (e.g., higher tidal height) to avoid infestation.
480
481

482 CONCLUSION

483 *Polydora* spp. have a long history of invasion via oyster translocation, of devaluing shellfish
484 products, and of necessitating treatments or changes to growing methods. Historically,
485 Washington State has been one of the few regions worldwide unaffected by shell-boring
486 *Polydora* spp., but that time has unfortunately passed, with the recent confirmation of *P. websteri*
487 in southern Puget Sound. To minimize the risk of *P. websteri* and other shell-boring *Polydora*
488 spp. to the Washington State shellfish industry, early signs of infestation should be addressed by
489 mapping current distribution, alerting the shellfish industry of the risk, and if warranted,
490 leveraging or augmenting regulations to control further spread and introduction of other shell-
491 boring polychaetes. More broadly, federal regulatory gaps should be addressed for better
492 monitoring of pest species harbored by and deleterious to cultured shellfish.

493

494 ACKNOWLEDGEMENTS

495 The authors thank Ralph Elston, Brent Vadopalas, and Sandy Shumway for providing comments
496 and guidance during the research and development of this manuscript.

497

498 DATA AVAILABILITY STATEMENT

499 Data sharing is not applicable to this article as no new data were created or analyzed in this
500 study.

501

502

503 CONFLICT OF INTEREST STATEMENT

504 We have no conflict of interest to disclose.

505 **Tables**

506 **Table 1:** Reports of *Polydora* spp. infestations in cultured shellfish. Studies include those that
 507 identified *Polydora* spp. in shellfish grown on farms or in culture experiments, and omits
 508 infestations documented in wild-collected shellfish.

Country	Region	<i>Polydora</i> species	Cultured host species	Reference
Australia	New South Wales	spp.	<i>Saccostrea glomerata</i>	Wisely, Holiday & Reid 1979
Australia	South Australia	<i>P. haswelli</i> <i>P. hoplura</i> <i>P. websteri</i>	<i>Mytilus edulis</i>	Pregenzer 1983
Australia	New South Wales, southern Queensland	spp.	<i>Saccostrea glomerata</i>	Nell 1993
Australia	Tasmania	<i>P. hoplura</i>	<i>Haliotis rubra</i> ; <i>Haliotis laevigata</i>	Leonart, Handlinger & Powell 2003a
Australia	Southwest	<i>P. cinata</i> <i>P. haswelli</i> <i>P. aura</i>	<i>Haliotis laevigata</i> ; <i>Haliotis roei</i> ; <i>Saccostrea commercialis</i>	Sato-Okoshi, Okoshi & Shaw 2008
Belgium	Bassin do Chasse, Ostend harbour	<i>P. ciliata</i>	<i>Ostrea edulis</i>	Daro & Bofill 1972
Brazil	Southern Brazil	spp.	<i>Crassostrea rhizophorae</i>	Nascimento 1983
Brazil	Santa Catarina, Ribeirão da Ilha	spp.	<i>Crassostrea gigas</i>	Sabry et al. 2011
Brazil	São Francisco River estuary, Sergipe state, northeastern Brazil	spp.	<i>Crassostrea gasar</i>	da Silva et al. 2015
Canada	New Brunswick	<i>P. websteri</i>	<i>Crassostrea virginica</i>	Clements et al. 2017a
Chile	Herradura Bay	spp.	<i>Ostrea chilensis</i>	Di Salvo & Martinez 1985
Chile	Tongoy Bay, Coquimbo	Unknown species similar to <i>P. ciliata</i>	<i>Argopecten purpuratus</i>	Basilio, Canete & Rozbaczylo 1995
China	Shandong Peninsula and Shanghai in eastern China	<i>P. onagawaensis</i> <i>P. brevipalpa</i> <i>P. websteri</i>	<i>Patinopecten yessoensis</i> ; <i>Haliotis discus hannai</i> ; <i>Chlamys farreri</i> ; <i>Crassostrea gigas</i>	Sato-Okoshi, Okoshi & Abe 2013

Costa Rica	Chomes, Gulf of Nicoya	spp.	<i>Crassostrea rhizophorae</i> ; <i>Crassostrea gigas</i>	Zuniga, Zurburg & Zamora 1998
France	Bay of Arcachon	spp.	<i>Ostrea edulis</i>	Robert, Borel, Pichot & Trut 1991
France	Bay of Brest, Brittany	spp.	<i>Crassostrea gigas</i>	Mazurie et al. 1995
France	Brittany	<i>P. ciliata</i> <i>P. hoplura</i>	<i>Crassostrea gigas</i>	Fleury et al. 2001
France	Brittany	spp,	<i>Crassostrea gigas</i>	Fleury et al. 2003
France	Normandy	spp.	<i>Crassostrea gigs</i>	Ropert, Pien, Mary & Bouchaud 2007
India	Gulf of Mannar	spp.	<i>Pinctada fucata</i>	Alagarswami & Chellam 1976
Indonesia	Padang Cermin Bay, Lampung.	spp.	<i>Pinctada maxima</i>	Hadiroseyan, Djokosetyianto & Iswadi 2007
Ireland	Guernsey, Kent	spp.	<i>Crassostrea gigas</i>	Steele & Mulcahy 1999
Ireland	Dungarvan, County Waterford	spp.	<i>Crassostrea gigas</i>	Steele & Mulcahy 2001
Italy	Adriatic Sea	<i>P. ciliata</i>	<i>Tapes philippinarum</i>	Boscolo & Giovanardi 2002
Italy	Venice Lagoon, North Adriatic Sea	<i>P. ciliata</i>	<i>Tapes philippinarum</i>	Boscolo & Giovanardi 2003
Japan	Abashiri Bay	<i>P. variegata</i>	<i>Patinopecten yessoensis</i>	Sato-Okashi, Sugawara & Nomura 1990
Japan	Unknown, not in english	spp.	<i>Pinctada fucata</i>	Wada & Masuda 1997
Japan	10 sites across Japan	<i>P. brevipalpa</i> <i>P. uncinata</i> <i>P. aura</i>	<i>Crassostrea gigas</i> ; <i>Patinopecten yessoensis</i> ; <i>Haliothis discus hannai</i> ; <i>Haliothis discus discus</i> ; <i>Haliothis gigantea</i> ; <i>Haliothis laevigata</i> ; <i>Haliothis roei</i> ; <i>Haliothis diversicolor supertexta</i> ; <i>Pinctada fucata</i>	Sato-Okoshi & Abe 2012
Korea	South and West coasts	<i>P. haswelli</i> <i>P. aura</i> <i>P. uncinata</i>	<i>Crassostrea gigas</i> ; <i>Pinctada fucata</i> ; <i>Haliothis discus discus</i>	Sato-Okoshi et al. 2012

Mexico	Baja California	spp.	<i>Crassostrea gigas</i>	Caceres-Martinez, Macias-Montes De Oca & Vasquez-Yeomans 1998
New Zealand	Bay of Islands	spp.		Curtin 1982
New Zealand	Marlborough Sound	<i>P. websteri</i> <i>P. hoplura</i>	<i>Crassostrea gigas</i>	Handley 1995
New Zealand	Houhora Harbour	<i>P. websteri</i> <i>P. hoplura</i>	<i>Crassostrea gigas</i>	Handley & Bergquist 1997
New Zealand	Houhora Harbour	spp.	<i>Crassostrea gigas</i>	Handley 2002
New Zealand	Manukau Harbour	Not a <i>Polydora</i> species, but related shell-boring polychaete, <i>Boccardia acus</i>	<i>Tiostrea chilensis</i>	Dunphy, Wells & Jeffs 2005
New Zealand	North Island & Coromandel	<i>P. websteri</i> <i>P. haswelli</i>	<i>Crassostrea gigas</i> ; <i>Perna canaliculus</i>	Read 2010
Russia	Sea of Japan	<i>P. brevipalpa</i>	<i>Patinopecten yessoensis</i>	Silina 2006
Russia	Sea of Japan	<i>P. brevipalpa</i>	<i>Mizuhopecten yessoensis</i>	Gabaev 2013
South Africa	Port Elizabeth	<i>P. hoplura</i>	<i>Crassostrea gigas</i>	Nel, Coetze & Van Niekerk 1996
South Africa	west, south, and east coasts	<i>P. hoplura</i>	<i>Haliotis midae</i>	Simon, Ludford & Wynne 2006
South Africa	Hermanus	Not a <i>Polydora</i> species, but related shell-boring polychaete - <i>Boccardia proboscidea</i>	<i>Haliotis sp.</i>	Simon, Bentley & Caldwell 2010
South Africa	Saldanha Bay	<i>P. hoplura</i>	<i>Crassostrea gigas</i>	David & Simon 2014
South Africa	Saldanha Bay	<i>P. hoplura</i>	<i>Crassostrea gigas</i>	David, Matthee & Simon 2014
South Africa	multiple sites	<i>P. hoplura</i>	<i>Haliotis midae</i>	Boonzaaier, Neethling, Mouton & Simon 2014

South Africa	Cape Point and Cape Agulhas: Kleinzee, Paternoster, Saldanha Bay and Port Elizabeth	<i>P. hoplura</i>	<i>Crassostrea gigas</i>	Williams, Matthee & Simon 2016
Thailand	Gulf of Thailand	spp.	<i>Molluscs living in shrimp ponds (converted mangrove)</i>	Yoshimi, Toru, & Chumpol 2007
USA	South Carolina	<i>P. ciliata</i>	<i>Crassostrea virginica</i>	Lunz 1941
USA	Connecticut	<i>P. websteri</i>	<i>Crassostrea virginica</i>	Loosanoff & Engle 1943
USA	Delaware Bay	spp.	<i>Crassostrea virginica</i>	Littlewood, Wargo & Kraeuter 1989
USA	Hawaii	<i>P. nuchalis</i>	<i>Crassostrea virginica;</i> <i>Penaeus vannamei</i>	Bailey-Brock 1990
USA	Delaware Bay	spp.	<i>Crassostrea virginica</i>	Littlewood, Wargo, Kraeuter & Watson 1992
USA	Chesapeake Bay	spp.	<i>Crassostrea gigas</i>	Burreson, Mann & Allen 1994
USA	Delaware Bay	<i>P. websteri</i>	<i>Crassostrea gigas;</i> <i>Crassostrea virginica</i>	Debrosse & Allen 1996
USA	Hawaii, shipped from Maine	Not a Polydora species, but related shell-boring polychaete - <i>Boccardia proboscidea</i>	<i>Ostrea edulis</i>	Bailey-Brock 2000
USA	Virginia	spp.	<i>Crassostrea virginica;</i> <i>Crassostrea ariakensis</i>	Calvo <i>et al.</i> 2001
USA	North Carolina	spp.	<i>Crassostrea ariakensis</i>	Bishop & Peterson 2005
USA	North Carolina	spp.	<i>Crassostrea virginica;</i> <i>Crassostrea ariakensis</i>	Bishop & Hooper 2005
USA	North Carolina	spp.	<i>Crassostrea ariakensis</i>	Grabowski <i>et al.</i> 2007
USA	Chesapeake Bay	spp.	<i>Crassostrea ariakensis;</i> <i>Crassostrea virginica</i>	McLean & Abbe 2008
USA	Maine	<i>P. websteri</i>	<i>Crassostrea virginica</i>	Brown 2012

USA	St. Charles River near the entrance of the Richibucto Estuary	<i>P. websteri</i>	<i>Crassostrea virginica</i>	Clements <i>et al.</i> 2017a
-----	--	--------------------	------------------------------	------------------------------

509

510

For Review Only

511 References Cited

- 512 Abe, H., W. Sato-Okoshi & Y. Endo. 2011. Seasonal changes of planktonic polychaete larvae
513 and chlorophyll a concentration in Onagawa Bay, northeastern Japan. *Ital. J. Zool.*
514 78(sup1), 255–266.
- 515 Alagarswami K & A. Chellam. 1976. On fouling and boring organisms and mortality of pearl
516 oysters in the farm at Veppalodai, Gulf of Mannar. *Indian J.Fish.* 23:10-22.
- 517 Alfjorden, A., M. Areskog, D. Bruno, R. Carnegie, D. Cheslett, S. Feist, S. Ford, et al. 2017.
518 New trends in important diseases affecting the culture of fish and molluscs in the ICES
519 area 2002 – 2015. International Council for the Exploration of the Sea (ICES). ICES
520 Cooperative Research Report No. 337.
- 521 Ambariyanto & R. Seed. 1991. The infestation of *Mytilus edulis* Linnaeus by *Polydora ciliata*
522 (Johnston) in the Conwy Estuary, North Wales. *J. Molluscan Stud.* 57:413–424.
- 523 Anger, K., V. Anger & E. Hagmeier. 1986. Laboratory studies on larval growth of Polydora
524 ligni, Polydora ciliata, and Pygospio elegans (Polychaeta, Spionidae). *Helgoländer
525 Meeresuntersuchungen* 40:377–395.
- 526 Bailey-Brock, J. H. & A. Ringwood. 1982. Methods for control of the mud blister worm,
527 *Polydora websteri*. Hawaiian oyster culture. *Sea Grant Quarterly* 4:6.
- 528 Bailey-Brock, J. H. 1990. *Polydora nuchalis* (Polychaeta: Spionidae), a New Hawaiian Record
529 from Aquaculture Ponds. *Pac. Sci.* 44:81–87.
- 530 Bailey-Brock, J. 2000. A new record of the polychaete *Boccardia proboscidea* (Family
531 Spionidae), imported to Hawai‘i with oysters. *Pac. Sci.* 54:27-30.
- 532 Barnabás, B., K. Jäger & A. Fehér. 2008. The effect of drought and heat stress on reproductive
533 processes in cereals. *Plant Cell Environ.* 31:11–38.

- 534 Basilio C. D., J. I. Canete & N. Rozbaczylo. 1995. *Polydora* sp. (spionidae), a polychaete borer
535 of the scallop *Argopecten purpuratus* (bivalvia: Pectinidae) valves from Tongoy Tay,
536 Chile. *Rev Biol Mar.* 30:71-77.
- 537 Benson, G.G. & A. Gyler. 1887. Report on the Hawkesbury River oyster beds. Commissioners
538 Of Fisheries. 1887. *Fisheries Of The Colony: Report Of Commissioners of Fisheries up*
539 *to 31st December, 1888.* Appendix G: Pp. 11–12. Charles Potter Govt. Pr., Sydney,
540 NSW. Pp. 73.
- 541 Bergman, K. M., R. W. Elner, & M. J. Risk. 1982. The influence of *Polydora websteri* borings
542 on the strength of the shell of the sea scallop, *Placopecten magellanicus*. *Can. J. Zool.*
543 60:2551–2556.
- 544 Bilbao, A. et al. 2011. Control of shell-boring polychaetes in *Haliotis Tuberculata Coccinea*
545 (Reeve 1846) aquaculture: Species identification and effectiveness of mebendazole. *J.*
546 *Shellfish Res.* 30:331–336.
- 547 Bishop, M. J. & P. J. Hooper. 2005. Flow, stocking density and treatment against *Polydora* spp.:
548 Influences on nursery growth and mortality of the oysters *Crassostrea virginica* and *C.*
549 *ariakensis*. *Aquaculture* 246:251–261.
- 550 Bishop, M. J. & C. H. Peterson. 2005. Constraints to *Crassostrea ariakensis* aquaculture: Season
551 and method of culture strongly influence success of grow-out. *J. Shellfish Res.* 24:995–
552 1006.
- 553 Blake, J. A. 1969a. Reproduction and larval development of *Polydora* from northern New
554 England (Polychaeta: Spionidae). *Ophelia* 7:1–63.
- 555 Blake, J. A. 1969b. Systematics and ecology of shell-boring polychaetes from New England.
556 *Integr. Comp. Biol.* 9:813–820.

- 557 Blake, J. A. & K. H. Woodwick. 1971. New species of *Polydora* (Polychaeta: Spionidae)
558 from the coast of California. *Bull. -South. Calif. Acad. Sci.* 70:72–79.
- 559 Blake, J. A. & P. L. Arnofsky. 1999. Reproduction and larval development of the spioniform
560 Polychaeta with application to systematics and phylogeny. *Hydrobiologia* 402:57–106.
- 561 Blake, J. A. 2017. Larval development of Polychaeta from the Northern California Coast.
562 fourteen additional species together with seasonality of planktic larvae over a 5-year
563 period. *J. Mar. Biol. Assoc. U. K.* 97:1081–1133.
- 564 Boonzaaier, M. K., S. Neethling, A. Mouton & C. A. Simon. 2014. Polydorid polychaetes
565 (Spionidae) on farmed and wild abalone (*Haliotis midae*) in South Africa: an
566 epidemiological survey. *Afr. J. Mar. Sci.* 36: 369–376.
- 567 Boscolo, R. & O. Giovanardi. 2002. *Polydora ciliata* shell infestation in *Tapes philippinarum*
568 Manila clam held out of the substrate in the Adriatic sea, Italy. *J. Invertebr. Pathol.*
569 79:197–198.
- 570 Boscolo R & O. Giovanardi. 2003. *Polydora ciliata* settlement on shells of the manila clam
571 *Tapes philippinarum* (Adams & Reeve). *Biologia marina mediterranea*. 10:1054-1056.
- 572 Bower, S. M., J. Blackbourn, G. R. Meyer & D. J. H. Nishimura. 1992. Diseases of cultured
573 Japanese scallops (*Patinopecten yessoensis*) in British Columbia, Canada. *Aquaculture*
574 107: 201–210.
- 575 Bower, S. M., S. E. McGladdery & I. M. Price. 1994. Synopsis of infectious diseases and
576 parasites of commercially exploited shellfish. *Annu. Rev. Fish Dis.* 4: 1–199.
- 577 Brown, Shannon W. 2012. Salinity tolerance of the oyster mudworm *Polydora websteri*. Honors
578 thesis. Honors college, University of Maine, Orono, Maine. Available at:
579 <https://digitalcommons.library.umaine.edu/honors/41>

- 580 Burreson, E. M., R. Mann & S. J. Allen. 1994. Field exposure of triploid *Crassostrea gigas* to
581 *Haplosporidium nelsoni* (MSX) and *Perkinsus marinus* (dermo) in the lower Chesapeake
582 Bay. *J. Shellfish Res.*, 13:293.
- 583 Caceres-Martinez, J., D. O. Macias-Montes & R. Vasquez-Yeomans. 1998. *Polydora* sp.
584 infestation and health of the Pacific oyster *Crassostrea gigas* cultured in Baja California,
585 NW Mexico. *J. Shellfish Res.* 17:259-264.
- 586 Calvo, G. W., M. W. Luckenbach & E. M. Burreson. 1999. A comparative field study of
587 *Crassostrea gigas* and *Crassostrea virginica* in relation to salinity in Virginia. Special
588 Reports in Applied Marine Science and Ocean Engineering (SRAMSOE) No. 349.
589 Virginia Institute of Marine Science, College of William and Mary.
- 590 Calvo, G. W., M. W. Luckenbach, S. J. Allen & E. M. Burreson 2001. A comparative
591 field study of *Crassostrea ariakensis* (Fujita 1913) and *Crassostrea virginica* (Gmelin
592 1791) in relation to salinity in Virginia. *J. Shellfish Res.* 20:221-229.
- 593 Calvo, L., B. Haskin, W. Schroer & R. Petrecca. 2014. Methods to control bio-fouling of
594 cultured eastern oysters, *Crassostrea virginica*, by the tube-building polychaete worm,
595 *Polydora cornuta*. Final report for FNE13-780. U.S. Department of Agriculture
596 Sustainable Agriculture Research and Education. Available at:
597 <https://projects.sare.org/project-reports/fne13-780/>
- 598 Chambon, C., A. Legeay, G. Durrieu, P. Gonzalez, P. Ciret & J-C. Massabuau. 2007. Influence
599 of the parasite worm *Polydora* sp. on the behaviour of the oyster *Crassostrea gigas*: a
600 study of the respiratory impact and associated oxidative stress. *Mar. Biol.* 152:329–338.
- 601 Çinar, M. E. 2013. Alien polychaete species worldwide: current status and their impacts. *J. Mar.*
602 *Biol. Assoc. U. K.* 93:1257–1278.

- 603 Clements, J. C., D. Bourque, J. McLaughlin, M. Stephenson & L. A. Comeau. 2017a. Siltation
604 increases the susceptibility of surface-cultured eastern oysters (*Crassostrea virginica*) to
605 parasitism by the mudworm *Polydora websteri*. *Aquac. Res.* 48:4707-4714.
- 606 Clements, J. C., D. Bourque, J. McLaughlin, M. Stephenson & L. A. Comeau. 2017b. Extreme
607 ocean acidification reduces the susceptibility of eastern oyster shells to a polydorid
608 parasite. *J. Fish Dis.* 40:1573–1585.
- 609 Clements, J. C., D. Bourque, J. McLaughlin, M. Stephenson & L. A. Comeau. 2018. Wanted
610 dead or alive: *Polydora websteri* recruit to both live oysters and empty shells of the
611 eastern oyster, *Crassostrea virginica*. *J. Fish Dis.* 41:855–858.
- 612 Cole, S. 2018. Mudblister worm infestation on farmed oysters along the Alabama coast. Masters
613 thesis., University of South Alabama. Available on ProQuest, number 10976739.
- 614 Cox, B., P. Kosmeyer, W. O'Connor, M. Dove & K. Johnstone. 2012. Oyster over-catch: cold
615 shock treatment. The Seafood CRC Company Ltd, the Fisheries Research and
616 Development Corporation, Port Stephens Fisheries Institute, Industry & Investment NSW
617 and Tasmanian Oyster Research Council Ltd. Project 734. Available at:
618 <http://www.frdc.com.au/Archived-Reports/FRDC%20Projects/2010-734-DLD.pdf>
- 619 Cox, J.C., 1889. Report of the Commissioners of Fisheries for the year ending 31st December
620 1889. In Commissioners Of Fisheries 1890. Charles Potter Govt. Pr., Sydney, NSW, Pp.
621 30.
- 622 Crooks, J. A. 2005. Lag times and exotic species: the ecology and management of biological
623 invasions in slow-motion1. *Écoscience*, 12:316–329.
- 624 Curtin, L. 1982. Longlines for improving oyster condition. *Catch' 82* 9(5):15. Retrieved from
625 <https://search.proquest.com/docview/13934320?accountid=14784>

- 626 Culver, C. S. & A. M. Kuris. 2000. The apparent eradication of a locally established introduced
627 marine pest. *Biol. Invasions* 2:245–253.
- 628 Da Silva, P. M., P. M. Scardua, C. B. Vieira, A. C. Alves & C. F. Dungan. 2015. Survey of
629 pathologies in *Crassostrea gasar* (Adanson, 1757) oysters from cultured and wild
630 populations in the São Francisco Estuary, Sergipe, Northeast Brazil. *J. Shellfish Res.*
631 34:289–296.
- 632 Di Salvo L. & E. Martinez. 1985. Culture of *Ostrea chilensis Philippi* 1845, in a north central
633 Chilean coastal bay. *Biologia Pesquera*. 14:16-22.
- 634 Daro, M. H. & J. Soroa Bofill. 1972. Study on the oyster culture biotope at Ostende in 1970.
635 *Aquaculture* 1:97–113.
- 636 David, A. A. & C. A. Simon. 2014. The effect of temperature on larval development of two non-
637 indigenous poecilogenous polychaetes (Annelida: Spionidae) with implications for life
638 history theory, establishment and range expansion. *J. Exp. Mar. Bio. Ecol.* 461:20–30.
- 639 David, A. A., C. A. Matthee & C. A. Simon. 2014. Poecilogony in *Polydora hoplura*
640 (Polychaeta: Spionidae) from commercially important molluscs in South Africa. *Mar.*
641 *Biol.* 161:887–898.
- 642 Debrosse, G. A., S. J. & Allen. 1996. The suitability of land-based evaluations of *Crassostrea*
643 *gigas* (Thunberg, 1793) as an indicator of performance in the field. *J. Shellfish Res.*,
644 15:291-295.
- 645 Diez, M. E., V. I. Radashevsky, J. M. Orensanz & F. Cremonte. 2011. Spionid polychaetes
646 (Annelida: Spionidae) boring into shells of molluscs of commercial interest in northern
647 Patagonia, Argentina. *Ital. J. Zool.*, 78:497–504.

- 648 Diggles, B. K. 2013. Historical epidemiology indicates water quality decline drives loss of oyster
649 (*Saccostrea glomerata*) reefs in Moreton Bay, Australia. *New Zeal. J. Mar. Fresh.*
650 47:561–581.
- 651 Dorsett, D. A. The reproduction and maintenance of *Polydora ciliata* (Johnst.) at Whitstable.
652 1961. *J. Mar. Biol. Assoc. U. K.* 41:383–396.
- 653 Dunphy, B. J., R. M. G. Wells & A. G. Jeffs. 2005. Polydorid infestation in the flat oyster,
654 *Tiostrea chilensis*: hyposaline treatment for an aquaculture candidate. *Aquac. Int.*
655 13:351–358.
- 656 Edgar, G. J. 2001. *Australian Marine Habitats In Temperate Waters*. Reed New Holland.
- 657 Eldredge, L. G. & J. D. Humphries. 1994. Perspectives in aquatic exotic species management in
658 the Pacific Islands. Pacific Science Association, South Pacific Commission.
- 659 Elston, R. A., C. A. Farley & M. L. Kent. 1986. Occurrence and significance of bonamiasis in
660 European flat oysters *Ostrea edulis* in North America. *Dis. Aquat. Org.* 2:49–54.
- 661 Ferner, M. C. & Jumars, P. A. 1999. Responses of deposit-feeding spionid polychaetes to
662 dissolved chemical cues. *J. Exp. Mar. Biol. Ecol.* 236:89–106.
- 663 Fleury P, E. Goyard & J. Mazurie et al. 2001. The assessing of Pacific oyster (*Crassostrea gigas*)
664 rearing performances by the IFREMER/REMORA network: Method and first results
665 (1993-98) in Brittany (France). *Hydrobiologia*. 465:195-208.
- 666 Fleury P, E. Le Ber, S. Claude et al. 2003. Comparison of pacific oyster (*Crassostrea gigas*)
667 rearing results (survival, growth, quality) in french farming areas, after a 10-years
668 monitoring (1993-2002) by the IFREMER/REMORA NETWORK. *J Shellfish Res.*
669 22:330.

- 670 Gabaev, D. D. 2013. Effects of fouling on the Japanese scallop *Mizuhopecten yessoensis* (Jay) in
671 Peter the Great Bay (Sea of Japan). *Oceanology* 53:183–191.
- 672 Gallo-García, M. C., M. G. Ulloa-Gómez & D. E. Godínez-Siordia. 2004. Evaluation of two
673 treatments in polychaete worm intensity associated with *Crassostrea gigas* (Thunberg,
674 1873) oyster valves. *Cienc. Mar.* 30:455–464.
- 675 Gamble, C. R. 2016. An evaluation of the floating cage system for Eastern oyster (*Crassostrea*
676 *virginica*) aquaculture production in the north-central Gulf of Mexico. Masters thesis.
677 University of Akureyri. Available at:
678 <https://pdfs.semanticscholar.org/c932/ab45f95675372813f8b921b38fa176ea3ba9.pdf>
- 679 Government of Canada, Fisheries & Ocean Services. 2017. Investigating *Polydora* outbreak in
680 New Brunswick off-bottom cultured oysters. Aquaculture Collaborative Research and
681 Development Program, Grant G-14-01-002.
- 682 Gentemann, C. L., M. R. Fewings & M. García-Reyes. 2017. Satellite sea surface temperatures
683 along the West Coast of the United States during the 2014-2016 northeast Pacific marine
684 heat wave: Coastal SSTs During “the Blob.” *Geophys. Res. Lett.* 44:312–319.
- 685 Grabowski, J. H., C. H. Peterson, M. J. Bishop & R. Conrad. 2007. The bioeconomic feasibility
686 of culturing triploid *Crassostrea ariakensis* in North Carolina. *J. Shellfish Res.* 26:529–
687 542.
- 688 Grant, J.D. 1889. Report on Georges River fisheries, Appendix A:5–15. Commissioners of
689 Fisheries 1889: Report of the Commissioners of Fisheries for the year ending 31st
690 December 1888. Charles Potter Govt. Pr., Sydney, NSW, Pp. 30.
- 691 Gryder, D. K. 2002. Control of mud blister formation in oysters. Final Report, Project FRG-00-
692 05, Virginia Fishery Resource Grant Program. Available at

- 693 http://139.70.23.11/research/units/centerspartners/map/frg/reports/docs_frg_reports/FRG-
- 694 [2000-05-Dennis-Gryder.pdf](#)
- 695 Hadiroseyan Y & D. Djokosetyanto dan Iswadi. 2007. Polychaete species infected pearl oyster
- 696 *Pinctada maxima* at Padang Cermin water, Lampung. *Jurnal akuakultur Indonesia*.
- 697 6:197-204.
- 698 Haigler, S. A. 1969. Boring mechanism of *Polydora websteri* inhabiting *Crassostrea virginica*.
- 699 *Am. Zool.* 9:821–828.
- 700 Handley, S. J. 1995. Spionid polychaetes in Pacific oysters, *Crassostrea gigas* (Thunberg) from
- 701 Admiralty Bay, Marlborough Sounds, New Zealand. *N. Z. J. Mar. Freshwater Res.*
- 702 29:305–309.
- 703 Handley a, S. J. & P. R. Bergquist. 1997. Spionid polychaete infestations of intertidal pacific
- 704 oysters *Crassostrea gigas* Thunberg) Mahurangi Harbour, northern New Zealand.
- 705 *Aquaculture* 153:191–205.
- 706 Handley, S. J. 1998. Power to the oyster: do spionid-induced shell blisters affect condition in
- 707 subtidal oysters? *J. Shellfish Res.* 17:1093–1100.
- 708 Handley, S. J. 2002. Optimizing intertidal Pacific oyster (Thunberg) culture, Houhora Harbour,
- 709 northern New Zealand. *Aquac. Res.* 33:1019–1030.
- 710 Hanley, T. C., J. W. White, C. D. Stallings & D. L. Kimbro. 2019. Environmental gradients
- 711 shape the combined effects of multiple parasites on oyster hosts in the northern Gulf of
- 712 Mexico. *Mar. Ecol. Prog. Ser.* 612:111–125.
- 713 Handlinger, J. H., M. Leonart & M. D. Powell. 2004. Development of an integrated
- 714 management program for the control of spionid mudworms in cultured abalone.
- 715 Australian Fisheries Research and Development Corporation Project No. 98/307.

- 716 Available at: [http://frdc.com.au/Archived-Reports/FRDC%20Projects/1998-307-](http://frdc.com.au/Archived-Reports/FRDC%20Projects/1998-307-DLD.pdf)
- 717 DLD.pdf
- 718 Hansen, B. W., H. H. Jakobsen, A. Andersen, R. Almeda, T. M. Pedersen, A. M. Christensen &
- 719 B. Nilsson. 2010. Swimming behavior and prey retention of the polychaete larvae
- 720 *Polydora ciliata* (Johnston). *J. Exp. Biol.* 213:3237–3246.
- 721 Hartman, O. 1940. *Boccardia proboscidea*, a new species of spionid worm from California. *J.*
- 722 *Wash. Acad. Sci.* 30:382–387.
- 723 Hatfield, P. A. 1965. *Polydora commensalis* Andrews - larval development and observations in
- 724 adults. *Biol. Bull.* 128:356–368.
- 725 Hidu, H., S. R. Chapman & W. Mook. 1988. Overwintering American oyster seed by cold humid
- 726 air storage. *J. Shellfish Res.* 7:47–50.
- 727 Hopkins, S. H. 1958. The planktonic larvae of *Polydora Websteri* Hartman (Annelida,
- 728 Polychaeta) and their settling on oysters. *Bull. Mar. Sci.* 8:268–277.
- 729 Jeffries, H. P. 1972. A stress syndrome in the hard clam, *Mercenaria mercenaria*. *J. Invertebr.*
- 730 *Pathol.* 20:242–251.
- 731 Kent, R. 1981. The effect of *Polydora ciliata* on the shell strength of *Mytilus edulis*. *ICES J.*
- 732 *Mar. Sci.* 39:252–255.
- 733 Kojima H., & M. Imajima. 1982. Burrowing polychaetes in the shells of the abalone *Haliotis*
- 734 *diversicolor aquatilis* chiefly on the species of *Polydora*. *Nihon Suisan Gakkai Shi* 48:31.
- 735 Korringa, P. 1976. Farming the flat oysters of the genus *Ostrea*: a multidisciplinary treatise
- 736 (Developments in aquaculture and fisheries science; 3), Amsterdam; Oxford: Elsevier.
- 737 Kristan, D. M. 2004. Intestinal nematode infection affects host life history and offspring
- 738 susceptibility to parasitism. *J. Anim. Ecol.* 73:227–238.

- 739 Lafferty, K. D., & A. M. Kuris. 1996. Biological control of marine pests. *Ecology* 77:1989–2000.
- 740 Lemasson, A. J., & A. M. Knights. 2019. Preferential parasitism of native oyster *Ostrea edulis*
- 741 over non-native *Magallana gigas* by a Polydorid worm. *Estuaries Coasts*.
- 742 Lie, U. 1968. A quantitative study of benthic infauna in Puget Sound, Washington, USA, in
- 743 1963-1964. *Fisk Dir. Skr. Ser. Havundersök*. 14:229–556.
- 744 Littlewood, D., R. N. Wargo & J. N. Kraeuter. 1989. Growth, mortality, MSX infection and yield
- 745 of intertidally grown *Crassostrea virginica*. *J. Shellfish Res.* 8:469.
- 746 Leonart, M., J. Handlinger & M. Powell. 2003a. Spionid mudworm infestation of farmed
- 747 abalone (*Haliotis* spp.). *Aquaculture* 221:85–96.
- 748 Leonart, M., J. Handlinger & M. Powell. 2003b. Treatment of spionid mud worm (*Boccardia*
- 749 *knoxi Rainer*) infestation of cultured abalone. *Aquaculture* 217:1–10.
- 750 Loosanoff, V. L. & J. B. Engle. 1943. *Polydora* in oysters suspended in the water. *Biol. Bull.*
- 751 85:69–78.
- 752 Lunz, G. R. 1941. *Polydora*, a pest in South Carolina oysters. Journal of the Elisha Mitchell
- 753 Scientific Society 57: 273–283.
- 754 Martinelli, J., H. Lopes, H., L. Hauser, I. Jimenez-Hidalgo, T. L. King, J. Padilla-Gamino, P.
- 755 Rawson, L. Spencer, J. Williams & C. Wood. 2019. First confirmation of the shell-boring
- 756 oyster parasite *Polydora websteri* (Polychaeta: Spionidae) in Washington State, USA. *In*
- 757 *press, Scientific Reports. PeerJ Preprint* available at:
- 758 <https://doi.org/10.7287/peerj.preprints.27621v2>
- 759 Mazurie J, J.F. Bouget, J. Barret, D. Blateau, R. Le Changour & B. Le Gall. 1995. Mussels and
- 760 oysters growth in Brest Bay, indicators of water quality and molluscs farming potential.
- 761 *Ing Eau Agric Territ.* 111-118.

- 762 McLean, R. I. & G. R. Abbe. 2008. Characteristics of *Crassostrea ariakensis* (Fujita 1913) and
763 *Crassostrea virginica* (Gmelin 1791) in the discharge area of a nuclear power plant in
764 central Chesapeake Bay. *J. Shellfish Res.* 27:517–523.
- 765 Meyer, F. P. 1991. Aquaculture disease and health management. *J. Anim. Sci.* 69:4201–4208.
- 766 Moreno, R. A., P. E. Neill & N. Rozbaczylo. 2006. Native and non-indigenous boring
767 polychaetes in Chile: a threat to native and commercial mollusc species. *Rev. Chil. Hist.*
768 *Nat.* 79:263-278.
- 769 Morse, D. L., P. D. Rawson & J. N. Kraeuter. 2015. Mud blister worms and oyster aquaculture.
770 *Maine Sea Grant Publications*. 46. Available at:
771 https://digitalcommons.library.umaine.edu/seagrant_pub/46/
- 772 Mortensen, S., T. Van der Meeren, A. Fosshagen, I. Hernar, L. Harkestad, and L. Torkildsen. &
773 Ø. Bergh. 2000. Mortality of scallop spat in cultivation, infested with tube dwelling
774 bristle worms, *Polydora* sp. *Aquac. Int.* 8:267–271.
- 775 Nascimento, A. I. 1983. Oyster Culture in Brazil: Problems and Perspectives. *Ciencia e Cultura*
776 (Sao Paulo). *Sao Paulo* 35:871-876.
- 777 Nell, J. A. 1993. Farming the Sydney rock oyster (*Saccostrea commercialis*) in Australia. *Rev.*
778 *Fish. Sci.* 1:97–120.
- 779 Nell, J. A. 2001. The history of oyster farming in Australia. *Mar. Fish. Rev.* 63:14–25.
- 780 Nell, J. 2007. Controlling mudworm in oysters. New South Wales Department of Primary
781 Industry Primefact 590. Available at:
782 [https://www.dpi.nsw.gov.au/__data/assets/pdf_file/0010/637633/Controlling-mudworm-](https://www.dpi.nsw.gov.au/__data/assets/pdf_file/0010/637633/Controlling-mudworm-in-oysters.pdf)
783 in-oysters.pdf

- 784 Nel, R., P. S. Coetzee & G. Van Niekerk. 1996. The evaluation of two treatments to reduce mud
785 worm (*Polydora hoplura* Claparede) infestation in commercially reared oysters
786 (*Crassostrea gigas* Thunberg). *Aquaculture* 141:31–39.
- 787 Ogburn, D. M., I. White & D. P. Mcphee. 2007. The disappearance of oyster reefs from eastern
788 Australian estuaries—Impact of colonial settlement or mudworm invasion? *Coast.
789 Manage.* 35:271–287.
- 790 Ogburn, D. M. 2011. The NSW oyster industry: A risk indicator of sustainable coastal policy and
791 practice. PhD Dissertation. The Australian National University. Available at:
792 <http://doi.org/10.25911/5d7a266d782dc>
- 793 Omel'yanenko, V. A., Kulikova, V. A. & Pogodin, A. G. 2004. The meroplankton of Amurskii
794 Bay (Peter the Great Bay, Sea of Japan). *Russ. J. Mar. Biol.* 30:159–174.
- 795 Orth, R. J. 1971. Observations on the planktonic larvae of *Polydora ligni* Webster (Polychaeta:
796 Spionidae) in the York River, Virginia. *Chesapeake Science* 12:121–124.
- 797 Oyarzun, F. X., Mahon, A. R., Swalla, B. J. & Halanych, K. M. 2011. Phylogeography and
798 reproductive variation of the poecilogenous polychaete *Boccardia proboscidea*
799 (Annelida: Spionidae) along the West Coast of North America. *Evol. Dev.* 13:489–503.
- 800 Paladini, G., M. Longshaw, A. Gustinelli & A. P. Shinn. 2017. Parasitic diseases in aquaculture:
801 Their biology, diagnosis and control. In B. A. Austin, & A. Newaj-Fyzul (Eds.),
802 Diagnosis and control of diseases of fish and shellfish (1st ed., pp. 37–107). Chichester
803 (UK): John Wiley & Sons Ltd.
- 804 Petersen, F. S. 2016. Addressing obstacles to developing oyster culture in Hawai'i. Masters
805 thesis. University of Hawai'i at Hilo. Available at: <http://hdl.handle.net/10790/2948>

- 806 Pregenzer, C. 1983. Survey of metazoan symbionts of *Mytilus edulis* (Mollusca: Pelecypoda) in
807 southern Australia. *Mar. Freshwater Res.* 34:387–396.
- 808 Quinan, J. 1883. Report on home fisheries, for February, 1883. Commissioners of fisheries,
809 1883. Fisheries of the colony: Report of commissioners of fisheries for year 1883.
810 Appendix T: Pp. 77–86. Charles Potter Govt. Pr., Sydney, NSW, Pp. 100.
- 811 Quinan, J. 1884. Report On Home Fisheries, For February, 1884. Commissioners of fisheries,
812 1884. Fisheries of the colony: Report of commissioners of fisheries for year 1884.
813 Appendix I: Pp. 11–12. Charles Potter Govt. Pr., Sydney, Nsw, Pp. 100.
- 814 Radashevsky, V. I., P. C. Lana & R. C. Nalessio. 2006. Morphology and biology of *Polydora*
815 species (Polychaeta: Spionidae) boring into oyster shells in South America, with the
816 description of a new species. *Zootaxa* 1353:1–37.
- 817 Radashevsky, V. I. & V. V. Pankova. 2006. The morphology of two sibling sympatric *Polydora*
818 species (Polychaeta: Spionidae) from the Sea of Japan. *J. Mar. Biol. Assoc. U. K.*
819 86:245–252.
- 820 Radashevsky, V. I. & Migotto, A. E. 2017. First report of the polychaete *Polydora hoplura*
821 (Annelida: Spionidae) from North and South America and Asian Pacific. *Mar. Biodivers.*
822 47:859–868.
- 823 Read, G. B. 2010. Comparison and history of *Polydora websteri* and *P. haswelli* (Polychaeta:
824 Spionidae) as mud-blister worms in New Zealand shellfish. *N. Z. J. Mar. Freshwater Res.*
825 44:83–100.
- 826 Riascos, J. M., O. Heilmayer, M. E. Oliva, J. Laudien & W. E. Arntz. 2008. Infestation of the
827 surf clam *Mesodesma donacium* by the spionid polychaete *Polydora bioccipitalis*. *J. Sea*
828 *Res.* 59:217–227.

- 829 Riascos, J. M., N. Guzmán, J. Laudien, M. E. Oliva, O. Heilmayer & L. Ortlieb. 2009. Long-
830 term parasitic association between the boring polychaete *Polydora bioccipitalis* and
831 *Mesodesma donacium*. *Dis. Aquat. Organ.* 85:209–215.
- 832 Robert R., M. Borel, Y. Pichot & G. Trut. 1991. Growth and mortality of the European oyster
833 *Ostrea edulis* in the Bay of Arcachon (France). *Aquat. living resour.* 4:265-274.
- 834 Ropert M., S. Pien, C. Mary & B. Bouchaud. 2007. REMONOR: Results of the year 2006
835 mortality assessment, growth and quality of *Crassostrea gigas* oysters. Ifremer, Plouzane
836 (France).[np].1 Nov 2007.
- 837 Royer, J., M. Ropert, M. Mathieu & K. Costil. 2006a. Presence of spionid worms and other
838 epibionts in Pacific oysters (*Crassostrea gigas*) cultured in Normandy, France.
839 *Aquaculture* 253:461–474.
- 840 Sabry, R. C., P. M. da Silva, T. C. Vasconcelos Gesteira, V. de Almeida Pontinha & A. R.
841 Magenta Magalhães. 2011. Pathological study of oysters *Crassostrea gigas* from culture
842 and *C. rhizophorae* from natural stock of Santa Catarina Island, SC, Brazil. *Aquaculture*
843 320:43–50
- 844 Sato-Okashi W, Y. Sugawara & T. Nomura. 1990. Reproduction of the boring polychaete
845 *Polydora variegata* inhabiting scallops in Abashiri Bay, North Japan. *Mar. Biol.* 104:61-
846 66.
- 847 Sato-Okoshi, W. & K. Okoshi. 1997. Survey of the genera *Polydora*, *Boccardiella* and
848 *Boccardia* (Polychaeta, Spionidae) in Barkley Sound (Vancouver Island, Canada), with
849 special reference to boring activity. *Bull. Mar. Sci.* 60:482–493.

- 850 Sato-Okoshi, W., K. Okoshi & J. Shaw. 2008. Polydorid species (Polychaeta: Spionidae) in
851 south-western Australian waters with special reference to *Polydora uncinata* and
852 *Boccardia knoxi*. *J. Mar. Biol. Assoc. U. K.* 88:491–501.
- 853 Sato-Okoshi, W. & H. Abe. 2012. Morphological and molecular sequence analysis of the
854 harmful shell boring species of *Polydora* (Polychaeta: Spionidae) from Japan and
855 Australia. *Aquaculture* 368-369:40–47.
- 856 Sato-Okoshi, W., K. Okoshi, B.-S. Koh, Y.-H. Kim & J.-S. Hong. 2012. Polydorid species
857 (Polychaeta: Spionidae) associated with commercially important mollusk shells in
858 Korean waters. *Aquaculture* 350-353:82–90.
- 859 Sato-Okoshi, W., Okoshi, K., Abe, H. & Li, J.-Y. 2013. Polydorid species (Polychaeta,
860 Spionidae) associated with commercially important mollusk shells from eastern China.
861 *Aquaculture* 406-407:153–159.
- 862 Schleyer, M. H. 1991. Shell-borers in the oyster, *Striostrea margaritacea*: Pests or symbionts?
863 *Symbiosis* 10:135–144.
- 864 Shinn, A. P., J. Pratoomyot, J. E. Bron, G. Paladini, E. E. Brooker & A. J. Brooker. 2015.
865 Economic costs of protistan and metazoan parasites to global mariculture. *Parasitology*
866 142:196–270.
- 867 Seaman, M. N. L. 1991. Survival and aspects of metabolism in oysters, *Crassostrea gigas*,
868 during and after prolonged air storage. *Aquaculture* 93:389–395.
- 869 Silina, A. V. 2006. Tumor-like formations on the shells of Japanese scallops *Patinopecten*
870 *yessoensis* (Jay). *Mar. Biol.* 148, 833–840.
- 871 Simon, C. A., A. Ludford & S. Wynne. 2006. Spionid polychaetes infesting cultured abalone
872 *Haliotis midae* in South Africa. *Afr. J. Mar. Sci.* 28:167–171.

- 873 Simon, C. A. 2011. *Polydora* and *Dipolydora* (Polychaeta: Spionidae) associated with molluscs
874 on the south coast of South Africa, with descriptions of two new species. *Afr. Invertebr.*
875 52:39–50.
- 876 Simon, C. A., M. G. Bentley & G. S. Caldwell. 2010. 2,4-Decadienol: Exploring a novel
877 approach for the control of polychaete pests on cultured abalone. *Aquaculture* 310:52–60.
- 878 Simon, C. A. & W. Sato-Okoshi. 2015. Polydorid polychaetes on farmed molluscs: distribution,
879 spread and factors contributing to their success. *Aquacult. Environ. Interact.* 7:147–166.
- 880 Smith, G. S. 1981. Southern Queensland's oyster industry. *J. Roy. His. Soc. Qld.* 11:45–58.
- 881 Steele, S. & M. F. Mulcahy. 1999. Gametogenesis of the oyster *Crassostrea gigas* in southern
882 Ireland. *J. Mar. Biol. Assoc. U. K.* 79:673–686.
- 883 Steele, S. & M. F. Mulcahy. 2001. Impact of the copepod *Mytilicola orientalis* on the Pacific
884 oyster *Crassostrea gigas* in Ireland. *Dis. Aquat. Organ.* 47:145–149.
- 885 Stephen, D. 1978. Mud blister formation by *Polydora ciliata* in the Indian backwater oyster
886 *Crassostrea madrasensis* (Preston). *Aquaculture* 13:347–350.
- 887 Teramoto, W., W. Sato-Okoshi, H. Abe, G. Nishitani & Y. Endo. 2013. Morphology, 18S rRNA
888 gene sequence and life history of a new *Polydora* species (Polychaeta: Spionidae) from
889 northeastern Japan. *Aquat. Biol.* 18:31–45.
- 890 U.S. Department of Agriculture. 2019. U.S. National List Of Reportable Animal Diseases, 2019.
891 Accessed November 2017. Available at:
892 https://www.aphis.usda.gov/animal_health/nahrs/downloads/2019_nahrs_dz_list.pdf
- 893 U.S. Department of the Interior & U.S. Geological Survey. n.d. Nonindigenous Aquatic Species
894 Search by State Tool. Accessed November 2019 at <https://nas.er.usgs.gov/>.

- 895 Victorian Fisheries Authority. 2015. Abalone Aquaculture Translocation Protocol. VFA.
- 896 Accessed November 2019, available at: <https://vfa.vic.gov.au/operational-policy/moving-and-stocking-live-aquatic-organisms/abalone-aquaculture-translocation-protocol>
- 897
- 898 Wada, Y. & A. Masuda. 1997. On the depth variation in occurrence patterns of *Polydora* sp. in
- 899 shells of cultured pearl oyster, *Pinctada fucata Martensii*. *Bull. Ehime Prefect. Fish. Exp. Stn.*
- 900
- 901 Walton, W. C., J. E. Davis, G. I. Chaplin, F. Scott Rikard, T. R. Hanson, P. J. Waters & D.
- 902 Ladon Swann. 2012. Timely information: off-bottom oyster farming. The Alabama
- 903 Cooperative Extension System. Available at:
- 904 <http://agrilife.org/fisheries/files/2013/09/Off-Bottom-Oyster-Farming.pdf>
- 905 Wargo, R. N. & S. E. Ford. 1993. The effect of shell infestation by *Polydora* sp. and infection by
- 906 *Haplosporidium nelsoni* (MSX) on the tissue condition of oysters, *Crassostrea virginica*.
- 907 *Estuaries* 16:229.
- 908 Washington Sea Grant. 2015. Shellfish aquaculture in Washington State. Final report to the
- 909 Washington State Legislature. 84 p. Available at:
- 910 <https://wsg.washington.edu/wordpress/wp-content/uploads/Shellfish-Aquaculture-Washington-State.pdf>
- 911
- 912 Washington State Department of Fish & Wildlife. ND. Shellfish import permit supplemental
- 913 information. Accessed March 2019. Available
- 914 at: <https://wdfw.wa.gov/sites/default/files/2019-03/ImportPermitExplanations.pdf>.
- 915 Washington State Department of Fish & Wildlife. 2019. WDFW 2019 Shellfish Import Approval
- 916 Requirements. Accessed March 2019. Available at:

- 917 https://wdfw.wa.gov/licensing/shellfish_import_transfer/wdfw_shellfish_import_guidelines_final_12122019.pdf.
- 918
- 919 Whitelegge, T. 1890. Report on the worm disease affecting the oysters on the coast of New
920 South Wales. *Records of the Australian Museum* 1(2):41–54, plates iii-vi.
- 921 Williams, L., C. A. Mathee. & C. A. Simon. 2016. Dispersal and genetic structure of *Boccardia*
922 *polybranchia* and *Polydora hoplura* (Annelida: Spionidae) in South Africa and their
923 implications for aquaculture. *Aquaculture* 465:235–244.
- 924 Wisely, B., J. E. Holliday & B. L. Reid. 1979. Experimental deepwater culture of the Sydney
925 rock oyster (*Crassostrea commercialis*): III. Raft cultivation of trayed oysters.
926 *Aquaculture* 17:25–32.
- 927 Wilson, D. P. 1928. The larvae of *Polydora ciliata* Johnston and *Polydora hoplura* Claparede. *J.
928 Mar. Biol. Assoc. U. K.* 15:567–603.
- 929 Yoshimi, F., S. Toru & S. Chumpol. 2007. Diversity and Community Structure of Macrofauna
930 Fauna in Shrimp Aquaculture Ponds of the Gulf of Thailand. *Jpn. Agric. Res. Q.* 41:163–
931 172.
- 932 Zottoli, R. A. & M. R. Carriker. 1974. Burrow morphology, tube formation, and
933 microarchitecture of shell dissolution by the spionid polychaete *Polydora websteri*. *Mar.
934 Biol.* 27:307–316.
- 935 Zuniga G., W. Zurburg & E. Zamora. 1998. Initiation of oyster culture on the pacific coast of
936 Costa Rica. *Proc Gulf Caribb Fish Inst.* 50:612-623.
- 937

938 **Figures Legends**

939 **Figure 1:** The percentage of shellfish produced by value in 2015 in each Washington State
940 Dept. of Fish and Wildlife aquaculture area, where NPS=North Puget Sound, CPS=Central Puget
941 Sound, SPS=South Puget Sound, HC=Hood Canal, SJF=Strait of Juan de Fuca, GH=Grays
942 Harbor, and WB=Willapa Bay. Inlay: locations in South Puget Sound (SPS), Oakland Bay and
943 Totten Inlet, where *Polydora* spp. were positively identified in 2017.

944 **Figure 2.** A. *Crassostrea gigas* valve with three active *Polydora* burrows (red arrows indicate
945 entry points), B. *Crassostrea virginica* valve with many burrows, and C. an exposed u-shaped
946 burrow (red arrow) occupied by a shell-boring polychaete. Oysters were sampled from Puget
947 Sound, WA in 2017 (Martinelli *et al.* 2019). Images courtesy of Julieta Martinelli and Heather
948 Lopes.

949 **Figure 3.** *Polydora websteri* found in *Crassostrea gigas* valve in Puget Sound, WA in 2017
950 (Martinelli *et al.* 2019). Image courtesy of Heather Lopes and Julieta Martinelli.

951 **Figure 4:** Phylogeny of shell-boring polychaete worms using 18S1 rRNA sequences extracted
952 from *Crassostrea gigas* oysters collected in South Puget Sound, Washington in 2017. Trees were
953 constructed using maximum likelihood estimates based on Kimura 2-parameter distances.
954 Individuals labeled with OAK and TOT were collected in Oakland Bay and Totten Inlet,
955 respectively. Reproduced from Martinelli *et al.* 2019.

956 **Fig. 5.** Phylogeny of shell-boring polychaete worms using mtCOI rRNA sequences extracted
957 from *Crassostrea gigas* oysters collected in South Puget Sound, Washington in 2017. Trees were
958 constructed using maximum likelihood estimates based on Kimura 2-parameter distances.
959 Individuals labeled with OAK and TOT were collected in Oakland Bay and Totten Inlet,
960 respectively. Reproduced from Martinelli *et al.* 2019.


Figure 1


Figure 2

Figure 3

View Only


Figure 4

Figure 5