



University of Washington Libraries
Interlibrary Loan and Document Delivery Services
Box 352900 - Seattle, WA 98195-2900
(206) 543-1878
interlib@uw.edu
WAU / WAUWAS / RAPID:WAU

8/7/2024 10:53:16 AM

Document Delivery Scan

Seattle

ILLiad TN: 2282132



Loansome Doc:



Customer Reference:

Billing Category:

Needed By: 09/18/2024

Maximum Cost: n/a

NOTICE OF COPYRIGHT

The document above is being supplied to you in accordance with United States Copyright Law (Title 17 US Code). It is intended only for your personal research or instructional use.

The document may not be posted to the web or retransmitted in electronic form.

Distribution or copying in any form requires the prior express written consent of the copyright owner and payment of royalties. Infringement of copyright law may subject the violator to civil fine and/or criminal prosecution.

Special Instructions:

Notes/Alternate Delivery:

Location: Health Sciences Library -
Serials

Call #: QH431 .B55

Work Order Location: s

Journal Title: Biochemical Genetics

Volume: 25

Issue: 7-8

Month/Year: , 1987

Pages: 535-541

Article Author: Cheng, Cheng M. -L.

Article Title: Cryopreservation of erythrocytes in small aliquots for isozyme electrophoresis

ISSN: 0006-2928

OCLC #: info:doi/10.1007/BF00554355

Email: lhs3@uw.edu

EMAIL: LHS3@UW.EDU

ILLiad

Cryopreservation of Erythrocytes in Small Aliquots for Isozyme Electrophoresis

M.-L. Cheng^{1,2} and J. L. VandeBerg¹

Received 7 Jan. 1987—Final 25 Mar. 1987

INTRODUCTION

A standard technique developed for preserving erythrocytes at blood banks (see Walter, 1984, for review) involves a piece of tubing as an integral part of the donor blood bag made of polyvinyl chloride (PVC). The tubing is filled with erythrocytes suspended in a cryopreservative and heat-sealed into small segments, each of which contains an aliquot of the sample. The samples in the segments are used for antigenic typing, cross-matching, and other assays. A variation of the tubing storage method was developed by Cheng *et al.* (1986) for storing large numbers of serum or plasma samples frozen in small aliquots. In addition to eliminating repeated freezing and thawing of samples, this method is particularly convenient for cataloging and retrieving samples, prolongs the shelf life of samples, and is economical in the use of freezer space.

We have examined the feasibility of storing cryopreserved erythrocytes by the storage method of Cheng *et al.* (1986) and compared the efficacy of two different freezing solutions. One of these is a citrate–ethylene glycol solution used by one of us (JLV) since 1970 for preserving erythrocytes from many species (VandeBerg and Johnston, 1977), and the other is a buffered glycerol solution (American Association of Blood Banks, 1974) that we now use in preference to the citrate–ethylene glycol solution for some (but not all) species. This communication presents the results obtained from erythrocytes

This work was supported by NIH Grants HL28972 and RR01124.

¹ Department of Genetics, Southwest Foundation for Biomedical Research, P.O. Box 28147, San Antonio, Texas 78284.

² To whom correspondence should be addressed at Special Chemistry Development, Beckman Instruments, Inc., Mail Code W-151, 200 South Kraemer Boulevard, Brea, California 92621.

of two primate species, the olive baboon (*Papio hamadryas anubis*; see VandeBerg and Cheng, 1986) and the rhesus macaque (*Macaca mulatta*).

MATERIALS AND METHODS

Blood Sample Collection. Blood samples from olive baboons and rhesus macaques were collected into Vacutainers containing acid-citrate-dextrose (Becton-Dickinson and Co., Rutherford, N.J.). Plasma was separated from erythrocytes by centrifugation and removed. Erythrocytes were washed in phosphate-buffered saline (PBS; 10 mM sodium phosphate, 0.15 M NaCl, pH 7.4) several times until no hemolysis was visible in the supernatant.

Erythrocyte Preservation. Washed erythrocytes were preserved in the following two freezing solutions: (1) citrate ethylene glycol—200 mM trisodium citrate and 400 ml ethylene glycol (certified grade, Fisher Scientific Co.) in 1000 ml, pH 8.4; and (2) buffered glycerol—33 mM sodium phosphate (monobasic), 33 mM sodium phosphate (dibasic), 106 mM tripotassium phosphate, and 400 ml glycerol in 1000 ml. One volume of packed erythrocytes was mixed with 2 vol of freezing solution and incubated at 4°C for 30–60 min. The Tygon tubing (i.d. 3.97 mm, o.d. 5.56 mm; R-3603, Norton Co., Akron, Ohio) was cut into 20-cm lengths, and each was attached to a 5-ml disposable syringe. The erythrocyte suspension was mixed thoroughly by gently inverting each tube several times before 4 ml of the mixture was drawn into a piece of tubing. Each piece of tubing containing erythrocytes was then heat-sealed into approximately 1.5-cm intervals with a Hematron dielectric sealer. Each tubing segment contained approximately 250 μ l of the erythrocyte suspension. Each series of attached tubing segments was inserted into a protective sleeve—a cellulose acetate butyrate tube (CAB tube; i.d. 0.95 cm, o.d. 1.11 cm., length 50 cm; Thermoplastics Inc., Wayne, N.J.). Both ends of the CAB tube were capped with Capsicles (Intermed, Denmark). To reduce friction between the Tygon tubing and the CAB tube, the inner wall of the CAB tube was prelubricated with mineral oil. After an inventory number had been assigned to each specimen and labels affixed on the CAB tube near both ends as well as on the Capsicles, each CAB tube was placed in a three-tiered aluminum rack (Cheng *et al.*, 1986) in a 9-ft³ constant-temperature chest freezer (Sears), maintained at -20°C.

Recovery of Erythrocytes for Isozyme Electrophoresis. Since neither ethylene glycol (VandeBerg and Johnston, 1977) nor glycerol (see Results and Discussion) affects electrophoretic mobility or histochemical staining procedures, it is not necessary to remove either cryoprotectant from erythrocytes. Each Tygon tubing segment was severed at the seal, placed in a prelabeled vessel (e.g., microfuge tube), and warmed to room temperature. One corner of the tubing segment was cut open with an incision scissor and centrifuged for

10 sec in a Beckman microfuge to sediment erythrocytes. The supernatant was removed by aspiration, and the erythrocytes were lysed in deionized water or appropriate lysing solution for electrophoresis.

Assessment of Hemolysis of Cryopreserved Erythrocytes. The extent of hemolysis of cryopreserved erythrocytes during storage was evaluated by the amount of cyanmethemoglobin present (Eilers, 1967) in the supernatant thawing (freezing) solution in comparison with that in the remaining intact erythrocytes at intervals of 3, 6, and 12 months. The hemoglobin assay kit was obtained from Sigma Chemical Co. (St. Louis, Mo.). Drabkin's solution and cyanmethemoglobin standard solution were prepared according to the manufacturer's instructions. Tygon tubing segments containing cryopreserved erythrocytes were warmed for 1 min in a 37°C water bath. The contents of two segments from each animal cryopreserved in the same freezing solution were pooled and spun for 5 min at 15,000g in a refrigerated centrifuge. The supernatant was removed and transferred into a borosilicate disposable culture tube (75 × 12 mm). Ten microliters of the supernatant was mixed with 990 μ l of Drabkin's solution in a borosilicate disposable culture tube. The packed erythrocyte pellet was lysed in 1 ml of Drabkin's solution in a borosilicate disposable culture tube. A 10- μ l aliquot of the lysate was mixed with 990 μ l of Drabkin's solution in a separate tube. Further dilution of this lysate in Drabkin's solution was frequently needed for accurate spectrophotometric measurement of cyanmethemoglobin in the solution. The tubes were allowed to stand at room temperature for 15 min before being read spectrophotometrically at a wavelength of 540 nm. The absorbance values were then converted to actual total amounts of hemoglobin as measured by cyanmethemoglobin present in the thawing supernatant and in the lysate of intact erythrocytes. The proportion of erythrocytes hemolyzed during storage in Tygon tubing segments was computed as $[T/(P + T)]$, where T is the amount of cyanmethemoglobin in the thawing supernatant, and P is the amount of cyanmethemoglobin in the lysate of intact erythrocytes. Comparison of the effects of the freezing solutions on the survival of erythrocytes of baboons or rhesus macaques over a period of 1 years was performed by analysis of variance.

Electrophoresis and Enzyme Staining Techniques. Lysates of cryopreserved erythrocytes and of freshly drawn blood were compared electrophoretically. Baboon erythrocyte adenine phosphoribosyl transferase (APRT; EC 2.4.2.7) isozymes were resolved following the method of Mowbray *et al.* (1972) using starch gel electrophoresis and stained with a solution containing 14 C-labeled adenine to develop an autoradiogram. Glucose phosphate isomerase (GPI; EC 5.3.1.9) isozymes of baboon erythrocytes also were resolved by starch gel electrophoresis and stained by the procedure of Harris and

Hopkinson (1976). However, the pH of the gel and bridge buffers was adjusted to 8.9 with NaOH to increase the anodal mobility of the baboon GPI isozymes. Erythrocyte mannose phosphate isomerase (MPI; EC 5.3.1.8) isozymes were separated by cellulose acetate electrophoresis in Helena Titan III plates (Helena Laboratories, Beaumont, Tex.) in 0.025 M Tris–0.192 M glycine, pH 8.35, and stained according to Harris and Hopkinson (1976).

RESULTS AND DISCUSSION

Hemolysis of Cryopreserved Erythrocytes. Table I shows the percentages of hemolysis of erythrocytes of four baboons and four rhesus macaques cryopreserved in citrate–ethylene glycol solution or in buffered glycerol solution and stored in small aliquots in heat-sealed Tygon tubing segments. Analysis of variance indicated significant differences in the extent of hemolysis of erythrocytes due to (1) different freezing solutions; (2) the species origin of erythrocytes; and (3) the duration of storage. Both rhesus macaque and baboon erythrocytes cryopreserved in buffered glycerol had substantially less hemolysis than those cryopreserved in citrate–ethylene glycol ($F_{1,12} = 450.28$; $P < 0.001$). Consistently more hemolysis occurred in baboon erythrocytes than in rhesus erythrocytes ($F_{1,12} = 205.28$; $P < 0.001$); however, the magnitude of difference was much greater for erythrocytes cryopreserved in citrate–ethylene glycol than for those cryopreserved in buffered glycerol. For instance, at the end of the 12-month period, the mean percentage hemolysis was 31 for rhesus and 74 for baboon erythrocytes cryopreserved in citrate–ethylene glycol. In contrast, the mean percentage hemolysis was 15 for rhesus

Table I. Percentages of Hemolysis of Erythrocytes of *Macaca mulatta* (MMA) and *Papio hamadryas anubis* (PHA) Cryopreserved in Citrate–Ethylene Glycol or Buffered Glycerol

	Ethylene Glycol			Glycerol		
	3 mo ^a	6 mo	12 mo	3 mo	6 mo	12 mo
MMA 1	11.5	32.4	32.6	7.5	15.0	16.0
MMA 2	21.6	26.2	26.6	2.7	12.9	14.2
MMA 3	21.2	27.0	28.5	2.5	6.3	6.6
MMA 4	25.4	33.6	36.0	11.4	20.7	21.4
Mean	19.9	29.8	30.9	6.0	13.7	14.6
PHA 1	66.3	68.4	68.1	10.6	13.4	14.4
PHA 2	50.9	77.2	76.8	7.8	23.4	24.1
PHA 3	56.1	78.2	78.5	6.7	19.8	21.0
PHA 4	54.6	72.5	73.0	7.9	20.1	20.9
Mean	57.0	74.1	74.1	8.3	19.2	20.1

^aMonths.

and 20 for baboon erythrocytes cryopreserved in buffered glycerol. The difference in percentage hemolysis for both species after 6 and 12 months of storage was not significant ($F_{1,24} = 0.214$; $P > 0.05$), but the difference between 3 months and the weighted mean of 6 months and 12 months was highly significant ($F_{1,24} = 40.6$; $P < 0.001$). Increasing hemolysis in the first 6 months may be due to initial shocks (e.g., changes in osmosis of erythrocyte membranes in freezing solutions, changes in temperature due to freezing, trauma induced by the heat-sealing process). After 6 months of storage, the rate of hemolysis of erythrocytes of both species cryopreserved in either freezing solution was negligible.

Isozyme Patterns. The electrophoretic zymograms of fresh erythrocytes and erythrocytes cryopreserved for 18 months or longer in buffered glycerol for three representative baboon erythrocyte enzyme systems, APRT, GPI, and MPI, are shown in Fig. 1. Comparable enzyme activities, judged from histochemical staining intensity and electrophoretic mobilities, were observed for the two types of erythrocytes. Furthermore, there was no increase in the number of subbands, indicating that enzymes were stable in erythrocytes stored in buffered glycerol.

VandeBerg and Johnston (1977) reported that ethylene glycol has no apparent effect on the mobility or activity of red kangaroo or rhesus macaque erythrocyte enzymes subjected to electrophoresis in starch or cellulose acetate media. We have confirmed this observation in rhesus macaques (data not shown) and extended it to baboons.

Effect of PVC on Erythrocytes. Tygon tubing is made of PVC and contains approximately 40% (w/w) of the plasticizer, di(2-ethylhexyl)phthalate (DEHP). DEHP is leached from the tubing and enters the red cells during storage. The potential toxicity of DEHP in laboratory animals has been of considerable concern. However, it also has been shown that DEHP facilitates the survival of erythrocytes during storage at low temperatures by decreasing their osmotic fragility (Horowitz *et al.*, 1985).

General Discussion. Our primary objectives for developing this storage method were to prolong the shelf life of stored erythrocytes, to protect the structural integrity and activity of enzymes, to eliminate contamination of erythrocyte samples, to ease the complexity in sample cataloging, and to reduce the labor involved in retrieving samples from and returning samples to freezers.

Our experimental results indicate that the stability of erythrocytes in citrate-ethylene glycol or buffered glycerol is species specific. This conclusion is reinforced by long-term observations on many thousands of cryopreserved erythrocyte samples in our laboratory. In fact, we initially were prompted to explore the use of glycerol as a cryopreservative when we noticed that




Fig. 1. Zymograms of three baboon erythrocyte enzymes. The anode is at the top of each zymogram. In all three zymograms, channels 1–3 contained lysates of glycerized baboon erythrocyte samples stored in small aliquots in Tygon tubing segments kept in an 18°C freezer, and channels 4–6 contained lysates of freshly collected baboon erythrocyte samples. A shows an autoradiogram of adenine phosphoribosyl transferase (APRT) isozymes separated by starch gel electrophoresis. The phenotypes, from left to right, are A, AB, AB, A, A, and AB. B illustrates glucose phosphate isomerase (GPI) isozymes also separated by starch gel electrophoresis. The phenotypes, from left to right, are B, AB, A, B, AB, and A. C shows mannose phosphate isomerase (MPI) isozymes separated by cellulose acetate electrophoresis. The phenotypes, from left to right, are BC, B, AB, BC, B, and AB. No differences in the electrophoretic mobilities or staining activities of the enzymes derived from stored and fresh erythrocytes were detected. Hb, hemoglobin.

erythrocytes from rats, mice, and rhesus macaques had undergone very little hemolysis during 10 years of storage in citrate-ethylene glycol, whereas erythrocytes of baboons [not one of the species investigated by VandeBerg and Johnston (1977)] were quite labile when stored under identical conditions. When we explored the use of glycerol as an alternative, we observed that erythrocytes of squirrel monkeys formed gummy aggregates when mixed with the buffered glycerol solution but could be adequately preserved in the citrate-ethylene glycol solution. Clearly, it is important to identify a cryoprotectant that is compatible with the erythrocytes to be preserved in preliminary experimental trials.

Based on our experience, we suggest several other precautions, although none has been rigorously evaluated in controlled experiments: (1) fragile erythrocytes in freshly collected blood should be removed by several washes in PBS before mixing with cryopreservative; (2) the ratio of packed erythrocytes to freezing solution should be no less than 1:1.5; and (3) the buffered glycerol- or citrate-ethylene glycol-preserved erythrocytes should be maintained above the freezing temperature of the cryopreservative, because the formation of ice can damage erythrocyte membranes and cause hemolysis; a household freezer maintained at -20°C is satisfactory.

ACKNOWLEDGMENTS

The authors thank J. A. Hilburn, M. J. Aivaliotis, and "Skip" Landry for technical assistance and Dr. W. R. Lowe for statistical analyses.

REFERENCES

- American Association of Blood Banks (1974). *Technical Methods and Procedures of the American Association of Blood Banks*, 6th ed., American Association of Blood Banks, Washington, D.C., pp. 342-344.
- Cheng, M.-L., Woodford, S. C., Hilburn, J. L., and VandeBerg, J. L. (1986). A novel system for storage of sera frozen in small aliquots. *J. Biochem. Biophys. Methods* **13**:47.
- Eilers, R. J. (1967). Notification of final adoption of an international method and standard solution for hemoglobinometry specifications for preparation of standard solution. *Am. J. Clin. Pathol.* **47**:212.
- Harris, H., and Hopkinson, D. A. (1976). *Handbook of Enzyme Electrophoresis in Human Genetics*, North-Holland, Amsterdam.
- Horowitz, B., Stryker, M., Waldman, A. A., Woods, K. R., Gass, J. D., and Drago, J. (1985). Stabilization of red blood cells by the plasticizer, diethylhexylphthalate. *Vox Sang.* **48**:150.
- Mowbray, S., Watson, B., and Harris H. (1972). A search for electrophoretic variants of human adenine phosphoribosyl transferase. *Ann. Hum. Genet.* **36**:153.
- VandeBerg, J. L., and Cheng, M.-L. (1986). The genetics of baboons in biomedical research. In Else, J. G., and Lee, P. C. (eds.), *Primate Evolution* (Selected Proceedings of the Tenth Congress of the International Primatological Society, Nairobi, Kenya, 1984, Vol. 1), Cambridge University Press, London, New York, pp. 317-327.
- VandeBerg, J. L., and Johnston, P. G. (1977). A simple technique for long-term storage of erythrocytes for enzyme electrophoresis. *Biochem. Genet.* **15**:213.
- Walter, C. W. (1984). Invention and development of the blood bag. *Vox Sang.* **47**:318.

Biochemical Genetics

A journal for the reporting of original research in biochemical genetics of any organism, from virus to man. Papers will deal with the molecular aspects of genetic variation and evolution, mutation, gene action and regulation, immunogenetics, somatic cell genetics, and nucleic acid function in heredity and development and with the biochemical aspects of genetic defects (provided these are not purely clinical).

Papers on new methods that apply to the above topics will be included.

Review articles will also be published occasionally.

EDITOR

Hugh S. Forrest, Department of Zoology, University of Texas at Austin, Austin, Texas

EUROPEAN EDITOR

D. A. Hopkinson, The Galton Laboratory, London, England

ASSOCIATE EDITORS

Charles R. Shaw, University of Texas M. D. Anderson Hospital, Houston, Texas

Richard E. Tashian, University of Michigan, Ann Arbor, Michigan

Allan C. Wilson, University of California, Berkeley, California

ADVISORY BOARD

- C. Baglioni, State University of New York, Albany, New York
- L. Beckman, University of Umeå, Umeå, Sweden
- G. J. Brewer, University of Michigan, Ann Arbor, Michigan
- D. J. Cove, University of Leeds, Leeds, England
- C. H. Doy, Australian National University, Canberra, Australia
- W. F. Feenstra, University of Groningen, Groningen, The Netherlands
- J. R. S. Fincham, University of Edinburgh, Edinburgh, Scotland
- W. M. Fitch, University of Southern California, Los Angeles, California
- H. H. Fudenberg, University of South Carolina, Charleston, South Carolina
- J. B. Harborne, University of Reading, Reading, England
- L. Hood, California Institute of Technology, Pasadena, California
- J. J. Hutton, Children's Hospital Research Foundation, Cincinnati, Ohio
- G. Johnson, Washington University, St. Louis, Missouri
- T. H. Jukes, University of California, Berkeley, California
- H. N. Kirkman, University of North Carolina, Chapel Hill, North Carolina
- L. I. Korochkin, Institute of Developmental Biology, Moscow, USSR
- R. Lewontin, Harvard University, Cambridge, Massachusetts
- E. Margoliash, Northwestern University, Evanston, Illinois
- K. Paigen, University of California, Berkeley, California
- F. H. Ruddle, Yale University, New Haven, Connecticut
- J. G. Scandalios, North Carolina State University, Raleigh, North Carolina
- D. C. Shreffler, Washington University, St. Louis, Missouri
- O. Smithies, University of Wisconsin, Madison, Wisconsin
- Z. Šormová, Institute of Organic Chemistry and Biochemistry, Prague, Czechoslovakia
- H. E. Sutton, University of Texas at Austin, Austin, Texas
- E. Vesell, Pennsylvania State University, Hershey, Pennsylvania
- R. P. Wagner, Santa Fe, New Mexico
- G. S. Whitt, University of Illinois, Urbana, Illinois
- James E. Womack, Texas A&M, College Station, Texas
- W. J. Young, University of Vermont, Burlington, Vermont

Biochemical Genetics is published bimonthly by Plenum Publishing Corporation, 233 Spring Street, New York, N.Y. 10013. Subscription orders should be addressed to the publisher. *Biochemical Genetics* is abstracted or indexed in Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Awareness in Biological Sciences, Current Contents, Excerpta Medica, Index Medicus, Mental Health Abstracts, Referativnyi Zhurnal, and Science Citation Index. © 1987 Plenum Publishing Corporation. *Biochemical Genetics* participates in the Copyright Clearance Center (CCC) Transactional Reporting Service. The appearance of a code line at the bottom of the first page of an article in this journal indicates the copyright owner's consent that copies of the article may be made for personal or internal use. However, this consent is given on the condition that the copier pay the flat fee of \$5.00 per copy per article (no additional per-page fees) directly to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, Massachusetts 01970, for all copying not explicitly permitted by Sections 107 or 108 of the U.S. Copyright Law. The CCC is a nonprofit clearinghouse for the payment of photocopying fees by libraries and other users registered with the CCC. Therefore, this consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale, nor to the reprinting of figures, tables, and text excerpts. 0006-2928/87 \$5.00

Subscription rates:

Volume 25, 1987 (12 issues) \$395.00 (outside the U.S., \$441.00).

Second-class postage paid at New York, N.Y., and at additional mailing offices. Postmaster: Send address changes to *Biochemical Genetics*, Plenum Publishing Corporation, 233 Spring Street, New York, N.Y. 10013.

Printed in U.S.A.