

# Status of SPECIES off the U.S. West Coast in 2025

Michael Kinneen<sup>1</sup>

1. NOAA Fisheries Northwest Fisheries Science Center, 2725 Montlake Boulevard East



U.S. Department of Commerce  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Northwest Fisheries Science Center

## Table of contents

<b>Disclaimer</b>	<b>i</b>
<b>Executive Summary</b>	<b>ii</b>
Stock . . . . .	ii
Catches . . . . .	ii
Data and Assessment . . . . .	ii
Stock biomass and dynamics . . . . .	ii
Recruitment . . . . .	ii
Exploitation status . . . . .	ii
Ecosystem considerations . . . . .	ii
Reference points . . . . .	ii
Management performance . . . . .	ii
Harvest projections . . . . .	ii
Decision table . . . . .	ii
Scientific uncertainty . . . . .	ii
Research and data needs . . . . .	ii
Rebuilding projections . . . . .	ii
<b>1 Introduction</b>	<b>1</b>
1.1 Distribution and Stock Structure . . . . .	1
1.2 Life History and Ecosystem Interactions . . . . .	1
1.3 Fishery description . . . . .	2
1.4 Management History and Performance . . . . .	3
1.5 Fisheries off Canada and Alaska . . . . .	4
<b>2 Data</b>	<b>5</b>
2.1 Fishery-independent data . . . . .	5
2.1.1 SWFSC and NWFSC/PWCC Midwater Trawl Survey . . . . .	5
2.1.2 Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey . . . . .	6
2.1.3 Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey . . . . .	6
2.2 Fishery-dependent data . . . . .	8
2.2.1 Landings . . . . .	8
2.2.2 Fishery length and age data . . . . .	9
2.2.3 Discards . . . . .	10
2.2.4 Biological data . . . . .	10
2.2.5 Abundance indices . . . . .	14
2.3 Environmental and ecosystem data . . . . .	14
2.4 History of modeling approaches . . . . .	15
2.5 Responses to SSC Groundfish Subcommittee requests . . . . .	15
2.6 Model Structure and Assumptions . . . . .	15
2.6.1 Model Changes from the Last Assessment . . . . .	15
2.6.2 Modeling Platform and Structure . . . . .	15

2.6.3	Model Parameters . . . . .	15
2.6.4	Key Assumptions and Structural Choices . . . . .	15
2.7	Base Model Results . . . . .	16
2.7.1	Parameter Estimates . . . . .	16
2.7.2	Fits to the Data . . . . .	17
2.7.3	Population Trajectory . . . . .	17
2.8	Model Diagnostics . . . . .	17
2.8.1	Convergence . . . . .	18
2.8.2	Parameter Uncertainty . . . . .	18
2.8.3	Sensitivity Analyses . . . . .	18
2.8.4	Retrospective Analysis . . . . .	20
2.8.5	Likelihood Profiles and key parameters . . . . .	20
2.9	Unresolved Problems and Major Uncertainties . . . . .	20
<b>3</b>	<b>Management</b>	<b>21</b>
3.1	Reference Points . . . . .	21
3.2	Unresolved problems and major uncertainties . . . . .	21
3.3	Harvest Projections and Decision Tables . . . . .	21
3.4	Evaluation of Scientific Uncertainty . . . . .	22
3.5	Regional management considerations . . . . .	22
3.6	Research and Data Needs . . . . .	22
3.7	Acknowledgements . . . . .	23
3.8	References . . . . .	24
<b>4</b>	<b>Tables</b>	<b>25</b>
4.1	Data . . . . .	25
4.1.1	Fishery-dependent data . . . . .	25
4.2	Figures . . . . .	35

Please cite this publication as:

Michael Kinneen, Maurice Goodman, . (2025) Status of Widow Rockfish off the U.S. West Coast in 2025. Pacific Fishery Management Council. [XX] p.

**Disclaimer**

These materials do not constitute a formal publication and are for information only. They are in a pre-review, pre-decisional state and should not be formally cited or reproduced. They are to be considered provisional and do not represent any determination or policy of NOAA or the Department of Commerce.

**Executive Summary**

Checking to see if this works Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey (Triennial Survey)

**Stock**

**Catches**

**Data and Assessment**

**Stock biomass and dynamics**

**Recruitment**

**Exploitation status**

**Ecosystem considerations**

**Reference points**

**Management performance**

**Harvest projections**

**Decision table**

**Scientific uncertainty**

**Research and data needs**

**Rebuilding projections**

## 1 Introduction

*Sebastes entomelas* (Widow Rockfish) is named after its black-lined gut cavity (*ento* meaning within and *melas* meaning black). It has been referred to as buda, beccafico (Italian bird), and viuva (widow) prior to the 1930s. More recently, the Widow Rockfish is also called brownie, belinda bass, brown bomber, and soft brown.

This is an assessment of Widow Rockfish that inhabit the waters off California, Oregon, and Washington from the U.S.-Canadian border in the north to the U.S.-Mexico border in the south, and does not include Puget Sound waters (Figure 1). This assessment represents a thorough reconsideration of the data, data preparation, and model structure for assessing Widow Rockfish, including reinvestigations of recent and historical catches (including discards), length and age data, and fleet structure.

### 1.1 Distribution and Stock Structure

Widow Rockfish inhabit water depths of 25–370 m from northern Baja California, Mexico to Southeastern Alaska, and are most abundant from British Columbia to Northern California. Although catches north of the U.S.-Canada border or south of the U.S.-Mexico border were not included in this assessment, it is possible that these populations contribute to the biomass of Widow Rockfish off of the U.S. West Coast through adult migration and/or larval dispersion.

There is little evidence of genetically separate stocks along the U.S. coast and past assessments have used a single area, coastwide model with multiple fisheries (He et al. 2011). In 2011, a two-area assessment model was brought forward for review, and was found to be similar to a coastwide model (He et al. 2011). There is some evidence of biological differences between areas. For example, Widow Rockfish collected off California tend to mature at a smaller length than Widow Rockfish collected off of Oregon (**barss\_maturity\_1987?**). This may be due to environmental or anthropogenic effects rather than genetic differences. The connectivity of Widow Rockfish populations throughout its range is unknown and it was decided to continue with a single area model for this assessment instead of potentially lose prediction power by splitting the data into two separate areas.

### 1.2 Life History and Ecosystem Interactions

Widow Rockfish are atypical for West Coast rockfish species because they form dense midwater aggregations at night, which were largely undetected until the late 1970s. They

are typically found over high relief strata and near cobblestone. The diet of Widow Rockfish is dominated by species that comprise the deep scattering layers, including salps, myctophids, *Sergestes similis* (a caridean shrimp), and euphausiids (Adams 1987).

Widow Rockfish are ovoviviparous with gestation lasting from 1 to 3 months. Parturition occurs earlier in southern latitudes (December-March off California) than in northern latitudes (April in British Columbia) and occur once a year (**barss\_maturity\_1987?**). Estimates of fecundity of Widow Rockfish range from 95,375 oocytes at 33 cm to 1,113,000 oocytes at 52 cm (**Boehlert\_fecundity\_1982?**).

There is little information regarding the movement of Widow Rockfish. Past assessments have assumed a two-area model because of differences in growth and maturity (see (He et al. 2011)). However, using recent observations from the NWFSC shelf/slope survey to follow two separate cohorts through time and space suggests that Widow Rockfish may recruit in the south and disperse northward as they age (Figure 2). Spatial recruitment and movement patterns of Widow Rockfish are uncertain and much more investigation and sampling is needed to fully understand them.

### 1.3 Fishery description

Widow Rockfish were lightly exploited by bottom trawl and hook-and-line gears prior to the 1980s. After many attempts to start trawl fisheries off the west coast of the United States in the late 1800s, the availability of otter trawl nets and the diesel engine in the mid-1920s helped trawl fisheries expand (Douglas 1998). The trawl fisheries really became established during World War II when demand increased for shark livers and bottomfish. A mink food fishery also developed during World War II (Jones and Harry Jr 1960). Foreign fleets began fishing for rockfish in the mid-1960s until the EEZ was implemented in 1977 (Rogers 2003). Longline catches of Widow Rockfish are present from the turn of the century and continue in recent years, mainly from fisheries targeting sablefish and halibut.

In the late 1960s and early 1970s, it is reported that foreign fishing vessels caught large numbers of Widow Rockfish (Rogers 2003). In the late 1970s a domestic midwater trawl fishery began developing off of Oregon when it was realized that Widow Rockfish form dense aggregations at night (**Gunderson\_great\_1984?**). The fishery expanded very quickly, with landings from trawl, net, and hook-and-line gears increasing more than 20 times by the early 1980s (Table 1). As early as 1982, trip limits were imposed to keep catches below recommended annual levels (Table 3). Trip limits became more restrictive over the years until Widow Rockfish was declared overfished in 2001. In 2002, harvest guidelines were greatly reduced and over the last decade have been small, although increasing since 2004 (Table 4).



Historical discarding practices are not well known, but it is believed that little discarding occurred prior to management restrictions. With the introduction of trip limits, limited data from the mid-1980s show occasional very high discard rates of Widow Rockfish from tows that occurred near the end of a trip.

More detailed information of the fisheries in each state is given in Section 2.2.1 where the reconstructed landings are discussed.

#### 1.4 Management History and Performance

Widow Rockfish has been a small large component of groundfish fisheries since the late 1970s. The landings of Widow Rockfish have been historically governed by harvest guidelines and trip limits, while recently management is imposed with total catch harvest limits in the form of overfishing limits (OFLs), acceptable biological catches (ABCs), and annual catch limits (ACLs). A trawl rationalization program, consisting of an individual fishing quota (IFQ) or catch shares system was implemented in 2011 for the limited entry trawl fleet targeting non-whiting groundfish, including Widow Rockfish, and the trawl fleet targeting and delivering whiting to shore-based processors. The limited entry at-sea trawl sectors (motherships and catch-processors) that target whiting and process at sea are managed in a system of harvest cooperatives.

Limits on Widow Rockfish were first established in 1982 (Table 3). These were implemented as trip limits and cumulative landing limits that were first imposed by trip, then week, then every 2 weeks, month, 2 months, and eventually into periods. In many years, the trip limits on Widow Rockfish were significantly reduced at the end of the year to avoid exceeding the harvest recommendations. Some important years were 1985 when trip limits were reduced to 30,000 pounds once per week or 60,000 pounds once every 2 weeks, 1990 when trip limits were reduced to 15,000 or 25,000 pounds every one or two weeks, respectively, 1998 when a 25,000 pound cumulative limit per two-month period was implemented, and 2011 when catch shares was implemented.

A sorting requirement was implemented for Widow Rockfish in the early 1980s with California beginning in 1982, Oregon in 1984, and Washington in 1988. Some important events that could affect fishery selectivity are the gear restrictions implemented in 2000, implementation of Rockfish Conservation Areas (RCA's) in 2002, seasonal changes to the RCA's in 2007, and the beginning of catch shares in 2011.

Table 4 shows that recent landings have been below recommended catch levels. Landings are a considerable amount below the ACL, and it is unlikely that total mortality has exceeded the ACL in the last 10 years.

### 1.5 Fisheries off Canada and Alaska

Widow Rockfish are distributed throughout Canada and Southeast Alaska and are commonly caught in trawl and hook-and-line fisheries. However, the landings from the fisheries in these areas are estimated to harvest Widow Rockfish at much smaller rate than has been observed off California, Oregon, and Washington mostly due to lower abundance of Widow Rockfish, but also partly due to precautionary behavior of Canadian managers after the large catches followed by management restrictions and concerns of the U.S. fishery in the early 1980s.

Alaska formed the “Other Rockfish” complex in 2012 from the combination of Other Slope Rockfish and the Widow and Yellowtail Rockfishes from the Pelagic Shelf Rockfish category. This new complex includes 18 species and Widow Rockfish are a small proportion of the catch (less than 5%). Total biomass estimates are provided by the Gulf of Alaska (GOA) triennial/biennial trawl survey. ABC’s and OFL’s were set for the Other Rockfish Complex and component species in 2013 with a recommended OFL in 2014 of 5,347 mt for the complex. Widow Rockfish comprise a small part of this complex in Alaska.

The fishery for Widow Rockfish in British Columbia, Canada started in 1986 although some very small landings occurred in the mid-1970s. Landings peaked at about 4,500 mt in 1990 and were around 2,000 mt throughout the 1990s [dfo\_widow\_1999]. Most landings occurred in a midwater trawl fishery, but there have also been reports of “nuisance catches in the salmon troll fishery”. An assessment of Widow Rockfish in Canada was completed in 1998 (**stanley\_shelf\_1999?**) as part of a shelf rockfish complex. Additional research has since been done on the estimation of biomass of particular aggregations of Widow Rockfish (Stanley et al. 2000), but no formal assessment has been done since.

## 2 Data

Many sources of data were available for this assessment, including indices of abundance (Table 5), length observations, and age observations from fishery-dependent and fishery-independent sources.

### 2.1 Fishery-independent data

Data from three fishery-independent surveys were used in this assessment: 1) the SWFSC and NWFSC/PWCC Midwater Trawl Survey (hereafter, “juvenile survey”); 2) the Alaska Fisheries Science Center (AFSC)/NWFSC Triennial Shelf Trawl Survey (hereafter, “triennial survey”); and 3) the NWFSC West Coast Groundfish Bottom Trawl Survey (hereafter, “WCGBTS”). These surveys employed different designs and sampling methodologies, were conducted during different years and time periods within years, and included coverage over different areas of the coast. In some instances, the survey frequency, depths, and geographic areas covered were not internally consistent within surveys. A brief description of each survey is provided below.

Strata were defined by latitude and depth to analyze the catch-rates, length compositions, and age compositions using stratified random sampling theory (Table 6 & Table 7). The latitude and depth breaks were chosen based on the design of the survey as well as by looking at biological patterns in relation to latitude and depth. Indices of abundance for all of the surveys were derived using model based approaches described below.

#### 2.1.1 SWFSC and NWFSC/PWCC Midwater Trawl Survey

We updated the coastwide pre-recruit index of abundance for widow rockfish using data from three midwater trawl surveys targeting young-of-the-year (YOY) rockfish (SWFSC and NWFSC/PWCC Midwater Trawl Survey (Juvenile Survey)), provided by Tanya Rogers (SWFSC, pers. comm.). All surveys used identical gear, enabling the construction of a consistent coastwide index spanning from 36°N to the U.S./Canada border since 2004. For building the widow rockfish pre-recruit index, we used data from 2001 to 2024 without spatial subsetting (including CA, OR, and WA). Sampling in 2020 was limited due to the COVID-19 pandemic and excluded from all models. In 2010 and 2012, coverage was incomplete, so these years were used to construct the index but excluded from the final model to align with the 2019 assessment. Data from 2001–2003 were also excluded following the 2015 assessment due to limited spatial coverage (36°30' to 38°20' N latitude). However, a sensitivity analysis was conducted to examine the impact of including those early years (see Figure: Sensitivity Analysis).

The index was built using a spatial GLM with the sdmTMB package [anderson\_sdmTMB\_2022], modeling 100-day standardized catch-per-tow as a function of year (fixed effect), Julian date (GAM smoother,  $k = 4$ ), spatial random field, and spatiotemporal random effects. Models with Tweedie, delta-lognormal, and delta-gamma error structures were compared; DHARMA residuals and simulation-based diagnostics indicated the Tweedie model performed best. The index shows a strong increasing trend in juvenile abundance from 2017 to 2023, with a slight decline in 2024. Despite the dip, recent values remain high relative to the previous decade, and uncertainty estimates support the robustness of this trend.

### **2.1.2 Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey**

The Triennial Survey was first conducted by the Alaska Fisheries Science Center (AFSC) in 1977, and the survey continued until 2004 (weinberg\_2001\_2002?). Its basic design was a series of equally-spaced east-to-west transects across the continental shelf from which searches for tows in a specific depth range were initiated. The survey design changed slightly over time. In general, all of the surveys were conducted in the mid summer through early fall. The 1977 survey was conducted from early July through late September. The surveys from 1980 through 1989 were conducted from the middle of July to late September. The 1992 survey was conducted from the middle of July through early October. The 1995 survey was conducted from early June through late August. The 1998 survey was conducted from early June through early August. Finally, the 2001 and 2004 surveys were conducted from May to July.

Haul depths ranged from 91–457 m during the 1977 survey. Due to haul performance issues and truncated sampling with respect to depth, the data from 1977 were omitted from this analysis. The surveys in 1980, 1983, and 1986 covered the U.S. West Coast south to 36.8°N latitude and a depth range of 55–366 m. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5°N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55–500 m and surveyed south to 34.5°N. In 2004, the final year of the Triennial Survey series, Northwest Fisheries Science Center (NWFSC) Fishery Resource and Monitoring Division (FRAM) conducted the survey following similar protocols to earlier years.

### **2.1.3 Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey**

The Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (WCGBTS) is based on a random-grid design; covering the coastal waters from a depth of 55–1,280 m (Bradburn, Keller, and Horness 2011). This design generally uses four industry-chartered vessels per year assigned to a roughly equal number of randomly

selected grid cells and divided into two ‘passes’ of the coast. Two vessels fish from north to south during each pass between late May to early October. There were only two vessels used in 2019 and three in 2013, with one of the three that year unable to complete its survey pass due to a government shutdown. No survey occurred in 2020 due to Coronavirus disease (COVID-19). This design therefore incorporates both vessel-to-vessel differences in catchability, as well as variance associated with selecting a relatively small number (approximately 700) of possible cells from a very large set of possible cells spread from the Mexican to the Canadian borders. Note that the Survey is not permitted to access the Cowcod Conservation Areas (CCAs) in Southern California.

Widow rockfish are not commonly caught in the WCGBTS. Higher catch rates occur north of 40° N latitude and catches are rare south of 36° N latitude (Figure 11). Few large fish are found shallower than 100 m and few small fish are found in the deeper water of the slope. There is no clear trend in length with latitude other than smaller fish tend to occur south of approximately 36° N latitude, and there appears to be some very small fish found near 39° N latitude.

Geostatistical models of biomass density were fit to survey data using the R package [Species Distribution Models with Template Model Builder](#) ([sdmTMB](#)) ([Anderson:2022:SRP?](#)). This approach reflects an updated approach compared to the 2015 assessment (non-spatial delta-GLMM) and the 2019 update assessment (VAST delta-lognormal model). These models can account for latent spatial factors with a constant spatial Gaussian random field and spatiotemporal deviations to evolve as a random walk Gaussian random field using a 200 knot grid of the survey area ([thorson\\_geostatistical\\_2015?](#)). The prediction grid was also truncated to only include available survey locations in depths between 55–500 m to limit extrapolating beyond the data and edge effects. Tweedie, delta-binomial, delta-gamma, and mixture distributions, which allow for extreme catch events, were investigated. The positive catch weight model includes survey pass (‘first’ or ‘second’) to account for missing data, i.e., the incomplete second pass of the 2013 survey. Vessel-year effects, which have traditionally been included in index standardization for this survey, were not included as the estimated variance for the random effect was close to zero. Vessel-year effects were more prominent when models did not include spatial effects and were included for each unique combination of vessel and year in the data to account for the random selection of commercial vessels used during sampling ([helsler\\_generalized\\_2004?](#); [thorson\\_accounting\\_2014?](#)).

Results are only shown for both the delta-gamma and delta-lognormal distributions. Both models converged (positive, definite Hessian matrix) but predicted data from both models showed slightly right-heavy tails compared to null expectations, with the gamma model having stronger divergence. The delta-lognormal distribution ultimately had to the best model diagnostics, e.g., similar distributions of theoretical normal quantiles and model quantiles, high precision, lack of extreme predictions that are incompatible with the life history, and low Akaike information criterion (AIC). Spatiotemporal estimates

of positive catch for the delta-lognormal distribution were then converted into annual indices using `sdmtmb::get_index()` function. (**Anderson:2022:SRP?**)

The index estimate is relatively stable, with a slightly increasing trend in recent years and a moderate peak in 2016. Overall, the lognormal index estimates were more comparable to the 2019 spatiotemporal VAST-based index and seemed less influenced by potential extreme catch events, particularly in 2013 and 2016; for these reasons, in addition to better model performance described above, the delta-lognormal sdmTMB-based index was used for the base model in this assessment. The delta-lognormal mean value (2262.824) was slightly lower than the means of the index values used in the 2015 assessment (2701.12) and the 2019 update assessment (3301.765). Comparisons of the different error structures, design-based estimate and the VAST index used in 2019 are in Figure XX.

## 2.2 Fishery-dependent data

### 2.2.1 Landings

Widow rockfish have been caught in trawl and hook-and-line fisheries since the early part of the 20th century. Widow rockfish are a desirable rockfish and are not likely to be discarded for market reasons. However, smaller widow rockfish are found at shallower depths and discarding practices in the early 1900s are uncertain. In data from the early 1980s, widow rockfish have had their own landing category, beginning in California in 1982, Oregon in 1984, and Washington in 1988. Estimates of historical landings of widow rockfish rely upon species-composition sampling data from each period. The uncertainty in species composition is greater in past years, with less systematic and extensive sampling occurring prior to 1980. Consequently, the precision with which landings of widow rockfish can be estimated likely decreases for earlier years.

The definitions of fishing fleets have not been changed from those in the 2015 and 2019 assessments. Five fishing fleets were specified within the model: 1) a shorebased bottom trawl fleet with coastwide catches from 1916–2024, 2) a shorebased midwater trawl fleet with coastwide catches from 1979–2024, 3) a mostly midwater trawl fleet that targets Pacific Hake/Whiting (*Merluccius productus*) and includes a foreign and at-sea fleet with catches from 1975–2024, a domestic shorebased fleet that targeted Pacific Hake with catches from 1991–2024, and foreign vessels that targeted Pacific Hake and rockfish between 1966–1976, 4) a net fishery consisting of catches mostly from California from 1981–2024, and 5) a hook-and-line fishery (predominantly longline) with coastwide catches from 1916–2024. As in previous assessments, catches from Puget Sound and those from commercial shrimp trawls, commercial pots, and recreational fisheries were excluded (as these are generally minimal).

Catches from all years (1916-2018) were carried forward into this assessment, with two exceptions. First, discards from the hook-and-line fleet were added to the removals for this fleet. The hook-and-line removals of widow rockfish are extremely minimal (Figure 1) and comprised only approximately 0.2% of the total removals over the last twenty years, with discard being a small fraction of that. The biological samples of the discard amount are also scarce, with input sample sizes not exceeding 6 and averaging around 3 per year (Table X). With this limited data, the model was unable to reliably estimate retention parameters and exhibited substantial sensitivity to even slight changes in discard amounts within the hook-and-line fleet. Therefore, in this assessment, we added hook-and-line discards to hook-and-line landings. Second, because PacFIN appear to underestimate midwater trawl catches in California in 1979-1980 (Edward Dick, Pers. Comms.) we adjusted midwater and bottom trawl catches from California in these years to reflect the ratio of California midwater to bottom trawl catches in 1981-1982. New catches (2019-onward) from PacFIN and ASHOP were otherwise appended onto the 1916-2018 landings and apportioned among fleets using the same criteria as those documented in the 2015 assessment.

### 2.2.2 Fishery length and age data

Biological data from commercial fisheries that caught widow rockfish were extracted from PacFIN (PSMFC) on July 3, 2019, from CALCOM on July 3, 2019 and from the NORPAC database on July 3, 2019. Lengths taken during port sampling in California, Oregon, and Washington were used to calculate length and age compositions. The data were classified into bottom trawl, midwater trawl, hake trawl, net, and hook-and-line fleets

Table 10 shows the number of landings sampled and Table 11 shows the number of lengths taken for each year, gear, and fleet from the three states. Table 12 shows these numbers for the at-sea fleet.

Consistent with the 2015 assessment, length and age samples from PacFIN and CALCOM were expanded up to the total landing then combined into state-specific frequencies (Table 13). Expansion factors were calculated in a way such that large expansions would not occur and based on ideas first presented by Owen Hamel (pers. comm., NWFSC). First the expansion factor ( $E_k$ ) was the total catch weight ( $W_k$ ) divided by the sample weight ( $w_k$ ), and raised to 0.9 to account for non-homogeneity within a trip. Then, expansion factors greater than 300 were capped (100 for net fisheries) to reduce the influence of small samples (i.e., a few fish representing a large catch). The predicted total numbers at length or age weighted by landings for each state were added to create a coast-wide length frequency. The effective sample sizes of the state combined length frequencies were determined from the following formula, which has been used in previous widow rockfish assessments as well as other west coast groundfish assessments.

Fishery Samples	Survey Samples
$N_{eff} = N_{sample} + 0.138N_{fish}, \frac{N_{fish}}{N_{sample}} < 44$	$N_{eff} = N_{sample} + 0.0707N_{fish}, \frac{N_{fish}}{N_{sample}} < 55$
$N_{eff} = 7.06N_{sample}, \frac{N_{fish}}{N_{sample}} \geq 44$	$N_{eff} = 4.89N_{sample}, \frac{N_{fish}}{N_{sample}} \geq 55$

This is slightly different than the sample size of 2.43 per haul for rockfish that (**stewart\_\_bootstrapping\_2014?**) report. Observed lengths were expanded to the tow from At-Sea Hake Observer Program samples (NORPAC). Tows are typically well sampled, thus expansion factors were not modified from what was calculated. Hake fishery length compositions were created by combining shoreside and at-sea length compositions, weighting by the catch from each sector. The effective sample sizes for hake fishery length and age comps were calculated using the above equations for the shoreside fleet and added to the number of tows sampled from the at-sea fleet.

Expanded length compositions for bottom trawl, midwater trawl, hake fisheries, net, and hook-and-line are shown in Figure 17 to Figure 21. It is quickly apparent that all of these fisheries rarely land fish less than 26 cm. All of the non-hake fleets show a strong cohort coming through in the late 1970s and early 1980s, and then another cohort coming through in the late 1980s. Sample sizes typically dropped off after 2000, except in the hake fishery where nearly every tow is sampled. Age compositions for the five fleets are shown in Figure 22 and Figure 26. Occasional cohorts appear to move through the population, indicating that widow rockfish population dynamics may be characterized by episodic recruitment events.

### 2.2.3 Discards

– TO DO!!! –

$$D_{y,f} = \frac{d_{y,f}}{r_{y,f}} R_{y,f}$$

### 2.2.4 Biological data

#### 2.2.4.1 Weight-length relationship

Weight-at-length data, which are the same used in the 2015 assessment, were collected from fisheries sampling and by the Triennial and NWFSC WCGBT Surveys, and were



used to estimate a weight-length relationship for widow rockfish (Figure 30). Weight-at-length was similar between sources with the fishery samples showing a slightly smaller weight at large sizes when compared to the survey data (Figure 31). WCGOP data were not used because only small fish were sampled, the weight of these small fish were typically less than from other sources (Figure 30), and the curves fitted to only WCGOP data were unable to estimate the slope. There were only 81 observations from the WCGOP data, which is a small amount of data compared to everything available. However, these observations may be useful to understand discards.

The weight-length relationship used in the 2011 assessment was similar for males but predicted slightly heavier females at larger sizes than the 2015 assessment (Figure 31). The following relationships between weight and length for females and males were estimated for the 2015 assessment from all of the data combined and were used in the current assessment:

$$\text{Females:} \quad \text{weight} = 1.7355 \times 10^{-5} \cdot \text{Length}^{2.9617}$$

$$\text{Males:} \quad \text{weight} = 1.4824 \times 10^{-5} \cdot \text{Length}^{3.0047}$$

where weight is measured in kilograms and length in cm. These relationships were used in the assessment as fixed relationships.

#### 2.2.4.2 Maturity schedule

Estimates of maturity used in this update were the same as the 2015 assessment. Estimates of maturity at length have been presented by (**barss\_maturity\_1987?**), (**echeverria\_thirty-four\_1987?**), and (**love\_life\_1990?**). (**barss\_maturity\_1987?**) supplied data collected from Oregon and California commercial and recreational samples, which allowed us to estimate the proportion mature-at-length and proportion mature-at-age for samples from each state (Figure 32). As noted by (**barss\_maturity\_1987?**), the samples from Oregon matured at older age and larger length. Estimates of maturity-at-length from California reported by (**barss\_maturity\_1987?**) are similar to estimates of length-at-50%-mature from samples collected in California reported by (**echeverria\_thirty-four\_1987?**) and (**love\_life\_1990?**), although (**barss\_maturity\_1987?**) show the smallest length-at-50%-mature. To maintain some consistency with the 2011 assessment and to avoid any potential growth issues by area, the 2015 assessment used maturity-at-age data from the 2011 assessment, but used the data provided by (**barss\_maturity\_1987?**) to estimate a new maturity curve following a logistic function with the data from California and Oregon equally weighted to avoid California dominating

the estimated relationship. This maturity-at-age curve falls between the estimated California and Oregon maturity-at-age curves (Figure 32, right), with the age-at-50%-mature estimated at 5.47 and with a slope of -0.7747 (as specified in SS). This logistic maturity-at-age curve was used in the 2015 and 2019 update assessment except that maturity-at-age for ages 2 and lower were set equal to zero (Table 19).

#### 2.2.4.3 Fecundity

Fecundity in rockfish is often not a linear function of weight, but increases faster at larger weights (Dick 2009). Therefore, this relationship is often accounted for in rockfish assessments by using spawning output (numbers of eggs) to determine current status. (Dick 2009) did not find a significant relationship between the number of eggs per gram of body weight and body weight for widow rockfish. Therefore, spawning output was assumed to be proportional to weight, which is the same as spawning biomass, and is reported here.

#### 2.2.4.4 Natural Mortality

Natural mortality used in this update differed from the 2015 assessment. Natural mortality ( $M$ ) is a parameter that is often highly uncertain in fish stocks. Past assessments of widow rockfish assumed constant natural mortality of 0.125 yr<sup>-1</sup> or 0.15 yr<sup>-1</sup>. The 2011 assessment estimated  $M$  with a prior developed by Owen Hamel (NWFSC, pers. comm.) using methods described in (Hamel 2014). This prior was based on a maximum age of 44 and 40 for females and males, respectively, a mean temperature of 8 degrees Celsius (about 150m deep off of Oregon), and a gonadosomatic index of 9.99% and 1.86% for females and males, respectively (**love\_life\_1990?**). The sex-specific lognormal priors for  $M$  have medians of 0.124 yr<sup>-1</sup> and 0.129 yr<sup>-1</sup> for females and males, respectively, and a coefficient of variation (CV) of 30.7% for each sex. In 2015, discussions with Owen Hamel (NWFSC) led to the development of a new prior based solely on maximum age to use when estimating  $M$ . Using all of the available age data, a maximum age of 54 was determined for both females and males, although it has been rare to observe widow rockfish older than about 45 years old (Figure 33). This resulted in a prior with a much smaller median (0.0810 or -2.513284 in log space) and a larger standard deviation in log space (0.523694). For the update assessment, an updated meta-analysis resulted in a prior with a slightly smaller median than the 2015 assessment (0.10 or -2.30 in log space) and a smaller standard deviation in log space (0.438). Figure 34 shows that these prior distributions are wide and not highly informative.

#### 2.2.4.5 Length-at-age

Estimates of length-at-age used in this update were the same as the 2015 assessment. Two different labs have aged the majority of processed otoliths for widow rockfish. The SWFSC has been aging widow rockfish otoliths for many years, including all of the fishery data prior to 2011 and otoliths collected from the NWFSC WCGBT survey in 2009 and 2010. The Cooperative Ageing Project (CAP) in Newport, Oregon aged 1,100 otoliths from the NWFSC WCGBT survey, 2,026 otoliths provided by ASHOP, and 3,467 otoliths collected by port samplers. All of the commercial fishery samples were collected in the years 2011–2014. In total, there are 105,814 paired age and length observations ranging from 1978 to 2014. Figure 35 shows the lengths and ages for all years and all data as well as predicted von Bertalanffy fits to the data. Females grow larger than males and sex specific growth parameters were estimated at the following values:

$$\text{Females:} \quad L_{\infty} = 50.34, \quad k = 0.15, \quad t_0 = -2.22$$

$$\text{Males:} \quad L_{\infty} = 44.19, \quad k = 0.21, \quad t_0 = -1.78$$

The data from each source (ASHOP, port sampling/BDS, Triennial survey, and NWFSC survey) are shown in Figure 36 with fitted von Bertalanffy lines. All of these sources are quite similar, especially observations from ASHOP and the NWFSC survey. The standard deviation (SD) and coefficient of variation (CV) of length-at-age are shown in Figure 37. Modelling the CV as a function of predicted length-at-age appears to be somewhat linear from a value just over 0.1 at small lengths and slightly less than 0.045 at larger lengths. However, variance in length-at-age was estimated separately in stock-synthesis.

#### 2.2.4.6 Sex ratios

Females tend to grow larger than males and it is expected that the proportion of females approaches one at large lengths and is less than 0.5 at intermediate lengths. Figure 38 shows that the proportion of females at length from survey data is approximately 50% until approximately 34 cm, when the proportion of females drops below 50%. At lengths larger than 46 cm, the proportion of females increases rapidly to one, suggesting that few males grow larger than 50 cm.

#### 2.2.4.7 Ageing bias and imprecision

Uncertainty surrounding the ageing-error process for widow rockfish used in the 2015 assessment was incorporated by estimating ageing error by age. No changes were made

from the 2015 assessment for the update. Age-composition data used in the model were from break-and-burn and surface reads and were aged by the Cooperative Ageing Project (CAP) in Newport, Oregon and the SWFSC in Santa Cruz, California. 12 Break-and-burn double reads of 1788 otoliths were performed by both the CAP and the SWFSC lab combined. Additionally, 100 otoliths were read both by surface and break-and-burn methods. An ageing error estimate was made based on these double reads using a computational tool specifically developed for estimating ageing error (Punt et al. 2008), and using release 1.0.0 of the R package `nwfscAgeingError` (**thorson\_nwfscageingerror\_2012?**) for input and output diagnostics, publicly available at: <https://github.com/nwfsc-assess/nwfscAgeingError>. The maximum aged fish read by the surface reading method was 10 years and the cross otolith reads between the surface and break-and-burn ageing methods showed limited variation. Therefore, a unique ageing error was not created for surface read otoliths. A non-linear standard error was estimated by age where there is more variability in the estimated age of older fish was estimated for each reading lab (Table 20 and Figure 39).

#### 2.2.5 Abundance indices

### 2.3 Environmental and ecosystem data

## Assessment model

An age-structured stock assessment model was used to predict the biomass trajectory of Widow Rockfish with an approach of balancing parsimony with complexity. This allowed for the determination of general trends in the biomass over time without introducing extraneous data partitions that explain little additional variation. The assessment followed the same model structure as the 2015 base assessment.

### **2.4 History of modeling approaches**

Refer to the most recent full assessment for additional information.

### **2.5 Responses to SSC Groundfish Subcommittee requests**

To be completed after review.

### **2.6 Model Structure and Assumptions**

For this update assessment, new versions of the previously used software were used. Stock Synthesis v3.30.13 was used to estimate the parameters in the 2019 model. R4SS, version 1.35.3, along with R version 3.5.3 were used to investigate and plot the 2019 model fits. For the update, Stock Synthesis v3.30.2 and R4SS, version 1.51.0, along with R version 4.4.2 were used. Bridging from Stock Synthesis v3.24U to v3.30.13 is illustrated in Figure 41. A summary of the data sources used in the model (details discussed above) is shown in Figure 40. Stock Synthesis has many options when setting up a model and the assessment model for Widow Rockfish was set up in the following manner.

#### **2.6.1 Model Changes from the Last Assessment**

#### **2.6.2 Modeling Platform and Structure**

#### **2.6.3 Model Parameters**

#### **2.6.4 Key Assumptions and Structural Choices**

Refer to the most recent full assessment for additional information.

## 2.7 Base Model Results

### 2.7.1 Parameter Estimates

#### 3.3.1 Parameter estimates

The estimates of natural mortality 0.124599 yr<sup>-1</sup> and NA yr<sup>-1</sup> for females and males, respectively) were higher than suggested by the medians of the prior distributions used in this assessment and the 2015 assessment. Fixing M at lower values than those estimates resulted in a pattern of reduced recruitment immediately before the fishery started. This suggests that the model is doing what it can to reduce the number of observations of older fish in the data. The estimates of M fall within the 95% confidence interval of the prior distribution (0.0425–0.237), and are shown in Figure 44.

Estimating M is difficult in stock assessments, and the estimated values may represent model misspecification instead of the actual life-history trait. However, in alternative models to the base case model, the estimates of M were rarely less than 0.14 yr<sup>-1</sup> (Table 29). Uncertainty in the estimated M was also much less than the range of the prior (Figure 44). The assumption that appeared to have the largest effect on M was introducing dome-shaped selectivity in the midwater trawl fleet, which made M smaller (Table 29).

Selectivity curves were estimated for commercial and survey fleets and parameter estimates are provided in Table 24. The final base model assumed asymptotic selectivity (double-normal selectivity curve ) for each fishery, except for the midwater trawl fishery. The NWFSC and Triennial surveys both used spline curves. All selectivity curves were length-based and are the same shape as in the 2015 benchmark. Time blocks were used for the bottom trawl, midwater trawl and hook-and-line fisheries as indicated in Table 21. The estimated selectivity, retention, and keep (the product of selectivity and retention) curves for the trawl and hook-and-line fleets are shown in Figure 45. The selectivity curves showed a shift to larger fish in 2002 for the bottom trawl fishery and a shift to smaller fish in 2003 for the hook-and-line fishery. The bottom trawl shift is consistent with the introduction of the RCA and gear restrictions (shoreward of the 18 RCA) that virtually eliminated fishing in shelf habitats where smaller Widow Rockfish would more likely be encountered. Around this same time, the fixed-gear RCA specifications began preventing fishing between 30 and 100 fm.

The retention curves showed a shift to retaining a lower percentage of fish since trip limits were introduced, but increases in recent years. The asymptote of the retention curve for the bottom trawl fishery sequentially decreased as more management restrictions were introduced to about 50% retention of larger fish in the 1998-2010 period.

Midwater trawl and hook-and-line fisheries estimated an asymptote to retention just above 80% for the period 1983-2010. Both the selectivity for the hake fleet and the selectivity

of the net fleet did not support dome-shaped selectivity (Figure 46). The estimated selectivity curves for the Triennial and NWFSC WCGBT surveys were similar to each other except that the triennial survey selected larger fish (Figure 46). The NWFSC WCGBT survey was no longer minimally dome-shaped as in the 2015 assessment.

In 2015, additional survey variability (process error added directly to each year's input variability) for the triennial and NWFSC WCGBT surveys was not estimated in the model because the estimate was zero. To avoid bound issues in estimation of the Hessian, the authors fixed these at zero because the model-based results provided reasonable estimates of variance. We retained the same modelling approach for the update assessment. The additional standard deviation added to the fishery-dependent indices was quite large, ranging from 0.16 for the bottom trawl index and 0.58 for the foreign at-sea hake fleet. The additional variability on the juvenile survey was the highest, at 0.83, giving the index very little weight in the model.

The estimates of maximum size for both females and males (Table 23) were not unexpected given the data in Figure 35. Estimates of  $k$  were slightly different in the model, but that is expected when accounting for selectivity. Estimated growth curves are shown in (Figure 47).

Estimates of recruitment suggest that the Widow Rockfish population is characterized by variable recruitment with occasional strong recruitments and periods of low recruitment (Figure 48). There is little information regarding recruitment prior to 1965, and the uncertainty in these estimates is expressed in the model. There are very large, but uncertain, estimates of recruitment in 2013, 1970, 2008, and 1971. Other large recruitment events (in descending order of magnitude) occurred in 1978, 2014, 1981, 2010, and 1991. The five lowest recruitments (in ascending order) occurred in 2012, 2011, 1976, 2007, and 1973. Estimates of recruitment appear to be episodic and characterized by periods of low recruitment. Two of the four largest estimated recruitments occurred in the last 11 years.

### **2.7.2 Fits to the Data**

### **2.7.3 Population Trajectory**

## **2.8 Model Diagnostics**

Three types of uncertainty are presented for the assessment of Widow Rockfish. First, uncertainty in the parameter estimates was determined using approximate asymptotic estimates of the standard error. These estimates were based on the maximum likelihood theory that the inverse of the Hessian matrix (the second derivative of the log-likelihood

function with respect to the parameter vector) approaches the true uncertainty of the parameter estimates as the sample size approaches infinity. This approach takes into account the uncertainty in the data and supplies correlation estimates between parameters, but does not capture possible skewness in the error distribution of the parameters and may not accurately estimate the standard error in some cases (see Stewart et al. 2013) UPDATE REF!!!!!!.

The second type of uncertainty that is presented is related to modeling and structural error. This uncertainty cannot be captured in the base model as it is related to errors in the assumptions used in specifying the base model. Therefore, sensitivity analyses were conducted where assumptions were modified to reveal the effect they have on the model results.

Lastly, a major axis of uncertainty was determined from a parameter or structural assumption that results in the greatest change in stock status and advice, and projections were made for different states of nature based upon that parameter or structural assumption.

### **2.8.1 Convergence**

### **2.8.2 Parameter Uncertainty**

Parameter estimates are shown in Table 22, Table 23, and Table 28 along with approximate asymptotic standard errors. The only parameters with an absolute value of correlation greater than 0.95 were the female and male natural mortality parameters, which is expected. Estimates of key derived quantities are given in Table 26 along with approximate 95% asymptotic confidence intervals. There is a reasonable amount of uncertainty in the estimates of biomass. The confidence interval of the 2019 estimate of depletion is XX%–XX% and above the management target of 40% of the unfished spawning biomass.

### **2.8.3 Sensitivity Analyses**

Sensitivity analysis was performed to determine the model behavior under different assumptions than those of the base case model. Seven sensitivity analyses were conducted to explore the potential differences in model structure and assumptions, including:

1. Fixed natural mortality at 0.1 for both sexes (2015 assessment prior)
2. Fixed natural mortality at 0.124 yr<sup>-1</sup> for females and 0.129 yr<sup>-1</sup> for males (2011 assessment prior)
3. Fixed steepness at 0.4



4. Fixed steepness at 0.6
5. Fixed steepness at 0.798 (2015 assessment value)
6. Forcing asymptotic selectivity on the midwater trawl fleet
7. Fitting logistic curves for NWFSC WCGBT survey selectivities
8. Weighting the composition data using the Francis method
9. Updated Washington catch reconstruction
10. Inclusion of previously excluded shrimp trawl data
11. Exclusion of triennial survey data

Likelihood values and estimates of key parameters are shown in Table 29. Predicted spawning biomass trajectories and estimated recruitment deviations are shown in Figure 65. The estimates of current stock depletion ranged from 55%-142% across the sensitivity runs, with fixing natural mortality at 0.1 resulting in the lowest estimate and forcing asymptotic selectivity on the midwater trawl fleet resulting in the highest estimate. Generally, the trajectory of the spawning biomass was qualitatively similar across all tested models, e.g., peak around late 1970s and late 2010s, projected decrease in biomass in 2025 followed by some recovery into the 2030s; the quantitative magnitude of these trends did vary across cases.

Fixing  $M$  at values lower than the base case estimate resulted in decreases in estimated spawning biomass, while fixing steepness across the values tested resulted in similar or increased estimated spawning biomass. The relative spawning biomass in 2025 changed to 93% with an  $M$  of 0.124 yr<sup>-1</sup> and 0.129 yr<sup>-1</sup> for females and males, respectively, and to 55% with an  $M$  of 0.1 yr<sup>-1</sup>. Fixing steepness at a value of 0.6 resulted in an increase of the spawning biomass to 125% and a decrease in equilibrium yield at a SPR50% reference harvest rate, while other tested values for fixed steepness had relatively minimal effects on spawning biomass and equilibrium yield at a SPR50% reference harvest rate. Fixing steepness at a value of 0.4 resulted in low recruitment deviations in the 2019-2024 period relative to other tested models and the base model.

Forcing asymptotic selectivity on the midwater fleet resulted in the largest estimated spawning biomass for 2025, while forcing logistic selectivity on the NWFSC WCGBT resulted in similar estimated spawning biomass to the base model. Including shrimp trawl data and updating WA catch reconstruction had almost no impact on the estimated spawning biomass. Excluding the triennial survey data lead to slight increases in estimated spawning biomass.

The alternative weighting using the Francis method generally increased the estimate of spawning biomass across the timeseries, but the estimated biomass for 2025 was similar between the Francis weighted model and the base model.

#### 2.8.4 Retrospective Analysis

A 5-year retrospective analysis was conducted by running the model using data only through 2020, 2021, 2022, 2023, and 2024 progressively (Table 30 and Figure 66). The initial scale of the spawning population was basically unchanged for all of these retrospectives. Removing 4–5 years of data led to slightly lower estimates of fishing mortality (F) and slightly higher spawning biomass over the last 15 years. In contrast, removing only 1–2 years resulted in higher F and lower biomass estimates. Despite these minor differences, all retrospectives showed a consistent declining trend in spawning biomass over the past decade. No concerning patterns were observed in the retrospective analysis.

#### 2.8.5 Likelihood Profiles and key parameters

Likelihood profiles were conducted for  $R_0$ , steepness (even though it was not estimated in the base case) and over male and female natural mortality values simultaneously. These likelihood profiles were conducted by fixing the parameter at specific values and removing the prior on the parameter being profiled. Without the original prior distribution the MLE estimates from the base case will likely be different than the MLE in the likelihood profile, but this displays what information the data have. There was some difficulty in achieving model convergence for many parameterizations in the likelihood profile. In some cases jittering was required.

As  $R_0$  increased, natural mortality also increased and the relative spawning biomass in 2015 was less depleted (Table 31). There was variable support for each likelihood component across the range of  $R_0$  evaluated. The total likelihood supported the estimated value (Table 31). Profiles are illustrated in Figure 68.

### 2.9 Unresolved Problems and Major Uncertainties

## 3 Management

### 3.1 Reference Points

catches (landings plus discards) have been below the point estimate of potential long-term yields calculated using an SPR50% reference point and the population has been increasing over the last decade. However, catches in 2018 were above the point estimate of potential long-term yields calculated using an SPR50 reference point.

The predicted spawning biomass from the base model generally showed a slight decline until the late 1970s, steep increase above unfished equilibrium levels, then a steep decline until the mid-1980s followed by less of a decline until 2001 (Figure 61). Since 2001, the spawning biomass has been increasing due to small catches, and recently, above average recruitment. The 2018 spawning biomass relative to unfished equilibrium spawning biomass is above the target of 40% of unfished spawning biomass (Figure 63). The fishing intensity (relative 1-SPR) exceeded the current estimates of the harvest rate limit (SPR50%) throughout the 1980s and early 1990s, as seen in Figure 73. Recent exploitation rates on Widow Rockfish were predicted to be much less than target levels. In recent years, the stock has experienced exploitation rates that have been below the target level while the biomass level has remained above the target level (Figure 74).

The equilibrium yield plot is shown in Figure 75, based on a steepness value fixed at 0.720. The predicted maximum sustainable yield under the assumptions of this assessment occurs near 25% of equilibrium unfished spawning biomass.

### 3.2 Unresolved problems and major uncertainties

### 3.3 Harvest Projections and Decision Tables

```
load(file=here::here("report","tables","projections.rda"))
projections <- as.data.frame(projections$table)

projections <- projections|>dplyr::mutate(across(2:ncol(projections), ~ round(.x)))
```

Projections with catches based on the predicted annual catch limit (ACL) using the SPR rate of 50%, the 40:10 control rule, and a 0.25 P\* adjustment using a sigma of 0.50 from 2021 onward suggest that the spawning biomass will decrease over the projection period

for all states of nature. Predicted ACL catches range from 10,961 mt in 2021 to 5,944 mt in 2030.

### ***nOT SURE WHERE acl CATCHES ARE***

#### **3.4 Evaluation of Scientific Uncertainty**

#### **3.5 Regional management considerations**

Widow Rockfish have shown latitudinal differences in life-history parameters, which has led past assessment authors to pursue a two-area model. Modelling a stock with two areas is difficult because it requires many assumptions about recruitment distribution, movement, and connectivity, while also splitting data into two areas that reduces sample sizes when compared to a coastwide model. The upside is that it can result in a better model that more accurately predicts regional status. This assessment is a coastwide model because not enough is known about the assumptions that would have to be made for a two-area model.

It is still important to consider regional differences when making management decisions. Following recent cohorts through time with survey data showed that older fish showed up in the north after younger fish were observed in the south (Figure 2). This may indicate connectivity between the north and the south and that this is truly one stock. However, more investigation is needed.

Widow Rockfish are managed on a coastwide basis and observed more often in the NWFSC WCGBT bottom trawl survey north of latitude 40° 10' N. Bottom trawl catches in California have historically been as large as in Oregon and larger than in Washington, but recently catches in California have been small. Rockfish Conservation Areas (RCAs) cover a significant proportion of Widow Rockfish habitat, but a midwater trawl fishery is beginning to re-develop that can fish in these areas. Future assessments and management of Widow Rockfish may want to monitor where catches are being taken to make sure that specific areas are not being overexploited. In addition, research on the connectivity along the coast as well as regional differences would help to inform the potential for overfishing specific areas. »»»> 1ed44512b0ba062144bae79bac8c50155d6b6fec

#### **3.6 Research and Data Needs**

### **3.7 Acknowledgements**

**3.8 References**

## 4 Tables

### 4.1 Data

#### 4.1.1 Fishery-dependent data

...

::: {#tbl-hake\_removals .cell tbl-cap='Landings (mt) from the foreign & domestic at-sea fleet and the domestic shoreside hake fleet. Catches (mt) from the Pacific whiting at-sea fishery as determined by onboard observers.'} ::: {.cell-output-display}

Year	For- eign...Do- mestic	Shore- side.hake	X	X.1
	At-sea	CA	OR	WA
1966	3670	0	0	0
1967	3902	0	0	0
1968	1956	0	0	0
1969	358	0	0	0
1970	554	0	0	0
1971	701	0	0	0
1972	421	0	0	0
1973	656	0	0	0
1974	418	0	0	0
1975	391.2	0	0	0
1976	718.5	0	0	0
1977	119.3	0	0	0
1978	191.9	0	0	0
1979	197.9	0	0	0
1980	272	0	0	0
1981	227.9	0	0	0
1982	157.5	0	0	0
1983	131.5	0	0	0
1984	294.7	0	0	0
1985	182.6	0	0	0
1986	256.8	0	0	0
1987	181.3	0	0	0
1988	231.6	0	0	0



Year	For- eign...Do- mestic	Shore- side.hake	X	X.1
1989	212	0	0	0
1990	230.2	0	0	0
1991	471.3	42.7	39	9.3
1992	389.6	13.5	42.1	6.2
1993	173.2	0.4	91.2	11
1994	370.7	2.1	210.8	28.6
1995	228.6	7.2	192.1	36.8
1996	252.2	5.7	475.1	104.7
1997	215.5	7.2	133.9	22.1
1998	268.5	40.4	278	28.1
1999	191.8	12.7	166.4	15.2
2000	205.4	7.7	70.9	4.7
2001	174	9.2	26.4	9
2002	154.9	1.2	2.6	1.4
2003	14.5	0.4	7.6	4.6
2004	21.2	7.4	12.4	8.5
2005	80.1	5.2	59.1	13.6
2006	143	3.6	11.3	35.3
2007	146	1	46.1	35.3
2008	115.2	29.2	36.1	37.5
2009	26.6	2.3	46.6	59.8
2010	44.6	9	35.3	17.5
2011	38.4	0	79.9	19.5
2012	79.2	0	85.1	17.1
2013	31.2	0	115.1	29.2
2014	56.2	0	250.1	35.9

∴ ∴

∴

Table 3: A subset of management actions of importance to fisheries that caught Widow Rockfish.

Year	Management action
1982	Establishment of a 75,000 pound trip limit on Widow Rockfish in October
1983	Per-trip and per-week limits implemented for Sebastes complex coastwide (north and south of 40° N)
1984	30,000 pound Widow Rockfish trip limit at the start of the year adjusted to 1,000 pound trip limit in September
	50,000 pound Widow Rockfish trip limit limited to once per week
	Trip limit lowered to 40,000 pounds once per week in May
	Directed fishery for Widow Rockfish closed in August and a full fishery closure in November
1985	30,000 pound trip limit once per week, or 60,000 pounds once every 2 weeks. Every 2 week option was rescinded in April
	Landings of Sebastes complex and Widow Rockfish smaller than 3,000 pounds unrestricted
	Widow Rockfish trip limit reduced to 3,000 pounds per trip without a trip frequency in July
1986	30,000 pound coastwide Widow Rockfish trip limit with no biweekly option
	Landings of Sebastes complex and Widow Rockfish smaller than 3,000 pounds unrestricted
	3,000 pound coastwide trip limit implemented in September when Widow Rockfish ABC reached
1987	30,000 pound coastwide Widow Rockfish trip limit with no biweekly option. Only one landing per week above 3,000 pounds.
	Reduced Widow Rockfish trip limit to 5,000 pounds in October
	Closed the Widow Rockfish fishery in November
1988	30,000 pound coastwide Widow Rockfish trip limit with no biweekly option. Only one landing per week above 3,000 pounds.
	Reduced Widow Rockfish trip limit to 3,000 pounds in October
	30,000 pound coastwide Widow Rockfish trip limit with no biweekly option. Only one landing per week above 3,000 pounds.

Year	Management action
1989	Reduced Widow Rockfish trip limit to 10,000 pounds in April
	Reduced Widow Rockfish trip limit to 3,000 pounds in October
1990	15,000 pound trip limit once per week, or 25,000 pounds once every 2 weeks. Only one landing per week above 3,000 pounds.
	Closed the Widow Rockfish fishery in December
1991	10,000 pound trip limit once per week, or 20,000 pounds once every 2 weeks. Only one landing per period above 3,000 pounds.
	Reduced Widow Rockfish trip limit to 3,000 pounds on my birthday in September
1992	30,000 pound coastwide Widow Rockfish trip limit per 4-week period. All landings apply to the 30,000 pounds.
	Reduced Widow Rockfish trip limit to 3,000 pounds in August
	Re-established the 30,000 pound cumulative landing limit for December
1993	30,000 pound coastwide Widow Rockfish trip limit per 4-week period. All landings apply to the 30,000 pounds.
	Reduced Widow Rockfish trip limit to 3,000 pounds in December
1994	Divided the commercial groundfish fishery in limited entry and open access fisheries.
	30,000 pound cumulative Widow Rockfish limit per calendar month.
	Reduced Widow Rockfish trip limit to 3,000 pounds in December
	Rockfish limit of 10,000 per vessel per trip in open access fisheries, not to exceed 30,000 pounds of Widow Rockfish (as in limited entry fisheries) cumulative per month.
1995	30,000 pound cumulative Widow Rockfish limit per calendar month.
	Monthly cumulative trip limit increased to 45,000 pounds for Widow Rockfish
1996	70,000 pound cumulative Widow Rockfish limit per two-month period.
	Reduced cumulative two-month period Widow Rockfish limit to 40,000 pounds in September.
	25,000 pound monthly cumulative limit implemented in November.
1997	70,000 pound cumulative Widow Rockfish limit per two-month period.
	Reduced cumulative two-month period Widow Rockfish limit to 60,000 pounds in May.

Year	Management action
1998	25,000 pound cumulative Widow Rockfish limit per two-month period.
	Increased cumulative two-month period Widow Rockfish limit to 30,000 pounds in May.
	Open access monthly cumulative trip limits reduced to 3,000 pounds in July.
1999	Dividing line between north and south management areas moved to 40° 10' N.
	Three-phase cumulative limit period system introduced.
	Phase 1: 70,000 pounds cumulative limit from January through March for Widow Rockfish.
	Phase 2: 16,000 pounds per 2-month period April through September for Widow Rockfish.
	Phase 3: 30,000 pounds per month October through December for Widow Rockfish.
	Open access limit to 2,000 pounds per month of Widow Rockfish
	Phase 2 two-month limits reduced to 11,000 pounds for Widow Rockfish starting in June.
	Open access month cumulative trip limit increased to 8,000 pounds of Widow Rockfish.
2000	WA and OR restrict landings applied to 30,000 monthly limit to have midwater gear. State imposed cumulative trip limit per month applied otherwise.
	Sorting of Widow Rockfish required before weighing in limited entry and open access fisheries.
	New limited entry trawl gear restrictions implemented for large footrope trawl gear, small footrope trawl gear, and midwater trawl gear.
	Cumulative trip limits allowed for Widow Rockfish only if small footrope or midwater trawl gear were used. Higher cumulative trip limits available to midwater gear.
	30,000 pound two-month cumulative trip limit for Widow Rockfish caught with mid-water gear.
	1,000 pound monthly trip limit allowed for small footrope trawl.
	3,000 pound monthly trip limits for Widow Rockfish caught with limited entry fixed gear, open access gear, and exempted trawl gear. Some closures south of 40° 10' N latitude in January through April

Year	Management action
2001	Similar actions as in 2000 with the following changes
	20,000 pound two-month cumulative trip limit for Widow Rockfish caught with mid-water gear in January through April and September through October. 10,000 pound two-month cumulative trip limit in other periods.
	Widow Rockfish limits reduced to 1,000 pounds per month in July-September unless landed with Pacific Whiting, which is 2,000 pounds per month with a 500 pound trip limit.
	Retention of Widow Rockfish prohibited beginning in October. For gears other than midwater trawl.
2002	Rockfish Conservation Areas (RCA) established. Large footrope gear prohibited inside 275 m. Widow fishery closed most of the year except for a small amount of bycatch and small monthly limits in some months.
2003	Widow fishery closed most of the year except for a small amount of bycatch and small monthly limits in some months.
2004	Widow fishery closed most of the year except for a small amount of bycatch and small monthly limits in some months
2005	Widow fishery closed most of the year except for a small amount of bycatch and small monthly limits in some months.
2006	Amendment 19 established essential fish habitat (EFH) boundaries and conservation areas. Widow bycatch cap in the non-tribal limited entry whiting trawl fishery increased from 200 mt to 220 mt in October
2007	Seasonal changes of trawl RCA boundaries and periodic closures within certain latitude boundaries (e.g., north of Cape Alava at 48°10' N latitude to the U.S.-Canada border) started in 2007. Small monthly limits for Widow Rockfish (less than 1,500 pounds per month)
	Widow bycatch cap in the non-tribal limited entry whiting trawl fishery increased from 200 mt to 220 mt in May.
	Limited entry whiting trawl fishery closed due to attainment of 220 mt widow bycatch in July
	Limited entry whiting trawl fishery re-opened with 275 mt widow bycatch cap in October
	Widow bycatch cap of 275 mt adopted for limited entry whiting trawl fishery.
	Limited entry whiting trawl fishery closed due to attainment of canary bycatch in August

Year	Management action
2008	Limited entry whiting trawl fishery re-opened with 284 mt widow bycatch cap in October
	Small monthly limits for Widow Rockfish (less than 1,500 pounds per month)
2009	Sector specific bycatch caps for Widow Rockfish in the limited entry whiting trawl fishery: 105 mt for shoreside fleet, 85 mt to catcher-processors, 60 mt to motherships
	Small monthly limits for Widow Rockfish (less than 1,500 pounds per month)
2010	Trawl rationalization began, establishing the IFQ fishery.
2011	Small monthly limits for Widow Rockfish (less than 1,500 pounds per month)

...

```

      Years      Depth_range_m Southern_latitude
"character"    "character"      "numeric"

```

```
[1] "data.frame"
```

Table 4: Depth ranges and limits of the southern latitude in the Triennial survey for the different years.

Years	Depth range (m)	Southern latitude
1977	91-457	34.05
1980-1986	55-366	36.80
1989-1992	55-366	34.50
1995-2004	55-500	34.50

Table 5: Stratifications used for the two surveys.

Triennial					
Strata	Area (km2)	Depth1	Depth2	Latitude1	Latitude2
A	33,730.25	55	183	34.5	49
B	11,062.63	183	400	34.5	49

Table 6: Stratifications used for the two surveys.

NWFSC WCGBT					
Strata	Area (km2)	Depth1	Depth2	Latitude1	Latitude2
A	10,687.86	55	183	34.5	40.5
B	3,394.82	183	400	34.5	40.5
C	23,042.39	55	183	40.5	49.0
D	7,667.81	183	400	40.5	49.0



## 4.2 Figures

- Adams, PB. 1987. "The Diet of Widow Rockfish *Sebastes Entomelas* in Northern California." *NOAA Tech. Rep. NMFS* 48: 37–41.
- Bradburn, Mark James, Aimee A Keller, and Beth Helene Horness. 2011. "The 2003 to 2008 US West Coast Bottom Trawl Surveys of Groundfish Resources Off Washington, Oregon, and California: Estimates of Distribution, Abundance, Length, and Age Composition."
- Dick, Edward Joseph. 2009. *Modeling the Reproductive Potential of Rockfishes (Sebastes Spp.)*. University of California, Santa Cruz.
- Douglas, David A. 1998. "Species Composition of Rockfish in Catches by Oregon Trawlers, 1963-93."
- Hamel, Owen S. 2014. "A Method for Calculating a Meta-Analytical Prior for the Natural Mortality Rate Using Multiple Life History Correlates." *ICES Journal of Marine Science* 72 (1): 62–69.
- He, Xi, Donald Pearson, E. J. Dick, John Field, Stephen Ralston, and Alec MacCall. 2011. "Status of the Widow Rockfish Resource in 2011." Portland, OR: Pacific Fishery Management Council.
- Jones, Walter G, and George Y Harry Jr. 1960. "The Oregon Trawl Fishery for Mink Food 1948-1957." *Fish Commission of Oregon Research Briefs* 8: 14–30.
- Punt, André E, David C Smith, Kyne KrusicGolub, and Simon Robertson. 2008. "Quantifying Age-Reading Error for Use in Fisheries Stock Assessments, with Application to Species in Australia's Southern and Eastern Scalefish and Shark Fishery." *Canadian Journal of Fisheries and Aquatic Sciences* 65 (9): 1991–2005.
- Rogers, Jean Beyer. 2003. "Species Allocation of *Sebastes* and *Sebastolobus* Sp. Caught by Foreign Countries from 1965 Through 1976 Off Washington, Oregon, and California, USA."
- Stanley, RD, R Kieser, K Cooke, AM Surry, and B Mose. 2000. "Estimation of a Widow Rockfish (*Sebastes Entomelas*) Shoal Off British Columbia, Canada as a Joint Exercise Between Stock Assessment Staff and the Fishing Industry." *ICES Journal of Marine Science* 57 (4): 1035–49.