



B

Facultat de Biologia Diagonal, 645 08028 Barcelona

Unitat 5. Models de probabilitat més frequents. SOLUCIONS

- 1. $X \sim \text{Bernouilli}(p=2/3)$. E(X)=2/3 i Var(X)=2/9.
- 2. 0.22.
- 3. 0.01966.
- 4. 0.0000059.
- 5. (a) 0.0159. (b) 0.3585. (c) 9.5*10⁻²⁷. (d) 19.
- 6. Si $p \le 1/3$ en cuadrimotor, però si p > 1/3 en bimotor.
- 7. 0.026.
- 8. 0.9604.
- 9. (a) 0.105. (b) 0.4219.
- 10. (a) 0.048. (b) 0.043. (c) $E(X_{(a)}) = 5.67$ i $E(X_{(b)}) = 17$.
- 11. 0.016.
- 12. (a) 0.024. (b) 0.976.
- 13. (a) 0.406. (b) 0.01655.
- 14. (a) $X \sim Poiss(2)$. (b) $X_3 \sim Poiss(6)$. (c) $X_{1/2} \sim Poiss(1)$. (d) $P(X \le 2) = 0.6767$, $P(X_3 \ge 5) = 0.7373$, $P(X_{1/2} = 0) = 0.3679$.
- 15. (a) E(X)=6, $X \sim Poiss(6)$. (b) 0.017.
- 16. 0.1353.
- 17. 0.645.
- 18. (a) 2/3. (b) 1/3.
- 19.0.29.