INTRO TO DATA SCIENCE LINEAR REGRESSION

I. LINEAR REGRESSION II. POLYNOMIAL REGRESSION

EXERCISES: III. LINEAR REGRESSION IN R IV. PREDICTING BASEBALL SALARIES

I. LINEAR REGRESSION

	continuous	categorical
supervised	???	???
unsupervised	???	???

REGRESSION PROBLEMS

supervised
unsupervisedregression
dimension reductionclassification
clustering

Q: What is a **regression** model?

Q: What is a **regression** model?

A: A functional relationship between input & response variables.

Q: What is a **regression** model?

A: A functional relationship between input & response variables

The **simple linear regression** model captures a linear relationship between a single input variable x and a response variable y:

Q: What is a **regression** model?

A: A functional relationship between input & response variables

The **simple linear regression** model captures a linear relationship between a single input variable x and a response variable y:

$$y = \alpha + \beta x + \varepsilon$$

Q: What do the terms in this model mean?

$$y = \alpha + \beta x + \varepsilon$$

$$y = \alpha + \beta x + \varepsilon$$

A: y =**response variable** (the one we want to predict)

$$y = \alpha + \beta x + \varepsilon$$

A: y =**response variable** (the one we want to predict)

x =input variable (the one we use to train the model)

$$y = \alpha + \beta x + \varepsilon$$

A: y =**response variable** (the one we want to predict)

x =input variable (the one we use to train the model)

 α = intercept (where the line crosses the y-axis)

$$y = \alpha + \beta x + \varepsilon$$

A: y =**response variable** (the one we want to predict)

x =input variable (the one we use to train the model)

 α = intercept (where the line crosses the y-axis)

 β = regression coefficient (the model "parameter")

$$y = \alpha + \beta x + \varepsilon$$

A: y =**response variable** (the one we want to predict)

x =input variable (the one we use to train the model)

 α = intercept (where the line crosses the y-axis)

 β = regression coefficient (the model "parameter")

 ε = **residual** (the prediction error)


We can extend this model to several input variables, giving us the multiple linear regression model:

We can extend this model to several input variables, giving us the **multiple linear regression** model:

$$y = \alpha + \beta_1 x_1 + \dots + \beta_n x_n + \varepsilon$$

Linear regression involves several technical assumptions and is often presented with lots of mathematical formality.

The details are not very important for our purposes, but you can check them out if you're interested.



Statistical Models Theory and Practice REVISED EDITION David A. Freedman

Q: How do we fit a regression model to a dataset?

Q: How do we fit a regression model to a dataset?

A: Minimize the sum of the squared residuals (Ordinary Least Squares).

Q: How do we fit a regression model to a dataset?

A: Minimize the sum of the squared residuals (Ordinary Least Squares).

In practice, any respectable piece of software will do this for you.

Q: How do we fit a regression model to a dataset?

A: Minimize the sum of the squared residuals (Ordinary Least Squares).

In practice, any respectable piece of software will do this for you.

And there are other ways.

Q: How do we fit a regression model to a dataset?

A: Minimize the sum of the squared residuals (Ordinary Least Squares).

In practice, any respectable piece of software will do this for you.

And there are other ways.

And software implements them as well.

II: POLYNOMIAL REGRESSION

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

A: Yes, because it's linear in the β 's!

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \varepsilon$$

Q: This represents a nonlinear relationship. Is it still a linear model?

A: Yes, because it's linear in the β 's!

"Although polynomial regression fits a *nonlinear* model to the data, as a statistical estimation problem it is *linear*, in the sense that the regression function E(y|x) is linear in the unknown parameters that are estimated from the data. For this reason, polynomial regression is considered to be a special case of multiple linear regression." — Wikipedia

POLYNOMIAL REGRESSION

Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

Polynomial regression allows us to fit very complex curves to data.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

But there is a problem with the model we've written down so far.

POLYNOMIAL REGRESSION



This model displays **collinearity**, which means the predictor variables are highly correlated with each other.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

```
> x <- seq(1, 10, 0.1)
> cor(x^9, x^10)
[1] 0.9987608
```

This model displays **collinearity**, which means the predictor variables are highly correlated with each other.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n x^n + \varepsilon$$

Collinearity causes the linear regression model to "break down", because it can't tell the predictor variables apart.

This model displays **collinearity**, which means the predictor variables are highly correlated with each other.

$$y = \alpha + \beta_1 x + \beta_2 x^2 + \dots + \beta_n$$

For identical features, this results in a singularity. We will see an example of this in just a minute!

Collinearity causes the linear regression model to "break down", because it can't tell the predictor variables apart.

POLYNOMIAL REGRESSION

Q: What can we do about this?

POLYNOMIAL REGRESSION

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

$$y = \alpha + \beta_1 f_1(x) + \beta_2 f_2(x^2) + ... + \beta_n f_n(x^n) + \varepsilon$$

Q: What can we do about this?

A: Replace the correlated predictors with uncorrelated predictors.

$$y = \alpha + \beta_1 f_1(x) + \beta_2 f_2(x^2) + \dots + \beta_n f_n(x^n) + \varepsilon$$

OPTIONAL NOTE

These polynomial functions form an *orthogonal basis* of the function space.

Q: So, how do we select a linear regression model?

- Q: So, how do we select a linear regression model?
- Techniques for producing designs

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection
 - Backward elimination

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection
 - Backward elimination
 - More

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection
 - Backward elimination
 - More
- Techniques for evaluating designs

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection
 - Backward elimination
 - More
- Techniques for evaluating designs
 - Training metrics

- Q: So, how do we select a linear regression model?
- Techniques for producing designs
 - Theory
 - Forward selection
 - Backward elimination
 - More
- Techniques for evaluating designs
 - Training metrics
 - Test set performance / Cross-validation

INTRO TO DATA SCIENCE

EXERCISES