

P08 Distinct Distance Route V1

Laureano Arcanio (February 2026)

Problem

Let $N = 75,001$. There are $2N - 1$ points placed on a line at integer positions

$$1, 2, \dots, 2N - 1.$$

A route is any permutation $(t_1, t_2, \dots, t_{2N-1})$ of these points.

Assume that the consecutive distances

$$|t_1 - t_2|, |t_2 - t_3|, \dots, |t_{2N-2} - t_{2N-1}|$$

are all distinct.

Let M be the minimum possible value of $\max(t_1, t_{2N-1})$ over all such routes.

Determine M .

Solution

1. The distance multiset is forced

There are $2N - 2$ consecutive distances. Each $|t_i - t_{i+1}|$ is a positive integer and

$$|t_i - t_{i+1}| \leq (2N - 1) - 1 = 2N - 2.$$

Since the $2N - 2$ distances are pairwise distinct, they must be exactly

$$\{1, 2, \dots, 2N - 2\}.$$

Hence the total distance is forced:

$$\begin{aligned} \sum_{i=1}^{2N-2} |t_i - t_{i+1}| &= 1 + 2 + \dots + (2N - 2) \\ &= \frac{(2N - 2)(2N - 1)}{2} \\ &= (N - 1)(2N - 1). \end{aligned} \tag{1}$$

2. Potential inequality and an endpoint-sum lower bound

Define

$$s(x) = |x - N|.$$

For all integers a, b , the triangle inequality gives

$$\begin{aligned} |a - b| &= |(a - N) - (b - N)| \\ &\leq |a - N| + |b - N| \\ &= s(a) + s(b). \end{aligned} \tag{2}$$

Summing (2) over all consecutive pairs:

$$\sum_{i=1}^{2N-2} |t_i - t_{i+1}| \leq \sum_{i=1}^{2N-2} (s(t_i) + s(t_{i+1})) = 2 \sum_{j=1}^{2N-1} s(t_j) - (s(t_1) + s(t_{2N-1})). \tag{3}$$

Because (t_j) is a permutation of $\{1, 2, \dots, 2N-1\}$, the multiset of scores is

$$\{0, 1, 1, 2, 2, \dots, N-1, N-1\},$$

so

$$\begin{aligned} \sum_{j=1}^{2N-1} s(t_j) &= 2(1 + 2 + \dots + (N-1)) \\ &= 2 \cdot \frac{(N-1)N}{2} \\ &= N(N-1). \end{aligned} \tag{4}$$

Substituting (1) and (4) into (3) yields

$$(N-1)(2N-1) \leq 2N(N-1) - (s(t_1) + s(t_{2N-1})),$$

hence

$$s(t_1) + s(t_{2N-1}) \leq N-1. \tag{5}$$

Since $N - x \leq |x - N| = s(x)$ for all x , we obtain

$$(N - t_1) + (N - t_{2N-1}) \leq N-1 \quad \Rightarrow \quad t_1 + t_{2N-1} \geq N+1. \tag{6}$$

3. Parity constraint

Modulo 2, we have $|x - y| \equiv x - y \equiv x + y \pmod{2}$. Therefore,

$$\sum_{i=1}^{2N-2} |t_i - t_{i+1}| \equiv \sum_{i=1}^{2N-2} (t_i + t_{i+1}) \equiv t_1 + t_{2N-1} \pmod{2},$$

since every interior t_2, \dots, t_{2N-2} appears twice.

Using (1),

$$t_1 + t_{2N-1} \equiv (N-1)(2N-1) \pmod{2}. \tag{7}$$

Because $N = 75,001$ is odd, $N-1$ is even, so $(N-1)(2N-1)$ is even and hence

$$t_1 + t_{2N-1} \equiv 0 \pmod{2}. \tag{8}$$

4. Impossibility below 37,502

Here $N + 1 = 75,002$. From (6),

$$t_1 + t_{2N-1} \geq 75,002. \quad (9)$$

If $\max(t_1, t_{2N-1}) \leq 37,501$, then $t_1 + t_{2N-1} \leq 75,002$. So equality holds and forces $t_1 = t_{2N-1} = 37,501$, impossible in a permutation. Therefore

$$\max(t_1, t_{2N-1}) \geq 37,502. \quad (10)$$

Answer

$$\boxed{37,502}$$